Characteristics	total population (n=4602)
Mean (SD) age, years	78.3 (9.3)
Women	2085 (45.3)
Mean (SD) CHADS ₂ score	2.1 (1.2)
Mean (SD) CHA ₂ DS ₂ -VASc score	3.6 (1.5)
Mean (SD) HAS-BLED score	2.6 (10)
Antithrombotic treatment the day before the inclusion event:	
Single: oral anticoagulation	1101 (23.9)
Single: antiplatelets	2450 (53.3)
Dual: oral anticoagulation+antiplatelets	893 (19.4)
Dual: aspirin+adenosine diphosphate receptor antagonists	117 (2.5)
Triple: oral anticoagulation+aspirin+adenosine diphosphate receptor antagonists	41 (0.9)
Concomitant drugs:	
Dipyridamole (persantin)	199 (4.3)
Heparin	20 (0.4)
Non-steroidal anti-inflammatory drug	1126 (24.5)
Proton pump inhibitor	693 (15.1)
H ₂ receptor antagonist	121 (2.6)
Comorbidities:	
Stroke or thromboembolism	1034 (22.5)
Myocardial infarction	681 (14.8)
Ischaemic heart disease	1749 (38.0)
Peripheral arterial disease	290 (6.3)
Vascular disease	903 (19.6)
Heart failure	1411 (30.7)
Hypertension	2058 (44.7)
Diabetes	743 (16.2)
Chronic kidney disease	304 (6.6)
Liverfailure	64 (14)
Previous bleeding	635 (13.8)
Alcohol misuse	219 (4.8)
Gastroesophageal reflux	45 (10)
Gastric or duodenal ulcer	302 (6.6)
Gastritis	37 (0.8)
Invasive and surgical procedures:	
Gastrointestinal surgery	935 (20.3)
Gastroscopy	1074 (23.3)