

Succ contraindications:

Burn/trauma/crush

Hyperkalemia

Myopathies

H/o spinal cord damage

Malignant hyperthermia

NEXUS:

No midline tenderness


No focal neuro deficit

No distracting injury

No decreased LOC

No intoxication

| BEHAVIOUR            | RESPONSE                |   |
|----------------------|-------------------------|---|
| Best Eye Response    | Spontaneously           | 4 |
|                      | To speech               | 3 |
|                      | To pain                 | 2 |
|                      | No response             | 1 |
| Best Verbal Response | Oriented                | 5 |
|                      | Confused                | 4 |
|                      | Inappropriate words     | 3 |
|                      | Inappropriate sounds    | 2 |
|                      | No response             | 1 |
| Best Motor Response  | Obeys commands          | 6 |
|                      | Moves to localized pain | 5 |
|                      | withdrawal from pain    | 4 |
|                      | Abnormal flexion        | 3 |
|                      | Abnormal extension      | 2 |
|                      | No response             | 1 |



- ☐ Physio Issues (HOp) Considered
- ☐ Induction Agent/Muscle Relaxant
- ☐ ± Push-Dose Epi
- ☐ Failed Plan Verbalized
- ☐ **Cric Evaluation**
- ☐ Post-Tube Analgesia/Sedation

- ☐ Denitrogenated  $\geq 3$  minutes
- ☐ **ApOx with NC @15**
- ☐ **Oxygenated  $\geq 95\%$  ( $\pm$  CPAP)**
- ☐ Look in Mouth · Dentures · Range Neck
- ☐ **Positioning**
- ☐ Pulse Ox Visible or Audible
- ☐ Access - Reliable & Tested

- ☐ BVM ( $\pm$  PEEP Valve) on Oxygen
- ☐ Video Laryngoscope
- ☐ Backup Laryngoscope
- ☐ **OPA, Bougie, SGA, Scalpel**
- ☐ Suction x 2