

Succ contraindications:

Burn/trauma/crush

Hyperkalemia

Myopathies

H/o spinal cord damage

Malignant hyperthermia

NEXUS:

No midline tenderness

No focal neuro deficit

No distracting injury

No decreased LOC

No intoxication

BEHAVIOUR	RESPONSE	
Best Eye Response	Spontaneously	4
	To speech	3
	To pain	2
	No response	1
Best Verbal Response	Oriented	5
	Confused	4
	Inappropriate words	3
	Inappropriate sounds	2
	No response	1
Best Motor Response	Obeys commands	6
	Moves to localized pain	5
	withdrawal from pain	4
	Abnormal flexion	3
	Abnormal extension	2
	No response	1



- ☐ Physio Issues (HOp) Considered
- ☐ Induction Agent/Muscle Relaxant
- ☐ ± Push-Dose Epi
- ☐ Failed Plan Verbalized
- ☐ **Cric Evaluation**
- ☐ Post-Tube Analgesia/Sedation

- ☐ Denitrogenated  $\geq 3$  minutes
- ☐ **ApOx with NC @15**
- ☐ **Oxygenated  $\geq 95\%$  ( $\pm$  CPAP)**
- ☐ Look in Mouth · Dentures · Range Neck
- ☐ **Positioning**
- ☐ Pulse Ox Visible or Audible
- ☐ Access - Reliable & Tested

- ☐ BVM ( $\pm$  PEEP Valve) on Oxygen
- ☐ Video Laryngoscope
- ☐ Backup Laryngoscope
- ☐ **OPA, Bougie, SGA, Scalpel**
- ☐ Suction x 2