

# HAOT: A Python package for hypersonic aero-optics analysis

Martin E. Liza <sup>1</sup>

<sup>1</sup> The University of Arizona

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

## Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Open Journals](#) 

## Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

## License

Authors of papers retain copyright<sup>14</sup>  
and release the work under a  
Creative Commons Attribution 4.0<sup>15</sup>  
International License ([CC BY 4.0](#))<sup>16</sup>.

## Summary

Hypersonic flows present a unique challenges due to the complex interplay of fluid dynamics, chemical reactions, and optical phenomena. As a signal from a Light Detection and Ranging (LiDAR) travels through a hypersonic flow field, the beam would be affected by the flow, this can lead to errors on targeting and detection measurements.

HAOT is a Hypersonic Aerodynamics Optics Tools Python package developed to calculate different aerodynamic properties in a hypersonic medium. Its source code is available on [GitHub](#), the documentation is available on [Read the Docs](#) and an example on the usage of the package is given on the GitHub repo under the example folder.

## Statement of Need

Many techniques used to calculate optical properties are scattered across various papers, but there is no centralized repository containing all these calculations. Furthermore, some of these calculations require spectroscopy constants, which are often unclear or inconsistently presented in the literature. This package includes a constants module that provides and documents numerous spectroscopy constants for diatomic molecules.

This package has been used by ([Liza et al., 2023](#)). In this work he used results from Computational Fluid Dynamics (CFD) to calculate index optic properties of interest.

## Algorithms

The HAOT package, contains five modules:

- Aerodynamics
- Optics
- Quantum Mechanics
- Constants
- Conversions

Each module can be imported independently. The [documentation](#) explains he functions in each module as well as their usage. Docstrings were used, so the function prototypes and usage are also available in an interactive Python session. Results from these algorithms were compared with literature and a unit test was developed and it is located under test. \

This section provide some of the capabilities of the packages but not all of them. For instance the package can calculate some compressible flow properties such as isentropic, normal shock, and oblique shock relations. Please refer to the [documentation](#) to see the complete list of available functions.

## Results

Equation 1 was introduced by (Smith & Weintraub, n.d.), and it provides approximation and approximation for the index of refraction as a function of atmospheric altitude.

$$n(h) \approx 1 + \frac{K_1}{T(h)} \left( p(h) + K_2 \frac{e(h)}{T(h)} \right) \quad (1)$$

Where:  $K_1$  and  $K_2$  are constants,  $T$  is the temperature as a function of altitude,  $p$  is pressure as a function of altitude, and  $e(h)$  is the partial pressure of water vapor. \

Results for equation 1 are provided in the figure 1. This approximation is a useful way of seen the region in which the index of refraction will have a higher impact in a seeker's performance. As expected the critical region is between 20[km] to 30[km] above sea level, which is a region where seeker's performance is very important.




Figure 1: Atmospheric index of refraction for dry air.

The Gladstone-Dale constant is an important constant used to calculate the index of refraction (equation 2). For a dilute gas, the index of refraction can be approximated as:

$$n - 1 = \sum_{s=1}^N K_s \rho_s \quad (2)$$

Where:  $n$  is the index of refraction,  $K_s$  is the species Gladstone-Dale constants, and  $\rho_s$  is the species density. \

Figure 2 shows the Gladstone-Dale constants for a five-species gas. The species density was calculated using Computational Fluid Dynamics (CFD) tool, SU2 (W. T. Maier et al., 2021), (W. Maier et al., 2023). These densities were then loaded into the HAOT tool to calculate those figures.




Figure 2: Species Gladstone-Dale constants for a five-species gas.




Figure 3: Polarizability of O2 using Kerl's method [Kerl1992].




Figure 4: Boltzmann Distribution for  $N_2$ .

## Acknowledgements

The author gratefully thank Kyle Hanquist, who supported with the tool verification.

## References

- Liza, M., Tumuklu, O., & Hanquist, K. M. (2023, June). Nonequilibrium effects on aero-optics in hypersonic flows. *AIAA AVIATION 2023 Forum*. <https://doi.org/10.2514/6.2023-3736>
- Maier, W. T., Needels, J. T., Garbacz, C., Morgado, F., Alonso, J. J., & Fossati, M. (2021). SU2-NEMO: An open-source framework for high-mach nonequilibrium multi-species flows. *Aerospace*, 8, 193. <https://doi.org/10.3390/aerospace8070193>
- Maier, W., Needels, J. T., Alonso, J. J., Morgado, F., Garbacz, C., Fossati, M., Tumuklu, O., & Hanquist, K. M. (2023). Development of physical and numerical nonequilibrium modeling capabilities within the SU2-NEMO code. *AIAA AVIATION 2023 Forum*. <https://doi.org/10.2514/6.2023-3488>
- Smith, E., & Weintraub, S. (n.d.). <https://doi.org/10.1109/JRPROC.1953.274297>