

My Thesis Title

Matthew Lawrence Joyce

April 9, 2020

CONTENTS

1. <i>The Large Hadron Collider</i>	3
1.1 scratch section to work out outline...	3
1.2 Introduction	3
1.3 Injection Complex	3
1.4 Tunnel and Magnets	4
1.5 Luminosity	5
2. <i>CMS Detector</i>	7
3. <i>The Standard Model</i>	8
4. <i>Supersymmetry</i>	9

1. THE LARGE HADRON COLLIDER

1.1 *scratch section to work out outline...*

First thing to do is introduce the LHC as the large hadron collider and mention where it is located. Now I should give a brief overview of what it does. Now I need to talk about how it does this. That means I need to mention how the protons are accelerated. Then I need to explain how we contain and collide them once they are accelerated. This leads into talking about the tunnel and magnets used for steering the bunches.

1.2 *Introduction*

The Large Hadron Collider (LHC) is a 26.7 kilometer-long, two-ring particle accelerator and collider located on the border of France and Switzerland at the European Organization for Nuclear Research (CERN). During normal operations the LHC maintains two counter-rotating beams of proton bunches that collide at four interaction points (IP) with up to $\sqrt{s} = 14$ TeV center of mass energy and a luminosity of $10^{34} \text{cm}^{-2} \text{s}^{-1}$. The ALICE (Point 2), ATLAS (Point 1), CMS (Point 5), and LHC-b experiments each have a detector at one of these interaction points as scene in Figure 1.1 . The CMS and ATLAS are general-purpose detectors while LHC-b specializes in beauty quark studies. ALICE is a heavy-ion experiment which uses $^{208}\text{Pb} - p$ or $^{208}\text{Pb} - ^{208}\text{Pb}$ collisions that can also be produced by the LHC.

1.3 *Injection Complex*

In order to bring the protons from rest up to their target collision energy a series of accelerators, as shown in Figure 1.2, are used. The acceleration sequence begins with the injection of hydrogen gas into a duoplasmatron. Here a bombardment of electrons ionize the hydrogen atoms while an electric field pushes them through the duoplasmatron cavity. The result is 100 keV protons being passed on to a quadrupole magnet which guides them into the aperture of a linear accelerator (LINAC2). The radio frequency (RF)




Fig. 1.1: Interaction points of the LHC

cavities in LINAC2 accelerate the protons up to 50 MeV. At this point the protons are sent into one of four rings in the Proton Synchrotron Booster (PSB). The PSB repeatedly accelerates the protons around a circular path until they reach an energy of 1.4 GeV. The bunches of protons from each PSB ring are then sequentially injected into the single-ringed Proton Synchrotron (PS). Each bunch injected into the PS are captured by one of the "buckets" (Figure 1.3) provided by the PS RF system which also manipulates the bunches into the desired profile and proton density. These proton bunches are accelerated to 25 GeV and injected into the Super Proton Synchrotron (SPS) where they are accelerated to 450 GeV. Finally the proton bunches are injected into the LHC ring where they are accelerated to 6.5 TeV and collided to yield a center of mass energy of $\sqrt{s} = 13$ TeV.

1.4 Tunnel and Magnets

The LHC was designed to produce collisions with up to $\sqrt{s} = 14$ TeV. That requires confining and guiding 7 TeV protons around the circumference of the LHC ring. The ring is housed in a 4 meter-wide underground tunnel




Fig. 1.2: Layout of LHC accelerator complex.

that ranges in depth between 45 and 170 meters below the surface. This tunnel was repurposed from the Large Electron-Positron (LEP) Collider which previously occupied the space. For this reason the tunnel is not completely circular but is instead made up of alternating curved and straight sections of 2500 m and 530 m in length respectively. The straight sections, labeled 1-8 in Figure 1.1, are used as either experimental facilities or sites for hardware necessary for LHC operations such as RF cavities for momentum cleaning, quadrupole magnets for beam focusing, and sextupole magnets for acceleration and betatron cleaning. Steering a 7 TeV proton beam around the curved sections requires a magnetic field of 8.33 Tesla which is provided by 1223 superconducting dipole magnets cooled to 1.9 K using liquid helium. A cross section of the LHC dipole is shown in Figure 1.4.

1.5 Luminosity





Fig. 1.3: Proton bunch capture onto RF bucket.

CROSS SECTION OF LHC DIPOLE



CERN AC_HE107A_V02/02/98

Fig. 1.4: Cross section of LHC dipole [1]

2. CMS DETECTOR

3. THE STANDARD MODEL

4. SUPERSYMMETRY

BIBLIOGRAPHY

- [1] J.-L. Caron, “Cross section of lhc dipole.. dipole lhc: coupe transversale..” ”AC Collection. Legacy of AC. Pictures from 1992 to 2002.”, May 1998.