


Introduction

What is a Starspot?

Sunspots/starspots are dark blotches that appear on the surface of stars.

The underlying processes that govern their formation and death, as well as their effects on the host star, are poorly understood



It is widely accepted that starspots are caused by magnetic field beneath the stellar surface.



Why Study Starspots?

Starspots serve to mask the identity of distant stars by causing either superficial changes to the stars observable properties or deep structural changes to the star.

Starspots are proposed to explain the discontinuity between theoretical model and observed properties of stars (i.e. low-mass stars always appear 5% larger than theory predicts).





Past Work

Ash and Feiden (in prep) derived starspot properties assuming two different formation scenarios:


- Transient spots require that a star have 75% of its surface covered by spots ($\rho = 0.90$, $w = 0.84$).
- Long-Term spots that caused deep structural changes require that around half of the star to be covered in spots ($\rho = .56$, $w = 0.88$).

Results & Conclusions



Results

- Due to the random nature of each light curve, it was necessary to run the simulation multiple times in order to perform a statistical analysis of the results.
- These simulations were performed 1,000 times each for two different test cases.
- These test cases ($\rho = 0.90$, $w = 0.85$ and $\rho = 0.60$, $w = 0.88$) were chosen to test Ash's conditions for transient spots and long-term spots.
- Corrections were applied to simulate the random orientation of stars.
- Minute variations that could not be observed were omitted in the corrected distributions.




Conclusions

- This statistical analysis shows that stars with higher surface coverage tend to show more brightness variability than those with lower surface coverage.
- Our findings, when compared to data collected by Jackson & Jeffries in the young open cluster NGC 2516, indicate that the transient (short-term) spot model more closely matches observed phenomena than the long-term spot model.
- This implies that spots are transient features, and they cause little if any changes to stellar structure.


References:
Gough, D. (2010). *Vainu Bappu Memorial Lecture: What is a Sunspot?* Cambridge, UK: Institute of Astronomy, University of Cambridge, U.K.
Jackson, R. J., & Jeffries, R. D. (2013). On the relationship between the size and surface coverage of starspots on magnetically active low-mass stars. *Monthly Notices of the Royal Astronomical Society*, 431(2), 1883-1890. doi:10.1093/mnras/sts04.
Jackson, R. J., & Jeffries, R. D. (2012). Why do some young cool stars show spot modulation while others do not? *Monthly Notices of the Royal Astronomical Society*, 423(3), 2966-2976. doi:10.1111/j.1365-2966.2012.21119.x

Methodology



3D Model to 2D Projection

- A 3D model sphere was created, and this sphere was divided along latitudinal and longitudinal lines to create a surface tessellated with small cells.
- Certain cells were randomly selected to be 'spot cells' at any desired surface coverage (ρ).
- A surface projection was chosen that conserved apparent surface area.
- The 2D projection was written to only display the cells in the visible hemisphere of the 3D model



Light Curve Generation and Amplitude Variation

- Spot cells were given a lower brightness value based on an assumed temperature ratio between the spots and the background photosphere (w).
- The model was then rotated, and a 2D projection was created for each discrete rotation. The brightness from each projection was recorded.
- This data was plotted against rotational degree (ϕ) to create a light curve, and the brightness variation was recorded.