

# What 50 Million Drawings Can Tell Us About Shared Meaning

Molly Lewis and Gary Lupyan, University of Wisconsin-Madison

## How to interrogate shared meaning?



## Drawings as a window into meaning



50M drawings across 345 categories ( $N = 15M$ )


[quickdraw.withgoogle.com](http://quickdraw.withgoogle.com)

## Quantifying Similarity


Hausdorff Distance — drawing similarity as the maximum Euclidean distance between two sets of points (Huttenlocher, Klanderman, & Rucklidge, 1993).

### Measure Validation:

- 1500 drawing pairs from two categories
- 20 pairs from each decile of similarity/item
- Rated for similarity on 7-pt. Likert Scale
- Each participant ( $N=100$ ) rated 50 drawing pairs
- Human judgements highly correlated with Hausdorff Distance ( $r = .29$ ;  $p < .001$ )



## Cross-Cultural Variability in Drawings



## Predicting Drawing Variability with Language

### Drawing Similarity:

- 50 drawings from each of 39 countries
- Items from 4 categories: food (e.g., "cake"), weather ("cloud"), artifact (e.g., "key"), place (e.g., "jail")
- For each country pair, calculated mean drawing similarity for each category.


### Independent Cultural Similarity Measures:

- Lexical semantics – Alignment from word embedding models (Bojanowski et al., 2016)
- Lexical forms – ASJP distance (Bakker, et al., 2009; Dediu, in press)
- Grammar – WALS distance (Dediu, in press)
- Physical distance

**Semantics predict independent variance in drawing distances, suggesting speakers of semantically more similar languages have more similar non-linguistic representations.**

Next: What properties characterize these differences?

### Drawing Distance vs. Semantic Distance



### What predicts drawing similarity?

