

# A Machine Learning Model For Predicting Deterioration of COVID-19 Inpatients

Omer Noy<sup>\*,1</sup>, Dan Coster<sup>\*,1,2</sup>, Maya Metzger<sup>1</sup>, Itai Attar<sup>2</sup>, Shani Shenhar-Tsafraty<sup>2,3</sup>, Shlomo Berliner<sup>2,3</sup>, Galia Rahav<sup>2,4</sup>, Ori Rogowski<sup>2,3</sup>, Ron Shamir<sup>1</sup>

<sup>1</sup>Blavatnik School of Computer Science and <sup>2</sup>Sackler Faculty of Medicine, Tel-Aviv University, <sup>3</sup>Department of Internal Medicine "E" & "C", Tel-Aviv Sourasky Medical Center, <sup>4</sup>Infectious Diseases Unit, Sheba Medical Center. \*Equal contribution. omernoy4@gmail.com | dancoster@gmail.com.

## Background

The COVID-19 pandemic has been spreading worldwide since December 2019, presenting an urgent threat to global health. Due to the limited understanding of disease progression and of the risk factors for the disease, it is a clinical challenge to predict which hospitalized patients will deteriorate. Early prediction can allow physicians to take measures that prevent or lessen condition worsening.

**Our goal** is to develop a predictive model for the early identification of patients at risk for clinical deterioration by analyzing Electronic Medical Records (EMR) data of COVID-19 inpatients.

## Datasets

The **development set** contains medical records of 662 inpatients admitted to the Sheba Medical Center (Sheba) between March and December 2020, who were diagnosed with COVID-19. The **validation set** contains medical records of 417 inpatients admitted to the Tel-Aviv Sourasky Medical Center (TASMC) between March and September 2020, who were diagnosed with COVID-19.



N = 662



N = 417

The data used was extracted from longitudinal EMRs and included both time-independent (static) and temporal (dynamic) features, such as demographics, background disease, vital signs and lab measurements. A total of **100 features** were selected per patient.

## Target Definition

**The National Early Warning Score 2 (NEWS2)** is used as a clinical risk score for early recognition of patients with severe infection. Liao et al. (2020) suggested an early warning score for COVID-19 patients termed "modified-NEWS2" (mNEWS2), which adds to the NEWS2 formula the factor age  $\geq 65$  years.


Points	3	2	1	0	1	2	3
Age				<65			$\geq 65$
Respiratory Rate	$\leq 8$		9-11	12-20		21-24	$\geq 25$
Oxygen Saturation	$\leq 91$	92-93	94-95	$\geq 96$			
Supplemental Oxygen		Yes		No			
Systolic BP	$\leq 90$	91-100	101-110	111-219			$\geq 220$
Heart Rate	$\leq 40$		41-50	51-90	91-110	111-130	$\geq 131$
Consciousness				Alert			Drowsiness Lethargy Coma Confusion
Temperature	$\leq 35.0$		35.1-36.0	36.1-38.0	38.1-39.0		$\geq 39.1$

The mNEWS2 score (Liao et al, 2020)


mNEWS2  $\geq 7$



Observations with a high mNEWS2 score ( $\geq 7$ ) recorded in the preceding 7-30 hours were called **positive**, and the rest were called negative. We did not use the data in 6 hours prior to the event.




## Model Development Overview




## Model Selection

The performance of 14 machine learning models that predict mNEWS2  $\geq 7$  was tested, using 10-fold cross-validation over the training set. We chose CatBoost as our final prediction model.



## Results



**External validation:** The final model achieved **AUROC=0.76** and **AUPR=0.7** in external validation on the TASMC data, slightly less than in the development dataset.

## Conclusions and Future Work

We utilized machine learning models for predicting deterioration within the next 7-30 hours based on EHR data of COVID-19 inpatients. On held-out data, the model achieved AUROC of 0.84 and AUPR of 0.74. It achieved values of 0.76 and 0.7 respectively in external validation on data from a different hospital.

### Future work:

- Models for early warning on approaching oxygen deficiency deterioration events.
- Multi-center model training.
- Comparative analysis across different COVID-19 waves.
- Existing models benchmarking.

Study supported in part by the Israel Science Foundation (grant 1339/18 and grant 3165/19 within the Israel Precision Medicine Partnership program) and German-Israeli Project DFG RE 4193/1-1. DC and ON were supported in part by fellowships from the Edmond J. Safra Center for Bioinformatics at Tel Aviv University.