

CSI - 3105 Design & Analysis of Algorithms

Course 16


Jean-Lou De Carufel

Fall 2019

Final Remark About Dynamic Programming

Consider a directed graph $G = (V, E)$, where all edges have weight 1. Let $u, v \in V$ be two vertices.


Assume we know a vertex w on the shortest path from u to v .



Final Remark About Dynamic Programming

Consider a directed graph $G = (V, E)$, where all edges have weight 1. Let $u, v \in V$ be two vertices.


Assume we know a vertex w on the shortest path from u to v .



Final Remark About Dynamic Programming

Consider a directed graph $G = (V, E)$, where all edges have weight 1. Let $u, v \in V$ be two vertices.


Assume we know a vertex w on the shortest path from u to v .



Final Remark About Dynamic Programming

Consider a directed graph $G = (V, E)$, where all edges have weight 1. Let $u, v \in V$ be two vertices.


Assume we know a vertex w on the shortest path from u to v .



Final Remark About Dynamic Programming

Consider a directed graph $G = (V, E)$, where all edges have weight 1. Let $u, v \in V$ be two vertices.

Assume we know a vertex w on the shortest path from u to v .




Optimal substructure!

Final Remark About Dynamic Programming

Consider a directed graph $G = (V, E)$, where all edges have weight 1. Let $u, v \in V$ be two vertices.


Assume we know a vertex w on the **longest** path from u to v .



Final Remark About Dynamic Programming

Consider a directed graph $G = (V, E)$, where all edges have weight 1. Let $u, v \in V$ be two vertices.


Assume we know a vertex w on the **longest** path from u to v .




Final Remark About Dynamic Programming

Consider a directed graph $G = (V, E)$, where all edges have weight 1. Let $u, v \in V$ be two vertices.

Assume we know a vertex w on the **longest** path from u to v .



Example:



NO!

In fact, computing the longest path is NP-Hard...

Chapter 6: P vs NP

§6.1 Introduction

An algorithm has *polynomial running time* if there is a constant $c \geq 1$ such that for every input of length n , the algorithm takes $O(n^c)$ time.

§6.1 Introduction

An algorithm has *polynomial running time* if there is a constant $c \geq 1$ such that for every input of length n , the algorithm takes $O(n^c)$ time.

An algorithm has *exponential running time* if there is a constant $c \geq 1$ such that for every input of length n , the algorithm takes $O(2^{n^c})$ time.

§6.1 Introduction

An algorithm has *polynomial running time* if there is a constant $c \geq 1$ such that for every input of length n , the algorithm takes $O(n^c)$ time.

An algorithm has *exponential running time* if there is a constant $c \geq 1$ such that for every input of length n , the algorithm takes $O(2^{n^c})$ time.

Intuitively speaking,

polynomial means good, fast, efficient, easy, ...

exponential means bad, slow, “try all possible solutions”, difficult, ...

Complexity Class P

A *decision problem* is a problem for which the answer is “yes” or “no”.

P = set of all decision problems that can be solved in polynomial time

Complexity Class P

A *decision problem* is a problem for which the answer is “yes” or “no”.

P = set of all decision problems that can be solved in polynomial time

Examples of problems that are in P :

- Is a given input graph connected?

Complexity Class P

A *decision problem* is a problem for which the answer is “yes” or “no”.

P = set of all decision problems that can be solved in polynomial time

Examples of problems that are in P :

- Is a given input graph connected?
- Is a given input graph bipartite?

Complexity Class P

A *decision problem* is a problem for which the answer is “yes” or “no”.

P = set of all decision problems that can be solved in polynomial time

Examples of problems that are in P :

- Is a given input graph connected?
- Is a given input graph bipartite?
- Is a given input sequence sorted?

Complexity Class P

A *decision problem* is a problem for which the answer is “yes” or “no”.

P = set of all decision problems that can be solved in polynomial time

Examples of problems that are in P :


- Is a given input graph connected?
- Is a given input graph bipartite?
- Is a given input sequence sorted?
- Does a given input graph contain an Euler cycle? An *Euler cycle* is a cycle that traverses each edge exactly once.

Other Problems

HAM-CYCLE

input: An undirected graph $G = (V, E)$ stored using adjacency lists.

question: Does G contain a Hamiltonian cycle? A *Hamiltonian cycle* is a cycle that traverses each vertex exactly once.




Other Problems

HAM-CYCLE

input: An undirected graph $G = (V, E)$ stored using adjacency lists.

question: Does G contain a Hamiltonian cycle? A *Hamiltonian cycle* is a cycle that traverses each vertex exactly once.




Other Problems

HAM-CYCLE

input: An undirected graph $G = (V, E)$ stored using adjacency lists.

question: Does G contain a Hamiltonian cycle? A *Hamiltonian cycle* is a cycle that traverses each vertex exactly once.



Not known if this problem is in *P*!

Other Problems

Traveling Salesman Problem (TSP)

input:

- A complete directed graph $G = (V, E)$, where each edge $(u, v) \in E$ has a weight $wt(u, v) > 0$.
- An integer K .

question: Does G contain a Hamiltonian cycle of total weight at most K ?


Other Problems

Traveling Salesman Problem (TSP)

input:

- A complete directed graph $G = (V, E)$, where each edge $(u, v) \in E$ has a weight $wt(u, v) > 0$.
- An integer K .

question: Does G contain a Hamiltonian cycle of total weight at most K ?




Other Problems

Traveling Salesman Problem (TSP)

input:

- A complete directed graph $G = (V, E)$, where each edge $(u, v) \in E$ has a weight $wt(u, v) > 0$.
- An integer K .

question: Does G contain a Hamiltonian cycle of total weight at most K ?



Not known if this problem is in P !

Other Problems

SUBSET-SUM

input: A set S of integers together with an integer t

question: Is there a subset S' of S such that

$$\sum_{x \in S'} x = t?$$

Other Problems

SUBSET-SUM

input: A set S of integers together with an integer t

question: Is there a subset S' of S such that

$$\sum_{x \in S'} x = t?$$

Example:

$$S = \{5, 7, 56, 23, 2902\}$$

$$t = 2963$$

Other Problems

SUBSET-SUM

input: A set S of integers together with an integer t

question: Is there a subset S' of S such that

$$\sum_{x \in S'} x = t?$$

Example:

$$S = \{5, 7, 56, 23, 2902\}$$

$$t = 2963$$

Yes! $S' = \{5, 56, 2902\}$

Other Problems

SUBSET-SUM

input: A set S of integers together with an integer t

question: Is there a subset S' of S such that

$$\sum_{x \in S'} x = t?$$

Example:

$$S = \{5, 7, 56, 23, 2902\}$$

$$t = 2963$$

$$\text{Yes! } S' = \{5, 56, 2902\}$$


Not known if this problem is in P !

Other Problems

CLIQUE

input: An undirected graph $G = (V, E)$ together with an integer K

question: Does G contain a clique (complete subgraph) with K vertices?




$$K = 4$$

Other Problems

CLIQUE

input: An undirected graph $G = (V, E)$ together with an integer K

question: Does G contain a clique (complete subgraph) with K vertices?




$$K = 4$$

Other Problems

CLIQUE

input: An undirected graph $G = (V, E)$ together with an integer K

question: Does G contain a clique (complete subgraph) with K vertices?



Not known if this problem is in P !

Other Problems

For each of the 4 previous problems,

- Not known if it can be solved in polynomial time.

Other Problems

For each of the 4 previous problems,

- Not known if it can be solved in polynomial time.
- If the answer to the question is YES, then
 - There is a “short” proof for this.

Other Problems

For each of the 4 previous problems,

- Not known if it can be solved in polynomial time.
- If the answer to the question is YES, then
 - There is a “short” proof for this.

Here, “short” means the length of the proof is “polynomial in the length of the input”.

Other Problems

For each of the 4 previous problems,

- Not known if it can be solved in polynomial time.
- If the answer to the question is YES, then
 - There is a “short” proof for this.

Here, “short” means the length of the proof is “polynomial in the length of the input”.

- If someone gives us such a short proof, then we can “easily” verify this proof.

Other Problems

For each of the 4 previous problems,

- Not known if it can be solved in polynomial time.
- If the answer to the question is YES, then
 - There is a “short” proof for this.

Here, “short” means the length of the proof is “polynomial in the length of the input”.

- If someone gives us such a short proof, then we can “easily” verify this proof.

Here, “easily” means “in polynomial time”.

HAM-CYCLE

proof/solution: sequence v_1, \dots, v_k of vertices.

HAM-CYCLE

proof/solution: sequence v_1, \dots, v_k of vertices.

verification: • $k = n?$

HAM-CYCLE

proof/solution: sequence v_1, \dots, v_k of vertices.

verification:

- $k = n$?
- no duplicates in v_1, \dots, v_k ?

HAM-CYCLE

proof/solution: sequence v_1, \dots, v_k of vertices.

- verification:
- $k = n$?
 - no duplicates in v_1, \dots, v_k ?
 - For each $1 \leq i < k - 1$, $\{v_i, v_{i+1}\}$ is an edge?

HAM-CYCLE

proof/solution: sequence v_1, \dots, v_k of vertices.

- verification:
- $k = n$?
 - no duplicates in v_1, \dots, v_k ?
 - For each $1 \leq i < k - 1$, $\{v_i, v_{i+1}\}$ is an edge?
 - $\{v_k, v_1\}$ is an edge?

TSP

proof/solution: sequence v_1, \dots, v_k of vertices.

TSP

proof/solution: sequence v_1, \dots, v_k of vertices.

verification: • $k = n?$

TSP

proof/solution: sequence v_1, \dots, v_k of vertices.

verification:

- $k = n$?
- no duplicates in v_1, \dots, v_k ?

TSP

proof/solution: sequence v_1, \dots, v_k of vertices.

- verification:
- $k = n$?
 - no duplicates in v_1, \dots, v_k ?
 - Do we have

$$\sum_{i=1}^{k-1} wt(v_i, v_{i+1}) + wt(v_k, v_1) \leq K?$$

SUBSET-SUM

proof/solution: A set S' .

SUBSET-SUM

proof/solution: A set S' .

verification: • Is S' a subset of S ?

SUBSET-SUM

proof/solution: A set S' .

verification:

- Is S' a subset of S ?
- Do we have

$$\sum_{x \in S'} x = t?$$

CLIQUE

proof/solution: A set V' .

CLIQUE

proof/solution: A set V' .

verification:

- Is V' a subset of V ?

CLIQUE

proof/solution: A set V' .

- verification:
- Is V' a subset of V ?
 - Do we have $|V'| = K$?

CLIQUE

proof/solution: A set V' .

- verification:
- Is V' a subset of V ?
 - Do we have $|V'| = K$?
 - For each $u, v \in V'$ such that $u \neq v$, is $\{u, v\}$ an edge in E ?