

# CSI - 3105 Design & Analysis of Algorithms

## Course 8

Jean-Lou De Carufel

Fall 2019

---

**Algorithm** *Dijkstra*( $G, s$ )

---

```
1: for each vertex  $v \in V$  do
2:    $d(v) = \infty$ 
3: end for
4:  $d(s) = 0$ 
5:  $S = \{\}$ 
6:  $Q = V$ 
7: while  $Q \neq \{\}$  do
8:    $u =$  vertex in  $Q$  for which  $d(u)$  is minimum
9:   delete  $u$  from  $Q$ 
10:  insert  $u$  into  $S$ 
11:  for each edge  $(u, v)$  do
12:     $d(v) = \min \{d(v), d(u) + wt(u, v)\}$ 
13:  end for
14: end while
```

---


# Dijkstra Algorithm is Correct

## Theorem

*Let  $G = (V, E)$  be a weighted directed graph and  $s \in V$  be a source vertex. Dijkstra algorithm finds the lengths of the shortest paths from  $s$  to all vertices in  $V$ .*

# Special Paths and Induction Hypotheses


PROOF: We say that a path from  $s$  to a vertex  $v$  is *special* if all vertices on that path belong to  $S$ , except maybe  $v$ .



A *special path* from  $s$  to  $v$ .

# Special Paths and Induction Hypotheses

PROOF: We say that a path from  $s$  to a vertex  $v$  is *special* if all vertices on that path belong to  $S$ , except maybe  $v$ .



A *special path* from  $s$  to  $v$ .

We prove by induction that

- (a) if a vertex  $u$  is in  $S$ , then  $d(u)$  gives the length of a shortest path from  $s$  to  $u$  and
- (b) if a vertex  $u$  is not in  $S$ , then  $d(u)$  gives the length of a shortest *special path* from  $s$  to  $u$ .

- (a) if a vertex  $u$  is in  $S$ , then  $d(u)$  gives the length of a shortest path from  $s$  to  $u$  and
- (b) if a vertex  $u$  is not in  $S$ , then  $d(u)$  gives the length of a shortest *special path* from  $s$  to  $u$ .

- (a) if a vertex  $u$  is in  $S$ , then  $d(u)$  gives the length of a shortest path from  $s$  to  $u$  and
- (b) if a vertex  $u$  is not in  $S$ , then  $d(u)$  gives the length of a shortest *special path* from  $s$  to  $u$ .

**Base case:**

- (a) At the beginning,  $S = \{ \}$ , so (a) is vacuously true.

- (a) if a vertex  $u$  is in  $S$ , then  $d(u)$  gives the length of a shortest path from  $s$  to  $u$  and
- (b) if a vertex  $u$  is not in  $S$ , then  $d(u)$  gives the length of a shortest *special path* from  $s$  to  $u$ .

**Base case:**

- (a) At the beginning,  $S = \{ \}$ , so (a) is vacuously true.
- (b) Since  $S = \{ \}$ , the other vertices simply cannot be reached by following a special path from  $s$ . Since  $d$  is initialized to  $\infty$ , then (b) also holds when the algorithm starts.



- (a) if a vertex  $u$  is in  $S$ , then  $d(u)$  gives the length of a shortest path from  $s$  to  $u$  and
- (b) if a vertex  $u$  is not in  $S$ , then  $d(u)$  gives the length of a shortest *special path* from  $s$  to  $u$ .

### Base case:

- (a) At the beginning,  $S = \{ \}$ , so (a) is vacuously true.
- (b) Since  $S = \{ \}$ , the other vertices simply cannot be reached by following a special path from  $s$ . Since  $d$  is initialized to  $\infty$ , then (b) also holds when the algorithm starts.

**Induction hypothesis:** Assume that both (a) and (b) hold right before we add a new vertex  $v$  to  $S$ .

- (a) if a vertex  $u$  is in  $S$ , then  $d(u)$  gives the length of a shortest path from  $s$  to  $u$  and
- (b) if a vertex  $u$  is not in  $S$ , then  $d(u)$  gives the length of a shortest *special path* from  $s$  to  $u$ .

### Base case:


- (a) At the beginning,  $S = \{ \}$ , so (a) is vacuously true.
- (b) Since  $S = \{ \}$ , the other vertices simply cannot be reached by following a special path from  $s$ . Since  $d$  is initialized to  $\infty$ , then (b) also holds when the algorithm starts.

**Induction hypothesis:** Assume that both (a) and (b) hold right before we add a new vertex  $v$  to  $S$ .

We address induction steps (a) and (b) separately.


## Induction Step (a)

**Induction step (a):** By the induction hypothesis (a), before the addition of  $v$ , we already know a shortest paths from  $s$  to all vertices that are in  $S$ . Adding  $v$  to  $S$  does not change these shortest paths.




## Induction Step (a)

**Induction step (a):** By the induction hypothesis (a), before the addition of  $v$ , we already know a shortest paths from  $s$  to all vertices that are in  $S$ . Adding  $v$  to  $S$  does not change these shortest paths.




As for node  $v$ , it is about to be inserted in  $S$ . Before adding it to  $S$ , we must check that  $d(v)$  gives the length of a shortest path from  $s$  to  $v$ .

## Induction Step (a)



So we compare  $d(v)$  to the lengths of all paths from  $s$  to  $v$ .


## Induction Step (a)



So we compare  $d(v)$  to the lengths of all paths from  $s$  to  $v$ .

There are two kinds of paths: (1) the paths that are special and (2) the ones that are not special.

# Induction Step (a)




So we compare  $d(v)$  to the lengths of all paths from  $s$  to  $v$ .

There are two kinds of paths: (1) the paths that are special and (2) the ones that are not special.

- (1) By the induction hypothesis (b), we already know that  $d(v)$  is less than or equal to the length of any special path from  $s$  to  $v$ .

# Induction Step (a)




So we compare  $d(v)$  to the lengths of all paths from  $s$  to  $v$ .

There are two kinds of paths: (1) the paths that are special and (2) the ones that are not special.

- (1) By the induction hypothesis (b), we already know that  $d(v)$  is less than or equal to the length of any special path from  $s$  to  $v$ .
- (2) A non-special path from  $s$  to  $v$  is one which contains at least one vertex  $x \neq v$  that is not in  $S$ .



# Induction Step (a)




So we compare  $d(v)$  to the lengths of all paths from  $s$  to  $v$ .

There are two kinds of paths: (1) the paths that are special and (2) the ones that are not special.

- (1) By the induction hypothesis (b), we already know that  $d(v)$  is less than or equal to the length of any special path from  $s$  to  $v$ .
- (2) A non-special path from  $s$  to  $v$  is one which contains at least one vertex  $x \neq v$  that is not in  $S$ . But such a path has length at least

$$d(x)$$

# Induction Step (a)



So we compare  $d(v)$  to the lengths of all paths from  $s$  to  $v$ .

There are two kinds of paths: (1) the paths that are special and (2) the ones that are not special.

- (1) By the induction hypothesis (b), we already know that  $d(v)$  is less than or equal to the length of any special path from  $s$  to  $v$ .
- (2) A non-special path from  $s$  to  $v$  is one which contains at least one vertex  $x \neq v$  that is not in  $S$ . But such a path has length at least

$$d(x) \geq d(v). \quad (\text{do you see why?})$$

## Induction Step (a)

Hence,  $d(v)$  is less than or equal to the length of any path from  $s$  to  $v$ ,

## Induction Step (a)

Hence,  $d(v)$  is less than or equal to the length of any path from  $s$  to  $v$ , which means that  $d(v)$  gives the length of a shortest path from  $s$  to  $v$ .


## Induction Step (a)

Hence,  $d(v)$  is less than or equal to the length of any path from  $s$  to  $v$ , which means that  $d(v)$  gives the length of a shortest path from  $s$  to  $v$ .

So the induction step is complete for (a).


## Induction Step (b)

**Induction step (b):** Consider a node  $w \neq v$  which is not in  $S$ .




## Induction Step (b)

**Induction step (b):** Consider a node  $w \neq v$  which is not in  $S$ . Let  $y$  be the last vertex in  $S$  encountered on a shortest special path from  $s$  to  $w$ .



## Induction Step (b)


**Induction step (b):** Consider a node  $w \neq v$  which is not in  $S$ . Let  $y$  be the last vertex in  $S$  encountered on a shortest special path from  $s$  to  $w$ . We consider two cases: (1)  $y \neq v$  or (2)  $y = v$ .





## Induction Step (b)


**Induction step (b):** Consider a node  $w \neq v$  which is not in  $S$ . Let  $y$  be the last vertex in  $S$  encountered on a shortest special path from  $s$  to  $w$ . We consider two cases: (1)  $y \neq v$  or (2)  $y = v$ .



(1) If  $y \neq v$ , then adding  $v$  to  $S$  does not change the value of  $d(w)$ .

## Induction Step (b)


**Induction step (b):** Consider a node  $w \neq v$  which is not in  $S$ . Let  $y$  be the last vertex in  $S$  encountered on a shortest special path from  $s$  to  $w$ . We consider two cases: (1)  $y \neq v$  or (2)  $y = v$ .



- (1) If  $y \neq v$ , then adding  $v$  to  $S$  does not change the value of  $d(w)$ . Hence, by the induction hypothesis (b),  $d(w)$  gives the length of a shortest special path from  $s$  to  $w$ .

## Induction Step (b)


**Induction step (b):** Consider a node  $w \neq v$  which is not in  $S$ . Let  $y$  be the last vertex in  $S$  encountered on a shortest special path from  $s$  to  $w$ . We consider two cases: (1)  $y \neq v$  or (2)  $y = v$ .



(2) If  $y = v$ , then by doing  $d(w) = \min \{d(w), d(v) + wt(v, w)\}$ ,

## Induction Step (b)


**Induction step (b):** Consider a node  $w \neq v$  which is not in  $S$ . Let  $y$  be the last vertex in  $S$  encountered on a shortest special path from  $s$  to  $w$ . We consider two cases: (1)  $y \neq v$  or (2)  $y = v$ .



- (2) If  $y = v$ , then by doing  $d(w) = \min \{d(w), d(v) + wt(v, w)\}$ , we ensure that after adding  $v$  to  $S$ ,  $d(w)$  gives the length of a shortest special path from  $s$  to  $w$ .

## Induction Step (b)

**Induction step (b):** Consider a node  $w \neq v$  which is not in  $S$ . Let  $y$  be the last vertex in  $S$  encountered on a shortest special path from  $s$  to  $w$ . We consider two cases: (1)  $y \neq v$  or (2)  $y = v$ .



- (2) If  $y = v$ , then by doing  $d(w) = \min \{d(w), d(v) + wt(v, w)\}$ , we ensure that after adding  $v$  to  $S$ ,  $d(w)$  gives the length of a shortest special path from  $s$  to  $w$ . □

Question: Design a deterministic algorithm to solve the following problem.

**input:** A weighted directed graph  $G = (V, E)$  together with a source vertex  $s \in V$ .

**output:** For each vertex  $v \in V$ , return `TRUE` if there is a *unique* shortest path from  $s$  to  $v$  and return `FALSE` if there is more than one shortest path from  $s$  to  $v$ .

What do you think of the following solution?

Initialize  $unique[v] = \text{TRUE}$  for all vertices  $v \in V$ .

Change the for-loop (inside the while-loop) by

```
1: for each edge  $(u, v)$  do  
2:   if  $d[u] + wt(u, v) < d[v]$  then  
3:      $d[v] = d[u] + wt(u, v)$   
4:   else if  $d[u] + wt(u, v) = d[v]$  then  
5:      $unique[v] = \text{FALSE}$ .  
6:   end if  
7: end for
```

What do you think of the following solution?

Initialize  $unique[v] = \text{TRUE}$  for all vertices  $v \in V$ .

Change the for-loop (inside the while-loop) by

```
1: for each edge  $(u, v)$  do  
2:   if  $d[u] + wt(u, v) < d[v]$  then  
3:      $d[v] = d[u] + wt(u, v)$   
4:      $unique[v] = \text{TRUE}$   
5:   else if  $d[u] + wt(u, v) = d[v]$  then  
6:      $unique[v] = \text{FALSE}$   
7:   end if  
8: end for
```



What do you think of the following solution?

Initialize  $unique[v] = \text{TRUE}$  for all vertices  $v \in V$ .

Change the for-loop (inside the while-loop) by

```
1: for each edge  $(u, v)$  do  
2:   if  $d[u] + wt(u, v) < d[v]$  then  
3:      $d[v] = d[u] + wt(u, v)$   
4:     if  $unique[u] = \text{FALSE}$  then  
5:        $unique[v] = \text{FALSE}$   
6:     end if  
7:   else if  $d[u] + wt(u, v) = d[v]$  then  
8:      $unique[v] = \text{FALSE}$   
9:   end if
```

What do you think of the following solution?

Initialize  $unique[v] = \text{TRUE}$  for all vertices  $v \in V$ .

Change the for-loop (inside the while-loop) by

```
1: for each edge  $(u, v)$  do  
2:   if  $d[u] + wt(u, v) < d[v]$  then  
3:      $d[v] = d[u] + wt(u, v)$   
4:     if  $unique[u] = \text{FALSE}$  then  
5:        $unique[v] = \text{FALSE}$   
6:     else  
7:        $unique[v] = \text{TRUE}$   
8:     end if  
9:   else if  $d[u] + wt(u, v) = d[v]$  then  
10:     $unique[v] = \text{FALSE}$   
11:   end if  
12: end for
```

What if you want to know whether or not there are at least 3 different shortest paths?

What if you want to know the exact number of shortest paths from  $s$  to any vertex  $v \in V$ ?