

CSI - 3105 Design & Analysis of Algorithms

Course 19

Jean-Lou De Carufel

Fall 2019

Example of a Reduction


$VERTEX - COVER = \{(G, K) \mid \text{graph } G = (V, E) \text{ contains a vertex cover with } K \text{ vertices.}\}$

Vertex cover: subset V' of V such that for each edge $\{u, v\}$ of G , at least one of u and v is in V' .

Example of a Reduction

$VERTEX - COVER = \{(G, K) \mid \text{graph } G = (V, E) \text{ contains a vertex cover with } K \text{ vertices.}\}$

Vertex cover: subset V' of V such that for each edge $\{u, v\}$ of G , at least one of u and v is in V' .




$\{2, 3, 5, 8\}$: vertex cover of size 4

Example of a Reduction

$VERTEX - COVER = \{(G, K) \mid \text{graph } G = (V, E) \text{ contains a vertex cover with } K \text{ vertices.}\}$

Vertex cover: subset V' of V such that for each edge $\{u, v\}$ of G , at least one of u and v is in V' .





$\{2, 3, 5, 8\}$: vertex cover of size 4



We want to show that

$$CLIQUE \leq_P VERTEX - COVER.$$



What is the connection between clique and vertex cover?

 G  \overline{G}



What is the connection between clique and vertex cover?





What is the connection between clique and vertex cover?




What is the connection between clique and vertex cover?




What is the connection between clique and vertex cover?



Is this a coincidence?




To prove the polynomial-time reduction $CLIQUE \leq_P VERTEX - COVER$, we need a function f which satisfies the *famous 3 properties*.



To prove the polynomial-time reduction $CLIQUE \leq_P VERTEX - COVER$, we need a function f which satisfies the *famous 3 properties*. Let $n = |V|$. We take

$$f(G, K) = (\bar{G}, n - K)$$




To prove the polynomial-time reduction $CLIQUE \leq_P VERTEX - COVER$, we need a function f which satisfies the *famous 3 properties*. Let $n = |V|$. We take

$$f(G, K) = (\bar{G}, n - K)$$

We have

- ① f maps inputs for CLIQUE to inputs for VERTEX-COVER.




To prove the polynomial-time reduction $CLIQUE \leq_P VERTEX - COVER$, we need a function f which satisfies the *famous 3 properties*. Let $n = |V|$. We take

$$f(G, K) = (\overline{G}, n - K)$$

We have

- ① f maps inputs for CLIQUE to inputs for VERTEX-COVER.
- ② Time to construct $(\overline{G}, n - K)$, when given (G, K) , is $O(|V| + |E|)$ which is polynomial in the size of (G, K) .



To prove the polynomial-time reduction $CLIQUE \leq_P VERTEX - COVER$, we need a function f which satisfies the *famous 3 properties*. Let $n = |V|$. We take

$$f(G, K) = (\overline{G}, n - K)$$

We have

- ① f maps inputs for CLIQUE to inputs for VERTEX-COVER.
- ② Is it true that G has a clique of size K if and only if \overline{G} has a vertex cover of size $n - K$?
- ③ Time to construct $(\overline{G}, n - K)$, when given (G, K) , is $O(|V| + |E|)$ which is polynomial in the size of (G, K) .

Let us prove Property 2.

Let us prove Property 2. We must prove that G has a clique of size K if and only if \overline{G} has a vertex cover of size $n - K$.

Let us prove Property 2. We must prove that G has a clique of size K if and only if \overline{G} has a vertex cover of size $n - K$.

$[\Rightarrow]$

Assume $G = (V, E)$ has a clique V' of size K .

Let us prove Property 2. We must prove that G has a clique of size K if and only if \overline{G} has a vertex cover of size $n - K$.

$[\Rightarrow]$

Assume $G = (V, E)$ has a clique V' of size K .

We will prove that $V \setminus V'$ is a vertex cover of size $n - K$.

Let us prove Property 2. We must prove that G has a clique of size K if and only if \overline{G} has a vertex cover of size $n - K$.

$[\Rightarrow]$

Assume $G = (V, E)$ has a clique V' of size K .

We will prove that $V \setminus V'$ is a vertex cover of size $n - K$.

Let (u, v) be an edge in \overline{G} . Then (u, v) is not an edge in G .

Let us prove Property 2. We must prove that G has a clique of size K if and only if \overline{G} has a vertex cover of size $n - K$.

$[\Rightarrow]$

Assume $G = (V, E)$ has a clique V' of size K .

We will prove that $V \setminus V'$ is a vertex cover of size $n - K$.

Let (u, v) be an edge in \overline{G} . Then (u, v) is not an edge in G . Hence, since V' is a clique, u and v are not both in V' .

Let us prove Property 2. We must prove that G has a clique of size K if and only if \overline{G} has a vertex cover of size $n - K$.

$[\Rightarrow]$

Assume $G = (V, E)$ has a clique V' of size K .

We will prove that $V \setminus V'$ is a vertex cover of size $n - K$.

Let (u, v) be an edge in \overline{G} . Then (u, v) is not an edge in G . Hence, since V' is a clique, u and v are not both in V' . Hence, at least one of u and v is in $V \setminus V'$.

Let us prove Property 2. We must prove that G has a clique of size K if and only if \overline{G} has a vertex cover of size $n - K$.

$[\Rightarrow]$

Assume $G = (V, E)$ has a clique V' of size K .

We will prove that $V \setminus V'$ is a vertex cover of size $n - K$.

Let (u, v) be an edge in \overline{G} . Then (u, v) is not an edge in G . Hence, since V' is a clique, u and v are not both in V' . Hence, at least one of u and v is in $V \setminus V'$.

Hence, $V \setminus V'$ is a vertex cover in \overline{G} .

Let us prove Property 2. We must prove that G has a clique of size K if and only if \overline{G} has a vertex cover of size $n - K$.

[\Rightarrow]

Assume $G = (V, E)$ has a clique V' of size K .

We will prove that $V \setminus V'$ is a vertex cover of size $n - K$.

Let (u, v) be an edge in \overline{G} . Then (u, v) is not an edge in G . Hence, since V' is a clique, u and v are not both in V' . Hence, at least one of u and v is in $V \setminus V'$.

Hence, $V \setminus V'$ is a vertex cover in \overline{G} .

And we have

$$|V \setminus V'| = |V| - |V'| = n - K.$$

$[\Leftarrow]$

Assume $\overline{G} = (V, \overline{E})$ has a vertex cover V^* of size $n - K$.

[\Leftarrow]

Assume $\overline{G} = (V, \overline{E})$ has a vertex cover V^* of size $n - K$.

We will prove that $V \setminus V^*$ is a clique of size K .

[\Leftarrow]

Assume $\overline{G} = (V, \overline{E})$ has a vertex cover V^* of size $n - K$.

We will prove that $V \setminus V^*$ is a clique of size K .

Let $u, v \in V \setminus V^*$ be two different vertices.

[\Leftarrow]

Assume $\overline{G} = (V, \overline{E})$ has a vertex cover V^* of size $n - K$.

We will prove that $V \setminus V^*$ is a clique of size K .

Let $u, v \in V \setminus V^*$ be two different vertices. If $\{u, v\}$ is an edge of \overline{G} , then V^* is not a vertex cover of \overline{G} and we get a contradiction.

[\Leftarrow]

Assume $\overline{G} = (V, \overline{E})$ has a vertex cover V^* of size $n - K$.

We will prove that $V \setminus V^*$ is a clique of size K .

Let $u, v \in V \setminus V^*$ be two different vertices. If $\{u, v\}$ is an edge of \overline{G} , then V^* is not a vertex cover of \overline{G} and we get a contradiction. Hence, $\{u, v\}$ is not an edge of \overline{G} .

[\Leftarrow]

Assume $\overline{G} = (V, \overline{E})$ has a vertex cover V^* of size $n - K$.

We will prove that $V \setminus V^*$ is a clique of size K .

Let $u, v \in V \setminus V^*$ be two different vertices. If $\{u, v\}$ is an edge of \overline{G} , then V^* is not a vertex cover of \overline{G} and we get a contradiction. Hence, $\{u, v\}$ is not an edge of \overline{G} . Hence, $\{u, v\}$ is an edge of G .

[\Leftarrow]

Assume $\overline{G} = (V, \overline{E})$ has a vertex cover V^* of size $n - K$.

We will prove that $V \setminus V^*$ is a clique of size K .

Let $u, v \in V \setminus V^*$ be two different vertices. If $\{u, v\}$ is an edge of \overline{G} , then V^* is not a vertex cover of \overline{G} and we get a contradiction. Hence, $\{u, v\}$ is not an edge of \overline{G} . Hence, $\{u, v\}$ is an edge of G .

We just showed that two arbitrary vertices in $V \setminus V^*$ are connected by an edge in G . Hence, $V \setminus V^*$ is a clique in G .

[\Leftarrow]

Assume $\overline{G} = (V, \overline{E})$ has a vertex cover V^* of size $n - K$.

We will prove that $V \setminus V^*$ is a clique of size K .

Let $u, v \in V \setminus V^*$ be two different vertices. If $\{u, v\}$ is an edge of \overline{G} , then V^* is not a vertex cover of \overline{G} and we get a contradiction. Hence, $\{u, v\}$ is not an edge of \overline{G} . Hence, $\{u, v\}$ is an edge of G .

We just showed that two arbitrary vertices in $V \setminus V^*$ are connected by an edge in G . Hence, $V \setminus V^*$ is a clique in G .

And we have

$$|V \setminus V^*| = |V| - |V^*| = n - (n - K) = K.$$



About 3SAT

$$\phi = (x_1 \vee \neg x_1 \vee \neg x_2) \wedge (x_2 \vee x_3 \vee x_4) \wedge (x_1 \vee x_3 \vee \neg x_4) \wedge (\neg x_1 \vee x_3 \vee \neg x_4)$$

About 3SAT

Consider a Boolean formula ϕ with variables x_1, x_2, \dots, x_n of the form

$$\phi = C_1 \wedge C_2 \wedge \dots \wedge C_m,$$

where each C_i is of the form

$$C_i = \ell_1^i \vee \ell_2^i \vee \ell_3^i.$$

Each ℓ_j^i is a variable or the negation of a variable. C_i is called a *clause* and ℓ_j^i is called a *literal*.

$$\phi = (x_1 \vee \neg x_1 \vee \neg x_2) \wedge (x_2 \vee x_3 \vee x_4) \wedge (x_1 \vee x_3 \vee \neg x_4) \wedge (\neg x_1 \vee x_3 \vee \neg x_4)$$

About 3SAT

Consider a Boolean formula ϕ with variables x_1, x_2, \dots, x_n of the form

$$\phi = C_1 \wedge C_2 \wedge \dots \wedge C_m,$$

where each C_i is of the form

$$C_i = \ell_1^i \vee \ell_2^i \vee \ell_3^i.$$

Each ℓ_j^i is a variable or the negation of a variable. C_i is called a *clause* and ℓ_j^i is called a *literal*.

$$\phi = (x_1 \vee \neg x_1 \vee \neg x_2) \wedge (x_2 \vee x_3 \vee x_4) \wedge (x_1 \vee x_3 \vee \neg x_4) \wedge (\neg x_1 \vee x_3 \vee \neg x_4)$$

We say that ϕ is *satisfiable* if there exists a truth value for each of x_1, x_2, \dots, x_n such that ϕ is true.

For the example, if $x_1 = 0$, $x_2 = 1$, $x_3 = 0$ and $x_4 = 0$, then $\phi = 1$, hence ϕ is satisfiable.

About 3SAT

$$3SAT = \{\phi \mid \phi \text{ is of the form above and } \phi \text{ is satisfiable}\}$$

About 3SAT

$$3SAT = \{\phi \mid \phi \text{ is of the form above and } \phi \text{ is satisfiable}\}$$

We want to show that

$$3SAT \leq_P INDEPENDENT-SET.$$

About 3SAT

$$3SAT = \{\phi \mid \phi \text{ is of the form above and } \phi \text{ is satisfiable}\}$$

We want to show that

$$3SAT \leq_P INDEPENDENT\ SET.$$

What is the connection between boolean formulas and graphs? We need a function f such that

About 3SAT

$$3SAT = \{\phi \mid \phi \text{ is of the form above and } \phi \text{ is satisfiable}\}$$

We want to show that

$$3SAT \leq_P INDEPENDENT-SET.$$

What is the connection between boolean formulas and graphs? We need a function f such that

- 1 f maps inputs for 3SAT to inputs for INDEPENDENT-SET.

$$f(\phi) = (G, K)$$

- 2 ϕ is satisfiable if and only if G has an independent set of size K .
- 3 Time to construct G , when given ϕ , is polynomial in the size of ϕ .

About 3SAT

$$3SAT = \{\phi \mid \phi \text{ is of the form above and } \phi \text{ is satisfiable}\}$$

We want to show that

$$3SAT \leq_P INDEPENDENT-SET.$$

What is the connection between boolean formulas and graphs? We need a function f such that

- ① f maps inputs for 3SAT to inputs for INDEPENDENT-SET.

$$f(\phi) = (G, K)$$

- ② ϕ is satisfiable if and only if G has an independent set of size K .
- ③ Time to construct G , when given ϕ , is polynomial in the size of ϕ .

Somehow, we have to “encode” a satisfiable formula ϕ as an independent set of size K in a graph.

Let ϕ be an input for 3SAT

$$\phi = C_1 \wedge C_2 \wedge \dots \wedge C_m,$$

each clause C_i is the disjunction (\vee) of 3 literals.




(G, K) is obtained as follows:

- $K = m$ (number of clauses)
- G has $3m$ vertices, one vertex for each literal.
 - For each clause, the literals in this clause form a triangle in G .
 - Additionally, there is an edge between any pair of opposite literals.


$$\phi = (\neg x \vee y \vee \neg z) \wedge (x \vee \neg y \vee z) \wedge (x \vee y \vee z)$$

 $\neg x$ x y $\neg z$ $\neg y$ z x y z


$$\phi = (\neg x \vee y \vee \neg z) \wedge (x \vee \neg y \vee z) \wedge (x \vee y \vee z)$$




$$\phi = (\neg x \vee y \vee \neg z) \wedge (x \vee \neg y \vee z) \wedge (x \vee y \vee z)$$



$$\phi = (\neg x \vee y \vee \neg z) \wedge (x \vee \neg y \vee z) \wedge (x \vee y \vee z)$$




$$\phi = (\neg x \vee y \vee \neg z) \wedge (x \vee \neg y \vee z) \wedge (x \vee y \vee z)$$



ϕ is satisfiable: take $x = 1$, $y = 0$ and $z = 0$


$$\phi = (\neg x \vee y \vee \neg z) \wedge (x \vee \neg y \vee z) \wedge (x \vee y \vee z)$$



ϕ is satisfiable: take $x = 1$, $y = 0$ and $z = 0$

- first clause: $\neg z = \text{TRUE}$, choose vertex 3


$$\phi = (\neg x \vee y \vee \neg z) \wedge (x \vee \neg y \vee z) \wedge (x \vee y \vee z)$$



ϕ is satisfiable: take $x = 1$, $y = 0$ and $z = 0$

- first clause: $\neg z = \text{TRUE}$, choose vertex 3
- second clause: $\neg y = \text{TRUE}$, choose vertex 5

$$\phi = (\neg x \vee y \vee \neg z) \wedge (x \vee \neg y \vee z) \wedge (x \vee y \vee z)$$



ϕ is satisfiable: take $x = 1$, $y = 0$ and $z = 0$

- first clause: $\neg z = \text{TRUE}$, choose vertex 3
- second clause: $\neg y = \text{TRUE}$, choose vertex 5
- third: $x = \text{TRUE}$, choose vertex 7

For the general case, we want to prove that if ϕ is satisfiable, then G has an independent set of size K .

\Rightarrow

For the general case, we want to prove that if ϕ is satisfiable, then G has an independent set of size K .

[\implies] Assume that ϕ is satisfiable. Then there is a truth assignment for the variables such that ϕ is true.

For the general case, we want to prove that if ϕ is satisfiable, then G has an independent set of size K .

[\implies] Assume that ϕ is satisfiable. Then there is a truth assignment for the variables such that ϕ is true.

Hence, every clause is true.

For the general case, we want to prove that if ϕ is satisfiable, then G has an independent set of size K .

[\implies] Assume that ϕ is satisfiable. Then there is a truth assignment for the variables such that ϕ is true.

Hence, every clause is true.

Hence, in every clause, at least one literal is true. Choose one literal which is true and pick the corresponding vertex.

For the general case, we want to prove that if ϕ is satisfiable, then G has an independent set of size K .

[\implies] Assume that ϕ is satisfiable. Then there is a truth assignment for the variables such that ϕ is true.

Hence, every clause is true.

Hence, in every clause, at least one literal is true. Choose one literal which is true and pick the corresponding vertex.

This gives $m = K$ vertices in G which form an independent set (do you see why?)

For the general case, we want to prove that if ϕ is satisfiable, then G has an independent set of size K .

[\implies] Assume that ϕ is satisfiable. Then there is a truth assignment for the variables such that ϕ is true.


Hence, every clause is true.

Hence, in every clause, at least one literal is true. Choose one literal which is true and pick the corresponding vertex.


This gives $m = K$ vertices in G which form an independent set (do you see why?)

Thus,

$$\phi \in 3SAT \implies (G, K) \in INDEPENDENT - SET.$$




In the example, take an independent set of size 3: $\{1, 6, 9\}$




In the example, take an independent set of size 3: $\{1, 6, 9\}$

- Vertex 1: set x such that $\neg x = 1$



In the example, take an independent set of size 3: $\{1, 6, 9\}$

- Vertex 1: set x such that $\neg x = 1$
- Vertex 6: set z such that $z = 1$



In the example, take an independent set of size 3: $\{1, 6, 9\}$

- Vertex 1: set x such that $\neg x = 1$
- Vertex 6: set z such that $z = 1$
- Vertex 9: set z such that $z = 1$

For the general case, we want to prove that if G has an independent set of size K , then ϕ is satisfiable.

[\Leftarrow]

For the general case, we want to prove that if G has an independent set of size K , then ϕ is satisfiable.

[\Leftarrow] Assume that G contains an independent set of size $K = m$.

For the general case, we want to prove that if G has an independent set of size K , then ϕ is satisfiable.

[\Leftarrow] Assume that G contains an independent set of size $K = m$.

Since G consists of m triangles, the independent set contains one vertex of each triangle.

For the general case, we want to prove that if G has an independent set of size K , then ϕ is satisfiable.

[\Leftarrow] Assume that G contains an independent set of size $K = m$.

Since G consists of m triangles, the independent set contains one vertex of each triangle.

For each triangle, let v be the vertex that is in the independent set.

- If v is of the form \boxed{x} , set $x = 1$.
- If v is of the form $\boxed{\neg x}$, set $x = 0$.

This way, (a subset of) the variables get a truth value such that $\phi = \text{TRUE}$.

For the general case, we want to prove that if G has an independent set of size K , then ϕ is satisfiable.

[\Leftarrow] Assume that G contains an independent set of size $K = m$.

Since G consists of m triangles, the independent set contains one vertex of each triangle.

For each triangle, let v be the vertex that is in the independent set.

- If v is of the form \boxed{x} , set $x = 1$.
- If v is of the form $\boxed{\neg x}$, set $x = 0$.

This way, (a subset of) the variables get a truth value such that $\phi = \text{TRUE}$.

Can the following happen? The vertices v and w are in the independent set and

- $v = \boxed{x}$ so we have to set $x = 1$.
- $w = \boxed{\neg x}$ so we have to set $x = 0$.

For the general case, we want to prove that if G has an independent set of size K , then ϕ is satisfiable.

[\Leftarrow] Assume that G contains an independent set of size $K = m$.

Since G consists of m triangles, the independent set contains one vertex of each triangle.

For each triangle, let v be the vertex that is in the independent set.

- If v is of the form \boxed{x} , set $x = 1$.
- If v is of the form $\boxed{\neg x}$, set $x = 0$.

This way, (a subset of) the variables get a truth value such that $\phi = \text{TRUE}$.

Can the following happen? The vertices v and w are in the independent set and

- $v = \boxed{x}$ so we have to set $x = 1$.
- $w = \boxed{\neg x}$ so we have to set $x = 0$.

No! Because v and w are connected by an edge in G .

For the general case, we want to prove that if G has an independent set of size K , then ϕ is satisfiable.

[\Leftarrow] Assume that G contains an independent set of size $K = m$.

Since G consists of m triangles, the independent set contains one vertex of each triangle.

For each triangle, let v be the vertex that is in the independent set.

- If v is of the form \boxed{x} , set $x = 1$.
- If v is of the form $\boxed{\neg x}$, set $x = 0$.

This way, (a subset of) the variables get a truth value such that $\phi = \text{TRUE}$.

Can the following happen? The vertices v and w are in the independent set and

- $v = \boxed{x}$ so we have to set $x = 1$.
- $w = \boxed{\neg x}$ so we have to set $x = 0$.

No! Because v and w are connected by an edge in G .

Thus,

$$(G, K) \in \text{INDEP} - \text{SET} \implies \phi \in 3\text{SAT}.$$

What is the time to compute $(G, K) = f(\phi)$? Using brute-force to compute the edges of G , we can do it in

$$O((3m)^2) = O(m^2) = O(\text{\#of clauses in } \phi)^2 = \text{polynomial time.}$$

