

# Topological Data Analysis: An Introduction to Persistent Homology

## Part 1: Inference and Stability

Ulrich Bauer



Sep 4, 2019

Machine learning summer school 2019  
Skoltech, Moscow



12



?




112




?


# Persistent homology






























## Gramicidin: an antibiotic functioning as ion channel





## Gramicidin: an antibiotic functioning as ion channel























# What is persistent homology?




# What is persistent homology?




# What is persistent homology?




# What is persistent homology?




# What is persistent homology?



# What is persistent homology?



# What is persistent homology?



# What is persistent homology?

## What is persistent homology?

Persistent homology is the homology of a filtration.

# What is persistent homology?

Persistent homology is the homology of a filtration.

- A filtration is a certain diagram  $K : \mathbb{R} \rightarrow \mathbf{Top}$  of topological spaces, indexed over the reals:

$$\dots \rightarrow K_s \hookrightarrow K_t \rightarrow \dots$$

- a topological space  $K_t$  for each  $t \in \mathbb{R}$
- an inclusion map  $K_s \hookrightarrow K_t$  for each  $s \leq t \in \mathbb{R}$

# What is persistent homology?

Persistent homology is the homology of a filtration.

- A filtration is a certain diagram  $K : \mathbb{R} \rightarrow \mathbf{Top}$  of topological spaces, indexed over the reals:

$$\dots \rightarrow K_s \hookrightarrow K_t \rightarrow \dots$$

- a topological space  $K_t$  for each  $t \in \mathbb{R}$
- an inclusion map  $K_s \hookrightarrow K_t$  for each  $s \leq t \in \mathbb{R}$
- Apply homology  $H_* : \mathbf{Top} \rightarrow \mathbf{Vect}$

# What is persistent homology?

Persistent homology is the homology of a filtration.

- A filtration is a certain diagram  $K : \mathbb{R} \rightarrow \mathbf{Top}$  of topological spaces, indexed over the reals:

$$\dots \rightarrow K_s \hookrightarrow K_t \rightarrow \dots$$

- a topological space  $K_t$  for each  $t \in \mathbb{R}$
- an inclusion map  $K_s \hookrightarrow K_t$  for each  $s \leq t \in \mathbb{R}$
- Apply homology  $H_* : \mathbf{Top} \rightarrow \mathbf{Vect}$
- Persistent homology is a diagram  $M : \mathbb{R} \rightarrow \mathbf{Vect}$  (*persistence module*).

In this talk, all vector spaces will be finite dimensional.

## Persistence modules

A *persistence module*  $M$  is a diagram  $M : \mathbb{R} \rightarrow \mathbf{vect}$  of vector spaces, indexed over the reals:

$$\dots \rightarrow M_s \longrightarrow M_t \dots \rightarrow$$

- a vector space  $M_t$  for each  $t \in \mathbb{R}$
- a linear map  $M_s \rightarrow M_t$  for each  $s \leq t \in \mathbb{R}$

## Persistence modules

A *persistence module*  $M$  is a diagram  $M : \mathbb{R} \rightarrow \mathbf{vect}$  of vector spaces, indexed over the reals:

$$\dots \rightarrow M_s \longrightarrow M_t \dots \rightarrow$$

- a vector space  $M_t$  for each  $t \in \mathbb{R}$
- an linear map  $M_s \rightarrow M_t$  for each  $s \leq t \in \mathbb{R}$

The maps (*morphisms*) between persistence modules are *natural transformations*:

$$\begin{array}{ccccc} \dots & \rightarrow & M_s & \longrightarrow & M_t & \dots & \rightarrow \\ & & f_s \downarrow & & \downarrow f_t & & \\ \dots & \rightarrow & N_s & \longrightarrow & N_t & \dots & \rightarrow \end{array}$$

- a linear map  $f_t : M_t \rightarrow N_t$  for each  $t \in \mathbb{R}$
- horizontal and vertical maps commute (for all  $s \leq t$ )

## Barcodes: the structure of persistence modules

Theorem (Crawley-Boevey 2015)

Any persistence module  $M : \mathbb{R} \rightarrow \mathbf{vect}$  (of finite dim. vector spaces over some field  $\mathbb{F}$ ) decomposes as a direct sum of interval modules

$$0 \rightarrow \cdots \rightarrow 0 \rightarrow \mathbb{F} \rightarrow \cdots \rightarrow \mathbb{F} \rightarrow 0 \rightarrow \cdots \rightarrow 0.$$

## Barcodes: the structure of persistence modules

Theorem (Crawley-Boevey 2015)

Any persistence module  $M : \mathbb{R} \rightarrow \text{vect}$  (of finite dim. vector spaces over some field  $\mathbb{F}$ ) decomposes as a direct sum of interval modules

$$0 \rightarrow \cdots \rightarrow 0 \rightarrow \mathbb{F} \rightarrow \cdots \rightarrow \mathbb{F} \rightarrow 0 \rightarrow \cdots \rightarrow 0.$$

- The supporting intervals form the *persistence barcode*  $B(M)$ .

## Barcodes: the structure of persistence modules

### Theorem (Crawley-Boevey 2015)

Any persistence module  $M : \mathbb{R} \rightarrow \mathbf{vect}$  (of finite dim. vector spaces over some field  $\mathbb{F}$ ) decomposes as a direct sum of interval modules

$$0 \rightarrow \cdots \rightarrow 0 \rightarrow \mathbb{F} \rightarrow \cdots \rightarrow \mathbb{F} \rightarrow 0 \rightarrow \cdots \rightarrow 0.$$

- The supporting intervals form the *persistence barcode*  $B(M)$ .
- The decomposition is not unique, but the barcode is.

## Barcodes: the structure of persistence modules

### Theorem (Crawley-Boevey 2015)

Any persistence module  $M : \mathbb{R} \rightarrow \mathbf{vect}$  (of finite dim. vector spaces over some field  $\mathbb{F}$ ) decomposes as a direct sum of interval modules

$$0 \rightarrow \cdots \rightarrow 0 \rightarrow \mathbb{F} \rightarrow \cdots \rightarrow \mathbb{F} \rightarrow 0 \rightarrow \cdots \rightarrow 0.$$

- The supporting intervals form the *persistence barcode*  $B(M)$ .
- The decomposition is not unique, but the barcode is.
- The barcode completely describes the persistence module (up to isomorphism).

## Barcodes: the structure of persistence modules

Theorem (Crawley-Boevey 2015)

Any persistence module  $M : \mathbb{R} \rightarrow \text{vect}$  (of finite dim. vector spaces over some field  $\mathbb{F}$ ) decomposes as a direct sum of interval modules

$$0 \rightarrow \cdots \rightarrow 0 \rightarrow \mathbb{F} \rightarrow \cdots \rightarrow \mathbb{F} \rightarrow 0 \rightarrow \cdots \rightarrow 0.$$

- The supporting intervals form the *persistence barcode*  $B(M)$ .
- The decomposition is not unique, but the barcode is.
- The barcode completely describes the persistence module (up to isomorphism).
- This is why we use homology with coefficients in a field.

## Barcodes: the structure of persistence modules

### Theorem (Crawley-Boevey 2015)

Any persistence module  $M : \mathbb{R} \rightarrow \mathbf{vect}$  (of finite dim. vector spaces over some field  $\mathbb{F}$ ) decomposes as a direct sum of interval modules

$$0 \rightarrow \cdots \rightarrow 0 \rightarrow \mathbb{F} \rightarrow \cdots \rightarrow \mathbb{F} \rightarrow 0 \rightarrow \cdots \rightarrow 0.$$

- The supporting intervals form the *persistence barcode*  $B(M)$ .
- The decomposition is not unique, but the barcode is.
- The barcode completely describes the persistence module (up to isomorphism).
- This is why we use homology with coefficients in a field.
- We rarely have a similarly clean structure for other parameter spaces, like  $\mathbb{R}^2 \rightarrow \mathbf{vect}$  (two-parameter persistence modules)

# Stability


# Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $f, g : X \rightarrow \mathbb{R}$  with  $\|f - g\|_\infty = \delta$  (and some regularity assumptions).

Consider the persistence barcodes of (sublevel set filtrations of)  $f$  and  $g$ .

Then there exists a matching between their intervals such that



# Stability of persistence barcodes for functions


Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $f, g : X \rightarrow \mathbb{R}$  with  $\|f - g\|_\infty = \delta$  (and some regularity assumptions).

Consider the persistence barcodes of (sublevel set filtrations of)  $f$  and  $g$ .

Then there exists a matching between their intervals such that

- matched intervals have endpoints within distance  $\leq \delta$ , and



# Stability of persistence barcodes for functions


Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $f, g : X \rightarrow \mathbb{R}$  with  $\|f - g\|_\infty = \delta$  (and some regularity assumptions).

Consider the persistence barcodes of (sublevel set filtrations of)  $f$  and  $g$ .

Then there exists a matching between their intervals such that

- matched intervals have endpoints within distance  $\leq \delta$ , and
- unmatched intervals have length  $\leq 2\delta$ .



# Homology inference

# Homology inference

Given: finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$

## Problem (Homology inference)

*Determine the homology  $H_*(X)$ .*

# Homology inference

Given: finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$

## Problem (Homology inference)

*Determine the homology  $H_*(X)$ .*

## Problem (Homological reconstruction)

*Construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .*

# Homology inference

Given: finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$

## Problem (Homology inference)

Determine the homology  $H_*(X)$ .

## Problem (Homological reconstruction)

Construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .

Approach:

- approximate the shape by a thickening  $P_\delta = \bigcup_{p \in P} B_\delta(p)$  covering  $X$

# Homology inference

Given: finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$

## Problem (Homology inference)

Determine the homology  $H_*(X)$ .

## Problem (Homological reconstruction)

Construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .

Approach:

- approximate the shape by a thickening  $P_\delta = \bigcup_{p \in P} B_\delta(p)$  covering  $X$

Requires strong assumptions:


## Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .




# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .




# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .




# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .




# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the induced map  $H_*(X \hookrightarrow X_\delta)$  is bijective, and
- the induced map  $H_*(X_\delta \hookrightarrow X_{2\delta})$  is injective.

Then  $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$ .



# Homology inference using persistence


Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the induced map  $H_*(X \hookrightarrow X_\delta)$  is bijective, and
- the induced map  $H_*(X_\delta \hookrightarrow X_{2\delta})$  is injective.

Then  $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$ .

Proof.



# Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the induced map  $H_*(X \hookrightarrow X_\delta)$  is bijective, and
- the induced map  $H_*(X_\delta \hookrightarrow X_{2\delta})$  is injective.

Then  $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$ .

Proof.

$$\begin{array}{ccccc} H_*(X) & & H_*(X_\delta) & & H_*(X_{2\delta}) \\ & \searrow & \nearrow & \searrow & \nearrow \\ & H_*(P_\delta) & & H_*(P_{2\delta}) & \end{array}$$

# Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the induced map  $H_*(X \hookrightarrow X_\delta)$  is bijective, and
- the induced map  $H_*(X_\delta \hookrightarrow X_{2\delta})$  is injective.

Then  $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$ .

Proof.

$$\begin{array}{ccccc} H_*(X) & \longleftrightarrow & H_*(X_\delta) & \hookrightarrow & H_*(X_{2\delta}) \\ & \searrow & \nearrow & \searrow & \nearrow \\ & & H_*(P_\delta) & & H_*(P_{2\delta}) \end{array}$$

# Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the induced map  $H_*(X \hookrightarrow X_\delta)$  is bijective, and
- the induced map  $H_*(X_\delta \hookrightarrow X_{2\delta})$  is injective.

Then  $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$ .

Proof.

$$\begin{array}{ccccc} H_*(X) & \longleftrightarrow & H_*(X_\delta) & \hookrightarrow & H_*(X_{2\delta}) \\ & \searrow & \nearrow & \searrow & \nearrow \\ & & H_*(P_\delta) & & H_*(P_{2\delta}) \end{array}$$

# Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the induced map  $H_*(X \hookrightarrow X_\delta)$  is bijective, and
- the induced map  $H_*(X_\delta \hookrightarrow X_{2\delta})$  is injective.

Then  $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$ .

Proof.

$$\begin{array}{ccccc} H_*(X) & \longleftrightarrow & H_*(X_\delta) & \hookrightarrow & H_*(X_{2\delta}) \\ & \searrow & \nearrow & \swarrow & \nearrow \\ & & H_*(P_\delta) & & H_*(P_{2\delta}) \end{array}$$

# Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the induced map  $H_*(X \hookrightarrow X_\delta)$  is bijective, and
- the induced map  $H_*(X_\delta \hookrightarrow X_{2\delta})$  is injective.

Then  $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$ .

Proof.

$$\begin{array}{ccccc} H_*(X) & \longleftrightarrow & H_*(X_\delta) & \hookrightarrow & H_*(X_{2\delta}) \\ & \searrow & \nearrow & \swarrow & \nearrow \\ & & H_*(P_\delta) & \longrightarrow & H_*(P_{2\delta}) \end{array}$$

# Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the induced map  $H_*(X \hookrightarrow X_\delta)$  is bijective, and
- the induced map  $H_*(X_\delta \hookrightarrow X_{2\delta})$  is injective.

Then  $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$ .

Proof.

$$\begin{array}{ccccc} H_*(X) & \longleftrightarrow & H_*(X_\delta) & \hookrightarrow & H_*(X_{2\delta}) \\ \searrow & & \nearrow & \uparrow \cong & \swarrow \\ H_*(P_\delta) & \xrightarrow{\quad} & H_*(P_{2\delta}) & \xleftarrow{\quad} & \\ \searrow & & \downarrow & \nearrow & \\ & & \text{im } H_*(P_\delta \hookrightarrow P_{2\delta}) & & \end{array}$$

# Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the induced map  $H_*(X \hookrightarrow X_\delta)$  is bijective, and
- the induced map  $H_*(X_\delta \hookrightarrow X_{2\delta})$  is injective.

Then  $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$ .

Proof.

$$\begin{array}{ccccc} H_*(X) & \xleftarrow{\cong} & H_*(X_\delta) & \hookrightarrow & H_*(X_{2\delta}) \\ & \searrow & \nearrow & \uparrow \cong & \swarrow \\ & H_*(P_\delta) & \xrightarrow{\quad} & H_*(P_{2\delta}) & \\ & \searrow & \downarrow & \nearrow & \\ & & \text{im } H_*(P_\delta \hookrightarrow P_{2\delta}) & & \end{array}$$

□

## Homological realization

This motivates the *homological realization problem*:

### Problem

Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that  $H_*(X) \cong \text{im } H_*(L \hookrightarrow K)$ :

$$H_*(L) \twoheadrightarrow H_*(X) \hookrightarrow H_*(K);$$

equivalently,  $\dim H_*(X) = \text{rank } H_*(L \hookrightarrow K)$ .

# Homological realization



This motivates the *homological realization problem*:

## Problem

Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that  $H_*(X) \cong \text{im } H_*(L \hookrightarrow K)$ :

$$H_*(L) \twoheadrightarrow H_*(X) \hookrightarrow H_*(K);$$

equivalently,  $\dim H_*(X) = \text{rank } H_*(L \hookrightarrow K)$ .



# Homological realization


This motivates the *homological realization problem*:

## Problem

Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that  $H_*(X) \cong \text{im } H_*(L \hookrightarrow K)$ :

$$H_*(L) \twoheadrightarrow H_*(X) \hookrightarrow H_*(K);$$

equivalently,  $\dim H_*(X) = \text{rank } H_*(L \hookrightarrow K)$ .



# Homological realization

This motivates the *homological realization problem*:


## Problem

Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that  $H_*(X) \cong \text{im } H_*(L \hookrightarrow K)$ :

$$H_*(L) \twoheadrightarrow H_*(X) \hookrightarrow H_*(K);$$

equivalently,  $\dim H_*(X) = \text{rank } H_*(L \hookrightarrow K)$ .

This is not always possible:



# Homological realization

This motivates the *homological realization problem*:


## Problem

Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that  $H_*(X) \cong \text{im } H_*(L \hookrightarrow K)$ :

$$H_*(L) \twoheadrightarrow H_*(X) \hookrightarrow H_*(K);$$

equivalently,  $\dim H_*(X) = \text{rank } H_*(L \hookrightarrow K)$ .

This is not always possible:



# Homological realization

This motivates the *homological realization problem*:


## Problem

Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that  $H_*(X) \cong \text{im } H_*(L \hookrightarrow K)$ :

$$H_*(L) \twoheadrightarrow H_*(X) \hookrightarrow H_*(K);$$

equivalently,  $\dim H_*(X) = \text{rank } H_*(L \hookrightarrow K)$ .

This is not always possible:



# Homological realization

This motivates the *homological realization problem*:


## Problem

Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that  $H_*(X) \cong \text{im } H_*(L \hookrightarrow K)$ :

$$H_*(L) \twoheadrightarrow H_*(X) \hookrightarrow H_*(K);$$

equivalently,  $\dim H_*(X) = \text{rank } H_*(L \hookrightarrow K)$ .

This is not always possible:



# Homological realization

This motivates the *homological realization problem*:


## Problem

Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that  $H_*(X) \cong \text{im } H_*(L \hookrightarrow K)$ :

$$H_*(L) \twoheadrightarrow H_*(X) \hookrightarrow H_*(K);$$

equivalently,  $\dim H_*(X) = \text{rank } H_*(L \hookrightarrow K)$ .


This is not always possible:




Theorem (Attali, B, Devillers, Glisse, Lieutier 2013)

The homological realization problem is NP-hard, even in  $\mathbb{R}^3$ .

# Simplification





## Sublevel set simplification

Let  $F_t = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

### Problem (Sublevel set simplification)

*Given a function  $f : X \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a function  $g$  with  $\|g - f\|_\infty \leq \delta$  minimizing  $\dim H_*(G_t)$ .*

## Sublevel set simplification

Let  $F_t = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

### Problem (Sublevel set simplification)

Given a function  $f : X \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a function  $g$  with  $\|g - f\|_\infty \leq \delta$  minimizing  $\dim H_*(G_t)$ .

- Since  $F_{t-\delta} \subseteq G_t \subseteq F_{t+\delta}$ , a lower bound is given by

$$\dim H_*(G_t) \geq \text{rank } H_*(F_{t-\delta} \hookrightarrow F_{t+\delta}).$$

## Sublevel set simplification

Let  $F_t = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

### Problem (Sublevel set simplification)

Given a function  $f : X \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a function  $g$  with  $\|g - f\|_\infty \leq \delta$  minimizing  $\dim H_*(G_t)$ .

- Since  $F_{t-\delta} \subseteq G_t \subseteq F_{t+\delta}$ , a lower bound is given by

$$\dim H_*(G_t) \geq \text{rank } H_*(F_{t-\delta} \hookrightarrow F_{t+\delta}).$$

- But it is NP-hard to decide whether this bound can be achieved.

## Sublevel set simplification

Let  $F_t = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

### Problem (Sublevel set simplification)

Given a function  $f : X \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a function  $g$  with  $\|g - f\|_\infty \leq \delta$  minimizing  $\dim H_*(G_t)$ .

- Since  $F_{t-\delta} \subseteq G_t \subseteq F_{t+\delta}$ , a lower bound is given by


$$\dim H_*(G_t) \geq \text{rank } H_*(F_{t-\delta} \hookrightarrow F_{t+\delta}).$$

- But it is NP-hard to decide whether this bound can be achieved.

### Theorem (Attali, B, Devillers, Glisse, Lieutier 2013)


Sublevel set simplification in  $\mathbb{R}^3$  is NP-hard.











# Topological simplification of functions

Consider the following problem:


## Problem (Topological simplification)

*Given a function  $f$  and a real number  $\delta \geq 0$ , find a function  $f_\delta$  subject to  $\|f_\delta - f\|_\infty \leq \delta$  with the minimal number of critical points.*

# Persistence and Morse theory

Morse theory (smooth or discrete):

- Relates critical points to homology of sublevel sets
- Provides a method for *cancelling* pairs of critical points




(from Milnor: *Lectures on the h-cobordism theorem*, 1965)

# Persistence and Morse theory

Morse theory (smooth or discrete):

- Relates critical points to homology of sublevel sets
- Provides a method for *cancelling* pairs of critical points



(from Milnor: *Lectures on the h-cobordism theorem*, 1965)

Persistent homology:

- Relates homology of different sublevel set
- Identifies pairs of critical points (birth and death of homological feature) and quantifies their *persistence*

# Persistence and discrete Morse theory

By stability of persistence barcodes:

## Proposition

*The intervals in the barcode of  $f$  with persistence  $> 2\delta$  provide a lower bound on the number of critical points of any function  $g$  with  $\|g - f\|_\infty \leq \delta$ .*

# Persistence and discrete Morse theory

By stability of persistence barcodes:

## Proposition

*The intervals in the barcode of  $f$  with persistence  $> 2\delta$  provide a lower bound on the number of critical points of any function  $g$  with  $\|g - f\|_\infty \leq \delta$ .*

## Theorem (B, Lange, Wardetzky, 2011)

*Let  $f$  be a function on a surface and let  $\delta > 0$ .*

*Then canceling all persistence pairs with persistence  $\leq 2\delta$  yields a function  $f_\delta$  with*

$$\|f_\delta - f\|_\infty \leq \delta$$

*that achieves the lower bound on the number of critical points.*

# Persistence and discrete Morse theory

By stability of persistence barcodes:

## Proposition

*The intervals in the barcode of  $f$  with persistence  $> 2\delta$  provide a lower bound on the number of critical points of any function  $g$  with  $\|g - f\|_\infty \leq \delta$ .*

## Theorem (B, Lange, Wardetzky, 2011)

*Let  $f$  be a function on a surface and let  $\delta > 0$ .*

*Then canceling all persistence pairs with persistence  $\leq 2\delta$  yields a function  $f_\delta$  with*

$$\|f_\delta - f\|_\infty \leq \delta$$

*that achieves the lower bound on the number of critical points.*

- Does not generalize to higher-dimensional manifolds!


# Persistence and stability: the big picture

Data


point cloud

$$P \subset \mathbb{R}^d$$


# Persistence and stability: the big picture




# Persistence and stability: the big picture




# Persistence and stability: the big picture




# Persistence and stability: the big picture



# Persistence and stability: the big picture




# Persistence and stability: the big picture



# The category of matchings

Consider the category **Mch** with

- objects: sets,
- morphisms: matchings (bijections between subsets).




# The category of matchings

Consider the category **Mch** with


- objects: sets,
- morphisms: matchings (bijections between subsets).

Composition:




## From barcodes to matching diagrams (and back)

- A barcode (collection of intervals) defines a diagram  $\mathbf{R} \rightarrow \mathbf{Mch}$ :



$t \mapsto \{\text{intervals in barcode containing } t\}$



$(s \leq t) \mapsto \{\text{intervals containing both } s, t\}$

## From barcodes to matching diagrams (and back)


- A barcode (collection of intervals) defines a diagram  $\mathbf{R} \rightarrow \mathbf{Mch}$ :



$t \mapsto \{\text{intervals in barcode containing } t\}$


$(s \leq t) \mapsto \{\text{intervals containing both } s, t\}$

- A matching diagram defines a barcode:




- intervals are formed by equivalence classes of matched elements


## Geometric interleavings




## Geometric interleavings




## Geometric interleavings




## Geometric interleavings



## Geometric interleavings



## Geometric interleavings




## Interleavings

Let  $\delta = \|f - g\|_\infty$ . Write  $F_t = f^{-1}(-\infty, t]$  for the  $t$ -sublevel set of  $f$ . Similarly,  $G_t = g^{-1}(-\infty, t]$

## Interleavings

Let  $\delta = \|f - g\|_\infty$ . Write  $F_t = f^{-1}(-\infty, t]$  for the  $t$ -sublevel set of  $f$ . Similarly,  $G_t = g^{-1}(-\infty, t]$


- The sublevel set filtrations  $F, G : \mathbb{R} \rightarrow \mathbf{Top}$  are  $\delta$ -interleaved:




## Interleavings

Let  $\delta = \|f - g\|_\infty$ . Write  $F_t = f^{-1}(-\infty, t]$  for the  $t$ -sublevel set of  $f$ . Similarly,  $G_t = g^{-1}(-\infty, t]$

- The sublevel set filtrations  $F, G : \mathbb{R} \rightarrow \mathbf{Top}$  are  $\delta$ -interleaved:



- Applying homology yields a  $\delta$ -interleaving of persistent homology (functors preserve commutativity):




# Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2012; B, Lesnick 2015)

If two persistence modules are  $\delta$ -interleaved,  
then there exists a  $\delta$ -matching of their barcodes:

- matched intervals have endpoints within distance  $\leq \delta$ ,
- unmatched intervals have length  $\leq 2\delta$ .




# Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2012; B, Lesnick 2015)

If two persistence modules are  $\delta$ -interleaved,  
then there exists a  $\delta$ -matching of their barcodes:

- matched intervals have endpoints within distance  $\leq \delta$ ,
- unmatched intervals have length  $\leq 2\delta$ .



Equivalently: there exists a  $\delta$ -interleaving of their barcodes (as diagrams  $\mathbf{R} \rightarrow \mathbf{Mch}$ ).

## Non-functoriality of persistence barcodes

Can a persistence module  $M$  be mapped to its barcode  $B(M)$  by a functor  $B : \mathbf{vect} \rightarrow \mathbf{Mch}$ ?

- This would preserve  $\delta$ -interleavings, and thus yield stability of persistence barcodes.

# Non-functoriality of persistence barcodes

Can a persistence module  $M$  be mapped to its barcode  $B(M)$  by a functor  $B : \mathbf{vect} \rightarrow \mathbf{Mch}$ ?

- This would preserve  $\delta$ -interleavings, and thus yield stability of persistence barcodes.



# Non-functoriality of persistence barcodes

Can a persistence module  $M$  be mapped to its barcode  $B(M)$  by a functor  $B : \mathbf{vect} \rightarrow \mathbf{Mch}$ ?

- This would preserve  $\delta$ -interleavings, and thus yield stability of persistence barcodes.



## Proposition

*There is no functor  $\mathbf{vect} \rightarrow \mathbf{Mch}$  sending a  $d$ -dimensional vector space to a set with  $d$  elements.*

# Non-functoriality of persistence barcodes

Can a persistence module  $M$  be mapped to its barcode  $B(M)$  by a functor  $B : \mathbf{vect} \rightarrow \mathbf{Mch}$ ?

- This would preserve  $\delta$ -interleavings, and thus yield stability of persistence barcodes.



## Proposition

*There is no functor  $\mathbf{vect} \rightarrow \mathbf{Mch}$  sending a  $d$ -dimensional vector space to a set with  $d$  elements.*

- In other words, there is no canonical way of assigning a basis to a vector space.

## Structure of persistence sub-/quotient modules

### Proposition

Let  $f : M \twoheadrightarrow N$  be an epimorphism of persistence modules:

- a surjective map  $f_t : M_t \twoheadrightarrow N_t$  for each  $t \in \mathbb{R}$  (commuting with horizontal maps)

$$\begin{array}{ccccccc} \dots & \rightarrow & M_s & \longrightarrow & M_t & \dots & \rightarrow \\ & & f_s \downarrow & & \downarrow f_t & & \\ \dots & \rightarrow & N_s & \longrightarrow & N_t & \dots & \rightarrow \end{array}$$

# Structure of persistence sub-/quotient modules

## Proposition

Let  $f : M \twoheadrightarrow N$  be an epimorphism of persistence modules:

- a surjective map  $f_t : M_t \twoheadrightarrow N_t$  for each  $t \in \mathbb{R}$  (commuting with horizontal maps)

$$\begin{array}{ccccccc} \dots & \rightarrow & M_s & \longrightarrow & M_t & \dots & \rightarrow \\ & & f_s \downarrow & & \downarrow f_t & & \\ \dots & \rightarrow & N_s & \longrightarrow & N_t & \dots & \rightarrow \end{array}$$

Then there is an injection of barcodes  $\chi(f) : B(N) \hookrightarrow B(M)$ .

If  $\chi(f)$  maps  $J$  to  $I$ , then

- $I$  and  $J$  are aligned below, and
- $I$  bounds  $J$  above.



# Structure of persistence sub-/quotient modules

## Proposition

Let  $f : M \twoheadrightarrow N$  be an epimorphism of persistence modules:

- a surjective map  $f_t : M_t \twoheadrightarrow N_t$  for each  $t \in \mathbb{R}$  (commuting with horizontal maps)

$$\begin{array}{ccccccc} \dots & \rightarrow & M_s & \longrightarrow & M_t & \dots & \rightarrow \\ & & f_s \downarrow & & \downarrow f_t & & \\ \dots & \rightarrow & N_s & \longrightarrow & N_t & \dots & \rightarrow \end{array}$$

Then there is an injection of barcodes  $\chi(f) : B(N) \hookrightarrow B(M)$ .

If  $\chi(f)$  maps  $J$  to  $I$ , then

- $I$  and  $J$  are aligned below, and
- $I$  bounds  $J$  above.



This construction respects composition:  $\chi(g \circ f) = \chi(g) \circ \chi(f)$

# Structure of persistence sub-/quotient modules

## Proposition

Let  $f : M \twoheadrightarrow N$  be an epimorphism of persistence modules:

- a surjective map  $f_t : M_t \twoheadrightarrow N_t$  for each  $t \in \mathbb{R}$  (commuting with horizontal maps)

$$\begin{array}{ccccccc} \dots & \rightarrow & M_s & \longrightarrow & M_t & \dots & \rightarrow \\ & & f_s \downarrow & & \downarrow f_t & & \\ \dots & \rightarrow & N_s & \longrightarrow & N_t & \dots & \rightarrow \end{array}$$

Then there is an injection of barcodes  $\chi(f) : B(N) \hookrightarrow B(M)$ .

If  $\chi(f)$  maps  $J$  to  $I$ , then

- $I$  and  $J$  are aligned below, and
- $I$  bounds  $J$  above.



This construction respects composition:  $\chi(g \circ f) = \chi(g) \circ \chi(f)$

Dually, there is an injection  $B(M) \hookrightarrow B(N)$  for monomorphisms  $M \hookrightarrow N$ .

## Induced matchings

For  $f : M \rightarrow N$  a morphism of pfd persistence modules, we always have a factorization through the image,

$$M \twoheadrightarrow \text{im } f \hookrightarrow N.$$

This gives an *induced matching*  $\chi(f)$  between their barcodes:

## Induced matchings


For  $f : M \rightarrow N$  a morphism of pfd persistence modules, we always have a factorization through the image,

$$M \twoheadrightarrow \text{im } f \hookrightarrow N.$$

This gives an *induced matching*  $\chi(f)$  between their barcodes:


- compose the injections  $B(M) \hookrightarrow B(\text{im } f) \hookrightarrow B(N)$  for epimorphisms and monomorphisms to a matching

$$\chi(f) : B(M) \not\rightarrow B(N).$$




## Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):




## Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):



## Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):




Induced  $\delta$ -matching of barcodes:

$$\underline{\hspace{1cm}} \qquad B(N)$$

- Left endpoints of matched intervals in  $B(M), B(N)$  are within distance  $\delta$ .

## Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):




Induced  $\delta$ -matching of barcodes:



- Left endpoints of matched intervals in  $B(M), B(N)$  are within distance  $\delta$ .

## Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):




Induced  $\delta$ -matching of barcodes:




- Left endpoints of matched intervals in  $B(M), B(N)$  are within distance  $\delta$ .

# Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):




Induced  $\delta$ -matching of barcodes:




- Left endpoints of matched intervals in  $B(M)$ ,  $B(N)$  are within distance  $\delta$ .

# Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):



Induced  $\delta$ -matching of barcodes:



$B(M)$   
 $B(\text{im } n(-\delta, \delta))$   
 $B(\text{im } f)$   
 $B(N(\delta))$   
 $B(N)$

- Left endpoints of matched intervals in  $B(M), B(N)$  are within distance  $\delta$ .


# Machine learning with persistence diagrams

## Extending the TDA pipeline


Mapping barcodes into a Hilbert space:

- desirable for machine learning methods (kernel methods, neural network) and statistics
- stability (Lipschitz continuity): important for reliable predictions
- inverse stability (bi-Lipschitz): avoid loss of information


# Persistence barcodes and persistence diagrams




## Persistence scale space kernel




## Persistence scale space kernel




## Persistence scale space kernel




## Persistence scale space kernel




## Persistence scale space kernel




## Persistence scale space kernel



## Persistence scale space kernel



## Persistence scale space kernel



## Metric distortion of the PSS kernel

Let  $d_{\text{PSS}}(D_f, D_g)$  be the  $L^2$  distance between smoothings of the persistence diagrams  $D_f, D_g$  (i.e., a *kernel distance*).

**Theorem (Reininghaus, Huber, B, Kwitt 2015)**

For two persistence diagrams  $D_f$  and  $D_g$  and  $\sigma > 0$ , we have

$$d_{\text{PSS}}(D_f, D_g) \leq \frac{C}{\sigma} d_1(D_f, D_g).$$

## Metric distortion of the PSS kernel

Let  $d_{\text{PSS}}(D_f, D_g)$  be the  $L^2$  distance between smoothings of the persistence diagrams  $D_f, D_g$  (i.e., a *kernel distance*).

**Theorem (Reininghaus, Huber, B, Kwitt 2015)**

For two persistence diagrams  $D_f$  and  $D_g$  and  $\sigma > 0$ , we have

$$d_{\text{PSS}}(D_f, D_g) \leq \frac{C}{\sigma} d_1(D_f, D_g).$$

- Here: stability with respect to *Wasserstein distance*:

$$d_p(D_f, D_g) = \left( \inf_{\mu: D_f \leftrightarrow D_g} \sum_{x \in D_f} \|x - \mu(x)\|_\infty^p \right)^{\frac{1}{p}}$$

- Note: bottleneck distance  $d_B(D_f, D_g) = \lim_{p \rightarrow \infty} d_p(D_f, D_g)$

## No bi-Lipschitz feature maps for persistence

Existing kernels for persistence diagrams in the literature:

- stability bounds only for 1–Wasserstein distance
- Lipschitz constants depend on bound on number and range of bars
- no bi-Lipschitz bounds known

## No bi-Lipschitz feature maps for persistence

Existing kernels for persistence diagrams in the literature:

- stability bounds only for 1–Wasserstein distance
- Lipschitz constants depend on bound on number and range of bars
- no bi-Lipschitz bounds known

Can we hope for something better?

## No bi-Lipschitz feature maps for persistence

Existing kernels for persistence diagrams in the literature:

- stability bounds only for 1–Wasserstein distance
- Lipschitz constants depend on bound on number and range of bars
- no bi-Lipschitz bounds known

Can we hope for something better?

### Theorem (B, Carrière 2018)

*There is no bi-Lipschitz map from the persistence diagrams (bottleneck or Wasserstein distance) into any Euclidean space  $\mathbb{R}^d$ , even when restricting to a bounded range or number of bars.*

## No bi-Lipschitz feature maps for persistence

Existing kernels for persistence diagrams in the literature:

- stability bounds only for 1–Wasserstein distance
- Lipschitz constants depend on bound on number and range of bars
- no bi-Lipschitz bounds known

Can we hope for something better?

### Theorem (B, Carrière 2018)

*There is no bi-Lipschitz map from the persistence diagrams (bottleneck or Wasserstein distance) into any Euclidean space  $\mathbb{R}^d$ , even when restricting to a bounded range or number of bars.*

### Theorem (B, Carrière 2018)

*If there was such a bi-Lipschitz map into any Hilbert space, the Lipschitz constant would have to go to  $\infty$  when the restrictions on the bars are lifted.*

Ripser

## Vietoris–Rips complexes

Consider a finite metric space  $(X, d)$  (given as a distance matrix).

The *Vietoris–Rips complex* is the simplicial complex

$$\text{Rips}_t(X) = \{S \subseteq X \mid \text{diam } S \leq t\}$$

- all edges with pairwise distance  $\leq t$
- all possible higher simplices

## An example computation

Example data set:

- 192 points on  $\mathbb{S}^2$
- persistent homology barcodes up to dimension 2
- over 56 mio. simplices in 3-skeleton

## An example computation

Example data set:

- 192 points on  $\mathbb{S}^2$
- persistent homology barcodes up to dimension 2
- over 56 mio. simplices in 3-skeleton

Some previous software:

- javaplex (Stanford): 3200 seconds, 12 GB
- Dionysus (Duke): 615 seconds, 3.4 GB
- DIPHA (IST Austria): 50 seconds, 6 GB
- GUDHI (INRIA): 60 seconds, 3 GB

## An example computation

Example data set:

- 192 points on  $\mathbb{S}^2$
- persistent homology barcodes up to dimension 2
- over 56 mio. simplices in 3-skeleton

Some previous software:

- javaplex (Stanford): 3200 seconds, 12 GB
- Dionysus (Duke): 615 seconds, 3.4 GB
- DIPHA (IST Austria): 50 seconds, 6 GB
- GUDHI (INRIA): 60 seconds, 3 GB

Demo: [live.ripser.org](http://live.ripser.org)

# An example computation

Example data set:

- 192 points on  $\mathbb{S}^2$
- persistent homology barcodes up to dimension 2
- over 56 mio. simplices in 3-skeleton

Some previous software:

- javaplex (Stanford): 3200 seconds, 12 GB
- Dionysus (Duke): 615 seconds, 3.4 GB
- DIPHA (IST Austria): 50 seconds, 6 GB
- GUDHI (INRIA): 60 seconds, 3 GB

Demo: [live.ripser.org](http://live.ripser.org)

- Ripser: 1.2 seconds, 160 MB

## Another example data set



(Columbia Object Image Library)