MLW / KUHeS Statistics and R short course

Session 5 - Practical

Marc Henrion

8 November 2022

Session 5 - Practical

Exercise 1

Using the adolescent_small.csv data from Session 4, fit a linear model regressing weight (variable a104wt) on age (variable a12age).

Test if the regression coefficient of age $\beta_{age} = 0$.

Note:

- deviance = sum of squares
- residual = error

Exercise 2

Repeat the ANOVA test from Session 4 as a linear regression model. Do you get the same results?

Note: glm will have the word deviance in its output. This is just another word for sum of squares.

Recall from Session 4:

aov(cd41~as.factor(hosp),data=Tbreg)

Exercise 3

Using the adolescent_small.csv data, fit the following GLM model:

Weight a104wt as a function of

- age a12age
- height a103ht
- hiv hiv
- sex a13sex

Produce:

- a residuals vs. fitted values graph
- histogram of the residuals
- a QQ plot.

Exercise 4

You are given the following data:

$$\mathbf{x} = (-6, -6, -4, -1, 0.5, 2, 8, 8, 11, 11.5)^T$$

$$\mathbf{y} = (-3.7, -4.3, -3.9, -4.6, 0.5, -6.9, 10.2, 16.1, 6, 19.5)^T$$

- a. Fit a linear regression model to these data and show the model output.
- b. Describe the resulting regression line:
 - What is the relationship between variables X and Y?
 - How much (on average) does Y change when X changes by 1?
 - What value does Y take (on average) when X = 0?
- c. Compute the coefficient of determination R^2 , the adjusted R^2 , the likelihood and the AIC. Which of these tell you how good your model fits the data?
- d. Compute the residuals $r_i = y_i \hat{y}_i$ and do a normal distribution QQ plot.
- e. What other diagnostic check(s) could you do? Do this and explain whether you think this is a good model.
- f. Re-fit the model, but now including a term for X^2 : $Y = \beta_0 + \beta_1 X + \beta_2 X^2 + \epsilon$. Check and discuss the resulting model and compare it to the previous one. Which model would you recommend for this dataset?

Exercise 5

Download (from GitHub) and load the dataset cuse.csv.

This is a dataset on contraceptive use. using, notUsing lists how many people in each group implied by combinations of age, education, wantsMore are currently using contraceptives. age, education are self-explanatory. wantsMore lists whether individuals want more children or not.

Model the binary variable specified by the 2 columns using, notUsing in terms of age, education, wantsMore.

- Discuss your results.
- What can you say about the deviance? Does it look like this is a good model?
- What happens if you include an interaction term between the age variable and the desire for more children variable?