

MATH 4720 / MSCS 5720

Instructor: Mehdi Maadooliat

Chapter 6 (Part C)



Department of Mathematics, Statistics and Computer Science

CHAPTER 6 (PART C)

- **Comparing Two Population Means**
 - Independent Samples
 - Dependent Samples
- **Two sample t-test (Independent Samples)**
 - Pooled t-test
 - Unequal variance t-test
- **Paired t-test (Dependent Samples)**
- **Power Analysis**
 - Independent Samples
 - Dependent Samples
- **Non-parametric Tests**
 - Sign test (from Chapter 5, test for median M)
 - Wilcoxon Rank-Sum (or Mann-Whitney) Test (two independent samples)
 - Wilcoxon Signed-Rank Test (dependent samples)

NONPARAMETRIC INFERENCE

- In both one-sample and two-sample t-tests, we assumed that either the sample size is ≥ 30 or the samples are drawn from normal populations.
- What if $n < 30$, and the distribution is **non-normal**?
- In such cases, we usually use **non-parametric** tests.
- No assumptions on the distribution means no parameters.

MOTIVATION

- **Example: Suppose the weights of cereal boxes is not normally distributed.**
- **Median weight of cereal boxes supposed to be 16.37 oz.**
- **Take a sample of 5 boxes: 16.01, 15.98, 16.23, 15.5, 16.2**
- **What is the probability that all of these boxes have weight less than 16.37 oz.?**
 - Ans. $\frac{1}{2^5} = 0.0315$.
- **This is the *p – value*.**
- **Note that to answer this, we did not need distributional assumption.**

EXAMPLE (CONT'D)

- **Now if the sample is:** 16.01, 15.98, 16.23, 15.5, 16.47
- **Here one 4 out of five values are less than 16.37 oz.**
- $H_0: \text{median} = 16.37$
- $H_a: \text{median} < 16.37$
- $p - \text{value}$
- $$= P \left(\begin{array}{c} \text{four or more values are less than 16.37} \\ \text{if } H_0 \text{ is true} \end{array} \right)$$
- $$= P(Y \geq 4), \quad Y \sim \text{Binomial}(n = 5, \pi = 0.5)$$
$$= 0.1875$$
- [Binomial Calculator](#)



NON-PARAMETRIC ONE SAMPLE INFERENCE (SECTION 5.9)



- **Sign Test**

Data: y_1, y_2, \dots, y_n

- $H_0: \text{median} = m_0$
- $H_a: \text{median} > m_0$
or $\text{median} < m_0$
or $\text{median} \neq m_0$
- **Test Statistics** $B = \# \text{ of data values} > m_0$
- **Decision Rule:**
 - $H_a: \text{median} > m_0$: **Reject H_0 in favor of H_a if $B \geq n - B_\alpha$**
 - $H_a: \text{median} < m_0$: **Reject H_0 in favor of H_a if $B \leq B_\alpha$**
 - $H_a: \text{median} \neq m_0$: **Reject H_0 in favor of H_a if $B \leq B_{\alpha/2}$ or $B \geq n - B_{\alpha/2}$**

BOOK EXAMPLE 5.20

- A landfill company wants to determine if the average weekly amount of household recyclable wastes material is more than 5 lbs. The data is collected from 25 households.



- Here, the word “average” should not be interpreted literally.
- [Minitab](#)

EXAMPLE CONT'D

- $n = 25 < 30$, and the data is **not normally** distributed.
- $H_0: \text{median} = 5$ vs. $H_a: \text{median} > 5$
- **Stat \rightarrow Nonparametrics \rightarrow 1- Sample Sign**

Sign Test for Median: Weekly Weight

Sign test of median = 5.000 versus > 5.000

	N	Below	Equal	Above	P	Median
Weekly Weight	25	12	0	13	0.5000	5.300

- T.S. $B = \#$ of data values greater 5 lbs = 13
- p-value is 0.5000, we **fail to reject** H_0 in favor of H_a .
- Thus we **cannot** conclude that the median household recyclable waste is greater than 5 pounds per week.

EXAMPLE CONT'D

- **Minitab Output**
- **95% Confidence Interval for the Median**

Sign CI: Weekly Weight

Sign confidence interval for median

			Achieved	Confidence		Position
			Confidence	Lower	Upper	
	N	Median				
Weekly Weight	25	5.300	0.8922	4.200	6.700	9
			0.9500	3.959	6.700	NLI
			0.9567	3.900	6.700	8

NON-PARAMETRIC TWO SAMPLES TEST

- Non-parametric Two **Independent** Samples Test

Group 1

$y_{11}, y_{12}, \dots, y_{1n_1}$

Group 2

$y_{21}, y_{22}, \dots, y_{2n_2}$

- $n_1 < 30$ **and/or** $n_2 < 30$
- Data is generated from **non-normal** distributions.





WILCOXON RANK-SUM TEST (MANN-WHITNEY U TEST)

- H_0 : Two distributions are identical (i. e. $med_1 = med_2$)
- H_a : dist. of 1 is shifted to the right of dist. 2 ($med_1 > med_2$)
- **or** dist. of 1 is shifted to the left of dist. 2 ($med_1 < med_2$)
- **or** Two distributions are not identical ($med_1 \neq med_2$)
- **T.S. Combine both the samples. Rank all values of the combined sample from lowest to the highest.**

T = sum of the ranks in sample 1

- **Decision Rule:**(We will use computer output)
 - $H_a: med_1 > med_2$: **Reject H_0 if $T > T_U$**
 - $H_a: med_1 < med_2$: **Reject H_0 if $T < T_L$**
 - $H_a: med_1 \neq med_2$: **Reject H_0 if $T > T_U^*$ or $T < T_L^*$**

A SIMPLE EXAMPLE

- $H_0: med_1 = med_2$

$$H_a: med_1 > med_2$$



- | | Group 1 | | | | | Group 2 | | | | |
|--------|--------------------|-----|---|---|----|--------------------|---|---|-----|---|
| | 32, 33, 55, 60, 61 | | | | | 23, 25, 56, 33, 21 | | | | |
| • Rank | 4 | 5.5 | 7 | 9 | 10 | 2 | 3 | 8 | 5.5 | 1 |
- **T.S.** $T =$ sum of the ranks of sample 1 $= 35.5$
 - If $T > T_U$, we would say that **dist. of group 1** is to the right of **dist. of group 2**
 - Note that T_U is determined in such a way that probability of false conclusion is $\alpha = 0.05$.

BOOK EXAMPLE 6.5

- An investigator is interested to study the effect of alcohol on reaction time. The following data is collected on the reaction time to an instruction.

Group 1: Alcohol
10 subjects

Group 2: Placebo
10 subjects



- [Minitab](#)

EXAMPLE 6.5: (CONT'D)

- $n_1 = 10$, $n_2 = 10$, and the distributions are **non-normal**
- $H_0: med_1 = med_2$ vs. $H_a: med_1 > med_2$
- **Minitab Output**

Stat → Nonparametrics → Mann-Whitney

Mann-Whitney Test and CI: Alcohol, Placebo

	N	Median
Alcohol	10	1.445
Placebo	10	0.845

Point estimate for ETA1-ETA2 is 0.610

95.5 Percent CI for ETA1-ETA2 is (0.250,1.080)

W = 140.0

Test of ETA1 = ETA2 vs ETA1 > ETA2 is significant at 0.0046

- **Conclusion:** Since p – value of 0.0046 is **small**, we reject H_0 in favor of H_a . Thus, we conclude that the reaction time for the Alcohol population is significantly higher than that for the Placebo population.

NON-PARAMETRIC TWO DEPENDENT SAMPLE TEST

Subject	y_1	y_2	$d = y_1 - y_2$
1	y_{11}	y_{21}	d_1
2	y_{12}	y_{22}	d_2
.	.	.	.
.	.	.	.
n	y_{1n}	y_{2n}	d_n

- $n < 30$ and the **differences are not normally distributed**
- In such case, a **nonparametric** method must be used.

WILCOXON SIGNED-RANK TEST

- $H_0: med_d = 0$ (**median of the difference = 0**)
- $H_a: med_d > 0$
 or $med_d < 0$
 or $med_d \neq 0$
- **T.S. Rank the absolute values of the differences.**
 - T_- = **sum of the ranks of negative differences**
 - T_+ = **sum of the ranks of positive differences**
 - T = **smaller of T_+ and T_-**
- **Decision Rule: (We will use computer output)**
 - $H_a: med_d > 0$: **Reject H_0 if $T_- < T_U^-$**
 - $H_a: med_d < 0$: **Reject H_0 if $T_+ < T_L^+$**
 - $H_a: med_d \neq 0$: **Reject H_0 if $T < T^*$**

A SIMPLE EXAMPLE

Subject	y_1	y_2	$d = y_1 - y_2$	Rank of $ y_1 - y_2 $
1	30	24	6	5
2	20	22	-2	1.5
3	32	30	2	1.5
4	41	37	4	4
5	27	30	-3	3

- **Test Statistics:**
- $T_- = 4.5$
- $T_+ = 10.5$
- $T = \text{smaller}(10.5, 4.5) = 4.5$

BOOK EXAMPLE 6.9



- Does Brand A fertilizer produce more grass than Brand B?

Field	Brand A	Brand B	Difference	Field	Brand A	Brand B	Difference
1	211.4	186.3	25.1	11	208.9	183.6	25.3
2	204.4	205.7	-1.3	12	208.7	188.7	20.0
3	202.0	184.4	17.6	13	213.8	188.6	25.2
4	201.9	203.6	-1.7	14	201.6	204.2	-2.6
5	202.4	180.4	22.0	15	201.8	181.6	20.1
6	202.0	202.0	0	16	200.3	208.7	-8.4
7	202.4	181.5	20.9	17	201.8	181.5	20.3
8	207.1	186.7	20.4	18	201.5	208.7	-7.2
9	203.6	205.7	-2.1	19	212.1	186.8	25.3
10	216.0	189.1	26.9	20	203.4	182.9	20.5

- Minitab



BOOK EXAMPLE 6.9 (CONT'D)



- $H_0: med_d = 0$ vs $H_a: med_d > 0$

BOOK EXAMPLE 6.9 (CONT'D)

- **Minitab Output**

- Stat → Nonparametrics → 1-Sample Wilcoxon

Wilcoxon Signed Rank Test: Difference

Test of median = 0.000000 versus median > 0.000000

	N	N for Test	Wilcoxon Statistic	P	Estimated Median
Difference	20	19	169.0	0.002	11.80

- **Conclusion:** Since the $p - value = 0.002$ is **small**, we reject H_0 in favor of H_a . Thus conclude that Brand A fertilizer produce more grass than Brand B.
- **95% Confidence interval**

Wilcoxon Signed Rank CI: Difference

	N	Estimated Median	Achieved Confidence	Confidence Interval Lower	Confidence Interval Upper
Difference	20	11.8	95.0	8.7	21.4