

Deep Learning

Lecture 3: Calculus

Dr. Mehrdad Maleki

A function $f : X \rightarrow Y$ is given.

A function $f : X \rightarrow Y$ is given. Then the derivative of f at $x_0 \in X$ means that if x_0 changes a little bit, how $f(x_0)$ changes accordingly.

A function $f : X \rightarrow Y$ is given. Then the derivative of f at $x_0 \in X$ means that if x_0 changes a little bit, how $f(x_0)$ changes accordingly. For example if the radius of a balloon of radius 50 cm change to 55 cm, how much change we have in the volume of the baloon?

A function $f : X \rightarrow Y$ is given. Then the derivative of f at $x_0 \in X$ means that if x_0 changes a little bit, how $f(x_0)$ changes accordingly. For example if the radius of a balloon of radius 50 cm change to 55 cm, how much change we have in the volume of the baloon? Since $V = \frac{4}{3}\pi R^3$ then $\frac{dV}{dR} = 4\pi R^2$.

A function $f : X \rightarrow Y$ is given. Then the derivative of f at $x_0 \in X$ means that if x_0 changes a little bit, how $f(x_0)$ changes accordingly. For example if the radius of a balloon of radius 50 cm change to 55 cm, how much change we have in the volume of the baloon? Since $V = \frac{4}{3}\pi R^3$ then $\frac{dV}{dR} = 4\pi R^2$. So $V_2 - V_1 \approx \frac{dV}{dR}(R_2 - R_1)$.

A function $f : X \rightarrow Y$ is given. Then the derivative of f at $x_0 \in X$ means that if x_0 changes a little bit, how $f(x_0)$ changes accordingly. For example if the radius of a balloon of radius 50 cm change to 55 cm, how much change we have in the volume of the baloon? Since $V = \frac{4}{3}\pi R^3$ then $\frac{dV}{dR} = 4\pi R^2$. So


$$V_2 - V_1 \approx \frac{dV}{dR}(R_2 - R_1). \text{ Since } R_2 - R_1 = 5 \text{ and}$$

$$\frac{dV}{dR}|_{R=50} = 4\pi \times 50^2 \approx 31400. \text{ So}$$


$$V_2 - V_1 \approx 31400 * 5 = 157000 \text{ cm}^3 \text{ but } V_1 \approx 52333 \text{ and}$$

$$V_2 \approx 157000 + 52333 \text{ thus } \frac{V_2}{V_1} = \frac{209333}{52333} \approx 4. \text{ Hence, the volume of the sphere increases by a factor of 4 if the radius of the sphere is increased by 5 cm.}$$


In one dimensional case when we have a scalar function of one variable $y = f(x)$ we can plot this on the $x - y$ coordinate and the slope of the graph of the function at point x_0 is the derivative of f at x_0 and it is denoted by $f'(x_0)$ or $\frac{dy}{dx}|_{x=x_0}$.




In one dimensional case when we have a scalar function of one variable $y = f(x)$ we can plot this on the $x - y$ coordinate and the slope of the graph of the function at point x_0 is the derivative of f at x_0 and it is denoted by $f'(x_0)$ or $\frac{dy}{dx}\big|_{x=x_0}$.




In one dimensional case when we have a scalar function of one variable $y = f(x)$ we can plot this on the $x - y$ coordinate and the slope of the graph of the function at point x_0 is the derivative of f at x_0 and it is denoted by $f'(x_0)$ or $\frac{dy}{dx}|_{x=x_0}$.




In one dimensional case when we have a scalar function of one variable $y = f(x)$ we can plot this on the $x - y$ coordinate and the slope of the graph of the function at point x_0 is the derivative of f at x_0 and it is denoted by $f'(x_0)$ or $\frac{dy}{dx}\big|_{x=x_0}$.




In one dimensional case when we have a scalar function of one variable $y = f(x)$ we can plot this on the $x - y$ coordinate and the slope of the graph of the function at point x_0 is the derivative of f at x_0 and it is denoted by $f'(x_0)$ or $\frac{dy}{dx}|_{x=x_0}$.




In one dimensional case when we have a scalar function of one variable $y = f(x)$ we can plot this on the $x - y$ coordinate and the slope of the graph of the function at point x_0 is the derivative of f at x_0 and it is denoted by $f'(x_0)$ or $\frac{dy}{dx}\big|_{x=x_0}$.




In one dimensional case when we have a scalar function of one variable $y = f(x)$ we can plot this on the $x - y$ coordinate and the slope of the graph of the function at point x_0 is the derivative of f at x_0 and it is denoted by $f'(x_0)$ or $\frac{dy}{dx}\big|_{x=x_0}$.



In one dimensional case when we have a scalar function of one variable $y = f(x)$ we can plot this on the $x - y$ coordinate and the slope of the graph of the function at point x_0 is the derivative of f at x_0 and it is denoted by $f'(x_0)$ or $\frac{dy}{dx}\big|_{x=x_0}$.



In one dimensional case when we have a scalar function of one variable $y = f(x)$ we can plot this on the $x - y$ coordinate and the slope of the graph of the function at point x_0 is the derivative of f at x_0 and it is denoted by $f'(x_0)$ or $\frac{dy}{dx}|_{x=x_0}$.



Chain Rule

If $y = f(z)$ and $z = g(x)$ then $y = h(x)$. The derivative of y with respect to x is,

Chain Rule

If $y = f(z)$ and $z = g(x)$ then $y = h(x)$. The derivative of y with respect to x is,

$$\frac{dy}{dx} =$$

Chain Rule

If $y = f(z)$ and $z = g(x)$ then $y = h(x)$. The derivative of y with respect to x is,

$$\frac{dy}{dx} = \frac{dy}{dz}$$

Chain Rule

If $y = f(z)$ and $z = g(x)$ then $y = h(x)$. The derivative of y with respect to x is,

$$\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx}$$

Chain Rule

If $y = f(z)$ and $z = g(x)$ then $y = h(x)$. The derivative of y with respect to x is,

$$\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx}$$

or equivalently,

$$h'(x) =$$

Chain Rule

If $y = f(z)$ and $z = g(x)$ then $y = h(x)$. The derivative of y with respect to x is,

$$\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx}$$

or equivalently,

$$h'(x) = f'(z)$$

Chain Rule

If $y = f(z)$ and $z = g(x)$ then $y = h(x)$. The derivative of y with respect to x is,

$$\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx}$$

or equivalently,

$$h'(x) = f'(z)g'(x)$$

Chain Rule

If $y = f(z)$ and $z = g(x)$ then $y = h(x)$. The derivative of y with respect to x is,

$$\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx}$$

or equivalently,

$$\begin{aligned} h'(x) &= f'(z)g'(x) \\ &= f'(g(x)) \end{aligned}$$

Chain Rule

If $y = f(z)$ and $z = g(x)$ then $y = h(x)$. The derivative of y with respect to x is,

$$\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx}$$

or equivalently,

$$\begin{aligned} h'(x) &= f'(z)g'(x) \\ &= f'(g(x))g'(x) \end{aligned}$$

Important Functions

$$f(x) = x^n$$

Important Functions

$$f(x) = x^n \Rightarrow f'(x) = nx^{n-1}$$

Important Functions

$$f(x) = x^n \Rightarrow f'(x) = nx^{n-1}$$

$$f(x) = \sin(x)$$

Important Functions

$$f(x) = x^n \Rightarrow f'(x) = nx^{n-1}$$

$$f(x) = \sin(x) \Rightarrow f'(x) = \cos(x)$$

Important Functions

$$f(x) = x^n \Rightarrow f'(x) = nx^{n-1}$$

$$f(x) = \sin(x) \Rightarrow f'(x) = \cos(x)$$

$$f(x) = \cos(x)$$

Important Functions

$$f(x) = x^n \Rightarrow f'(x) = nx^{n-1}$$

$$f(x) = \sin(x) \Rightarrow f'(x) = \cos(x)$$

$$f(x) = \cos(x) \Rightarrow f'(x) = -\sin(x)$$

Important Functions

$$f(x) = x^n \Rightarrow f'(x) = nx^{n-1}$$

$$f(x) = \sin(x) \Rightarrow f'(x) = \cos(x)$$

$$f(x) = \cos(x) \Rightarrow f'(x) = -\sin(x)$$

$$f(x) = e^x$$

Important Functions

$$f(x) = x^n \Rightarrow f'(x) = nx^{n-1}$$

$$f(x) = \sin(x) \Rightarrow f'(x) = \cos(x)$$

$$f(x) = \cos(x) \Rightarrow f'(x) = -\sin(x)$$

$$f(x) = e^x \Rightarrow f'(x) = e^x$$

Important Functions

$$f(x) = x^n \Rightarrow f'(x) = nx^{n-1}$$

$$f(x) = \sin(x) \Rightarrow f'(x) = \cos(x)$$

$$f(x) = \cos(x) \Rightarrow f'(x) = -\sin(x)$$

$$f(x) = e^x \Rightarrow f'(x) = e^x$$

$$f(x) = \log(x)$$

Important Functions

$$f(x) = x^n \Rightarrow f'(x) = nx^{n-1}$$

$$f(x) = \sin(x) \Rightarrow f'(x) = \cos(x)$$

$$f(x) = \cos(x) \Rightarrow f'(x) = -\sin(x)$$

$$f(x) = e^x \Rightarrow f'(x) = e^x$$

$$f(x) = \log(x) \Rightarrow f'(x) = \frac{1}{x}$$

Derivative Rules

$$(u(x) + v(x))'$$

Derivative Rules

$$(u(x) + v(x))' = u'(x) + v'(x)$$

Derivative Rules

$$(u(x) + v(x))' = u'(x) + v'(x)$$

$$(u(x)v(x))'$$

Derivative Rules

$$(u(x) + v(x))' = u'(x) + v'(x)$$

$$(u(x)v(x))' = u'(x)v(x) + u(x)v'(x)$$

Derivative Rules

$$(u(x) + v(x))' = u'(x) + v'(x)$$

$$(u(x)v(x))' = u'(x)v(x) + u(x)v'(x)$$

$$(cu(x))'$$

Derivative Rules

$$(u(x) + v(x))' = u'(x) + v'(x)$$

$$(u(x)v(x))' = u'(x)v(x) + u(x)v'(x)$$

$$(cu(x))' = cu'(x)$$

Chain Rule of multi-variables

If $y = f(z_1, \dots, z_n)$ and $z_i = g_i(x)$ then $y = h(x)$ and we have,

Chain Rule of multi-variables

If $y = f(z_1, \dots, z_n)$ and $z_i = g_i(x)$ then $y = h(x)$ and we have,

$$\frac{dy}{dx} =$$

Chain Rule of multi-variables

If $y = f(z_1, \dots, z_n)$ and $z_i = g_i(x)$ then $y = h(x)$ and we have,

$$\frac{dy}{dx} = \frac{\partial y}{\partial z_1} \frac{dz_1}{dx}$$

Chain Rule of multi-variables

If $y = f(z_1, \dots, z_n)$ and $z_i = g_i(x)$ then $y = h(x)$ and we have,

$$\frac{dy}{dx} = \frac{\partial y}{\partial z_1} \frac{dz_1}{dx} + \dots +$$

Chain Rule of multi-variables

If $y = f(z_1, \dots, z_n)$ and $z_i = g_i(x)$ then $y = h(x)$ and we have,

$$\frac{dy}{dx} = \frac{\partial y}{\partial z_1} \frac{dz_1}{dx} + \dots + \frac{\partial y}{\partial z_n} \frac{dz_n}{dx}$$

Chain Rule of multi-variables

If $y = f(z_1, \dots, z_n)$ and $z_i = g_i(x)$ then $y = h(x)$ and we have,

$$\begin{aligned}\frac{dy}{dx} &= \frac{\partial y}{\partial z_1} \frac{dz_1}{dx} + \dots + \frac{\partial y}{\partial z_n} \frac{dz_n}{dx} \\ &= \frac{\partial y}{\partial z_1} g'_1(x)\end{aligned}$$

Chain Rule of multi-variables

If $y = f(z_1, \dots, z_n)$ and $z_i = g_i(x)$ then $y = h(x)$ and we have,

$$\begin{aligned}\frac{dy}{dx} &= \frac{\partial y}{\partial z_1} \frac{dz_1}{dx} + \dots + \frac{\partial y}{\partial z_n} \frac{dz_n}{dx} \\ &= \frac{\partial y}{\partial z_1} g'_1(x) + \dots +\end{aligned}$$

Chain Rule of multi-variables

If $y = f(z_1, \dots, z_n)$ and $z_i = g_i(x)$ then $y = h(x)$ and we have,

$$\begin{aligned}\frac{dy}{dx} &= \frac{\partial y}{\partial z_1} \frac{dz_1}{dx} + \dots + \frac{\partial y}{\partial z_n} \frac{dz_n}{dx} \\ &= \frac{\partial y}{\partial z_1} g'_1(x) + \dots + \frac{\partial y}{\partial z_n} g'_n(x)\end{aligned}$$

Example

Let $y = z_1^2 + z_2^2$ and $z_1 = \log(x)$, $z_2 = \sin(x)$ then $\frac{dy}{dx}\big|_{x=1}$ is

Example

Let $y = z_1^2 + z_2^2$ and $z_1 = \log(x)$, $z_2 = \sin(x)$ then $\frac{dy}{dx}|_{x=1}$ is

$$\frac{dy}{dx} =$$

Example

Let $y = z_1^2 + z_2^2$ and $z_1 = \log(x)$, $z_2 = \sin(x)$ then $\frac{dy}{dx}|_{x=1}$ is

$$\frac{dy}{dx} = \frac{\partial y}{\partial z_1} \frac{dz_1}{dx}$$

Example

Let $y = z_1^2 + z_2^2$ and $z_1 = \log(x)$, $z_2 = \sin(x)$ then $\frac{dy}{dx}|_{x=1}$ is

$$\begin{aligned}\frac{dy}{dx} &= \frac{\partial y}{\partial z_1} \frac{dz_1}{dx} + \frac{\partial y}{\partial z_2} \frac{dz_2}{dx} \\ &= \end{aligned}$$

Example

Let $y = z_1^2 + z_2^2$ and $z_1 = \log(x)$, $z_2 = \sin(x)$ then $\frac{dy}{dx}|_{x=1}$ is

$$\begin{aligned}\frac{dy}{dx} &= \frac{\partial y}{\partial z_1} \frac{dz_1}{dx} + \frac{\partial y}{\partial z_2} \frac{dz_2}{dx} \\ &= (2z_1)\left(\frac{1}{x}\right)\end{aligned}$$

Example

Let $y = z_1^2 + z_2^2$ and $z_1 = \log(x)$, $z_2 = \sin(x)$ then $\frac{dy}{dx}\big|_{x=1}$ is

$$\begin{aligned}\frac{dy}{dx} &= \frac{\partial y}{\partial z_1} \frac{dz_1}{dx} + \frac{\partial y}{\partial z_2} \frac{dz_2}{dx} \\ &= (2z_1)\left(\frac{1}{x}\right) + (2z_2)(\cos(x)) \\ &= \end{aligned}$$

Example

Let $y = z_1^2 + z_2^2$ and $z_1 = \log(x)$, $z_2 = \sin(x)$ then $\frac{dy}{dx}\big|_{x=1}$ is

$$\begin{aligned}\frac{dy}{dx} &= \frac{\partial y}{\partial z_1} \frac{dz_1}{dx} + \frac{\partial y}{\partial z_2} \frac{dz_2}{dx} \\ &= (2z_1)\left(\frac{1}{x}\right) + (2z_2)(\cos(x)) \\ &= (2 \times 0)\left(\frac{1}{1}\right)\end{aligned}$$

Example

Let $y = z_1^2 + z_2^2$ and $z_1 = \log(x)$, $z_2 = \sin(x)$ then $\frac{dy}{dx}|_{x=1}$ is

$$\begin{aligned}\frac{dy}{dx} &= \frac{\partial y}{\partial z_1} \frac{dz_1}{dx} + \frac{\partial y}{\partial z_2} \frac{dz_2}{dx} \\ &= (2z_1)\left(\frac{1}{x}\right) + (2z_2)(\cos(x)) \\ &= (2 \times 0)\left(\frac{1}{1}\right) + (2 \times \sin 1)(\cos 1) \\ &= \end{aligned}$$

Example

Let $y = z_1^2 + z_2^2$ and $z_1 = \log(x)$, $z_2 = \sin(x)$ then $\frac{dy}{dx}|_{x=1}$ is

$$\begin{aligned}\frac{dy}{dx} &= \frac{\partial y}{\partial z_1} \frac{dz_1}{dx} + \frac{\partial y}{\partial z_2} \frac{dz_2}{dx} \\ &= (2z_1)\left(\frac{1}{x}\right) + (2z_2)(\cos(x)) \\ &= (2 \times 0)\left(\frac{1}{1}\right) + (2 \times \sin 1)(\cos 1) \\ &= \sin(2) \\ &\approx\end{aligned}$$

Example

Let $y = z_1^2 + z_2^2$ and $z_1 = \log(x)$, $z_2 = \sin(x)$ then $\frac{dy}{dx}|_{x=1}$ is

$$\begin{aligned}\frac{dy}{dx} &= \frac{\partial y}{\partial z_1} \frac{dz_1}{dx} + \frac{\partial y}{\partial z_2} \frac{dz_2}{dx} \\ &= (2z_1)\left(\frac{1}{x}\right) + (2z_2)(\cos(x)) \\ &= (2 \times 0)\left(\frac{1}{1}\right) + (2 \times \sin 1)(\cos 1) \\ &= \sin(2) \\ &\approx 0.034899497\end{aligned}$$

If $f : \mathbb{R}^n \rightarrow \mathbb{R}$, i.e., $y = f(x_1, \dots, x_n)$, i.e., scalar y depends on reals x_1, \dots, x_n , then the **Gradient** of f at \mathbf{x}_0 is

If $f : \mathbb{R}^n \rightarrow \mathbb{R}$, i.e., $y = f(x_1, \dots, x_n)$, i.e., scalar y depends on reals x_1, \dots, x_n , then the **Gradient** of f at \mathbf{x}_0 is

$$\nabla f(\mathbf{x}_0) = \begin{bmatrix} \left. \frac{\partial y}{\partial x_1} \right|_{\mathbf{x}_0} \\ \vdots \\ \left. \frac{\partial y}{\partial x_n} \right|_{\mathbf{x}_0} \end{bmatrix}$$

$y = x_1 + x_2$, then if $y = \text{sum}(x_1, x_2)$ then the Gradient of *sum* at $\mathbf{x}_0 = (2, 3)$ is,

$y = x_1 + x_2$, then if $y = \text{sum}(x_1, x_2)$ then the Gradient of *sum* at $\mathbf{x_0} = (2, 3)$ is,

$$\nabla \text{sum}(2, 3) =$$

$y = x_1 + x_2$, then if $y = \text{sum}(x_1, x_2)$ then the Gradient of *sum* at $\mathbf{x_0} = (2, 3)$ is,

$$\nabla \text{sum}(2, 3) = \begin{bmatrix} \frac{\partial \text{sum}}{\partial x_1} \\ \frac{\partial \text{sum}}{\partial x_2} \end{bmatrix}$$

$y = x_1 + x_2$, then if $y = \text{sum}(x_1, x_2)$ then the Gradient of *sum* at $\mathbf{x_0} = (2, 3)$ is,

$$\begin{aligned}\nabla \text{sum}(2, 3) &= \begin{bmatrix} \frac{\partial \text{sum}}{\partial x_1} \\ \frac{\partial \text{sum}}{\partial x_2} \end{bmatrix} \\ &= \begin{bmatrix} 1 \\ 1 \end{bmatrix}\end{aligned}$$

Directional Derivative

Let $\mathbf{u} \in \mathbb{R}^n$ and $\|\mathbf{u}\| = 1$, then the **Directional Derivative** of f in the direction of \mathbf{u} in \mathbf{x}_0 is denoted by $\mathcal{D}_{\mathbf{u}}f(\mathbf{x}_0)$ and is define by,

Directional Derivative

Let $\mathbf{u} \in \mathbb{R}^n$ and $\|\mathbf{u}\| = 1$, then the **Directional Derivative** of f in the direction of \mathbf{u} in \mathbf{x}_0 is denoted by $\mathcal{D}_{\mathbf{u}}f(\mathbf{x}_0)$ and is defined by,

$$\mathcal{D}_{\mathbf{u}}f(\mathbf{x}_0) = \nabla f(\mathbf{x}_0) \cdot \mathbf{u}$$

In which direction, we have maximum Directional Derivative?

In which direction, we have maximum Directional Derivative? If we choose $\mathbf{u} = \frac{\nabla f(\mathbf{x}_0)}{\|\nabla f(\mathbf{x}_0)\|}$ then we have maximum directional derivative and

In which direction, we have maximum Directional Derivative? If we choose $\mathbf{u} = \frac{\nabla f(\mathbf{x}_0)}{\|\nabla f(\mathbf{x}_0)\|}$ then we have maximum directional derivative and in the direction of $-\mathbf{u}$ we have minimum directional derivative.

What is the directional derivative of the function $f(x_1, x_2) = x_1^2 + x_2^2$ in the direction of $\mathbf{u} = \frac{1}{25}(3, 4)$ in $\mathbf{x}_0 = (1, 1)$?

What is the directional derivative of the function $f(x_1, x_2) = x_1^2 + x_2^2$ in the direction of $\mathbf{u} = \frac{1}{25}(3, 4)$ in $\mathbf{x}_0 = (1, 1)$?

Since $\nabla f(x_1, x_2) = \begin{bmatrix} 2x_1 \\ 2x_2 \end{bmatrix}$ then,

What is the directional derivative of the function $f(x_1, x_2) = x_1^2 + x_2^2$ in the direction of $\mathbf{u} = \frac{1}{25}(3, 4)$ in $\mathbf{x}_0 = (1, 1)$?

Since $\nabla f(x_1, x_2) = \begin{bmatrix} 2x_1 \\ 2x_2 \end{bmatrix}$ then,

$$\mathcal{D}_{\mathbf{u}}f(x_1, x_2) = \nabla f(x_1, x_2) \cdot \mathbf{u}$$

What is the directional derivative of the function $f(x_1, x_2) = x_1^2 + x_2^2$ in the direction of $\mathbf{u} = \frac{1}{25}(3, 4)$ in $\mathbf{x}_0 = (1, 1)$?

Since $\nabla f(x_1, x_2) = \begin{bmatrix} 2x_1 \\ 2x_2 \end{bmatrix}$ then,

$$\begin{aligned}\mathcal{D}_{\mathbf{u}}f(x_1, x_2) &= \nabla f(x_1, x_2) \cdot \mathbf{u} \\ &= \begin{bmatrix} 2 \\ 2 \end{bmatrix} \cdot \frac{1}{25} \begin{bmatrix} 3 \\ 4 \end{bmatrix}\end{aligned}$$

What is the directional derivative of the function $f(x_1, x_2) = x_1^2 + x_2^2$ in the direction of $\mathbf{u} = \frac{1}{25}(3, 4)$ in $\mathbf{x}_0 = (1, 1)$?

Since $\nabla f(x_1, x_2) = \begin{bmatrix} 2x_1 \\ 2x_2 \end{bmatrix}$ then,

$$\begin{aligned}\mathcal{D}_{\mathbf{u}}f(x_1, x_2) &= \nabla f(x_1, x_2) \cdot \mathbf{u} \\ &= \begin{bmatrix} 2 \\ 2 \end{bmatrix} \cdot \frac{1}{25} \begin{bmatrix} 3 \\ 4 \end{bmatrix} \\ &= \frac{14}{25}\end{aligned}$$

What is the directional derivative of the function $f(x_1, x_2) = x_1^2 + x_2^2$ in the direction of $\mathbf{u} = \frac{1}{25}(3, 4)$ in $\mathbf{x}_0 = (1, 1)$?

Since $\nabla f(x_1, x_2) = \begin{bmatrix} 2x_1 \\ 2x_2 \end{bmatrix}$ then,

$$\begin{aligned}\mathcal{D}_{\mathbf{u}}f(x_1, x_2) &= \nabla f(x_1, x_2) \cdot \mathbf{u} \\ &= \begin{bmatrix} 2 \\ 2 \end{bmatrix} \cdot \frac{1}{25} \begin{bmatrix} 3 \\ 4 \end{bmatrix} \\ &= \frac{14}{25} \\ &= 0.56\end{aligned}$$

What is the directional derivative of the function $f(x_1, x_2) = x_1^2 + x_2^2$ in the direction of $\mathbf{u} = \frac{1}{25}(3, 4)$ in $\mathbf{x}_0 = (1, 1)$?

Since $\nabla f(x_1, x_2) = \begin{bmatrix} 2x_1 \\ 2x_2 \end{bmatrix}$ then,

$$\begin{aligned}\mathcal{D}_{\mathbf{u}}f(x_1, x_2) &= \nabla f(x_1, x_2) \cdot \mathbf{u} \\ &= \begin{bmatrix} 2 \\ 2 \end{bmatrix} \cdot \frac{1}{25} \begin{bmatrix} 3 \\ 4 \end{bmatrix} \\ &= \frac{14}{25} \\ &= 0.56\end{aligned}$$

while $\mathcal{D}_{\frac{\nabla f(\mathbf{x}_0)}{\|\nabla f(\mathbf{x}_0)\|}} f(x_1, x_2) = \|\nabla f(\mathbf{x}_0)\| \approx 2.82$

Jacobian

For a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, such that $(y_1, \dots, y_m) = f(x_1, \dots, x_n)$, the **Jacobian** of f at \mathbf{x}_0 is a $m \times n$ matrix and denoted by $\mathbf{J}_f(\mathbf{x}_0)$ and is defined as follow:

Jacobian

For a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, such that $(y_1, \dots, y_m) = f(x_1, \dots, x_n)$, the **Jacobian** of f at \mathbf{x}_0 is a $m \times n$ matrix and denoted by $\mathbf{J}_f(\mathbf{x}_0)$ and is defined as follow:

$$\mathbf{J}_f(\mathbf{x}_0) =$$

Jacobian

For a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, such that $(y_1, \dots, y_m) = f(x_1, \dots, x_n)$, the **Jacobian** of f at \mathbf{x}_0 is a $m \times n$ matrix and denoted by $\mathbf{J}_f(\mathbf{x}_0)$ and is defined as follow:

$$\mathbf{J}_f(\mathbf{x}_0) = \begin{bmatrix} \left. \frac{\partial y_1}{\partial x_1} \right|_{\mathbf{x}_0} & \cdots & \left. \frac{\partial y_1}{\partial x_n} \right|_{\mathbf{x}_0} \\ \vdots & \ddots & \vdots \\ \left. \frac{\partial y_m}{\partial x_1} \right|_{\mathbf{x}_0} & \cdots & \left. \frac{\partial y_m}{\partial x_n} \right|_{\mathbf{x}_0} \end{bmatrix}$$

Jacobian

For a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, such that $(y_1, \dots, y_m) = f(x_1, \dots, x_n)$, the **Jacobian** of f at \mathbf{x}_0 is a $m \times n$ matrix and denoted by $\mathbf{J}_f(\mathbf{x}_0)$ and is defined as follow:

$$\mathbf{J}_f(\mathbf{x}_0) = \begin{bmatrix} \left. \frac{\partial y_1}{\partial x_1} \right|_{\mathbf{x}_0} & \cdots & \left. \frac{\partial y_1}{\partial x_n} \right|_{\mathbf{x}_0} \\ \vdots & \ddots & \vdots \end{bmatrix}$$

Jacobian

For a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, such that $(y_1, \dots, y_m) = f(x_1, \dots, x_n)$, the **Jacobian** of f at \mathbf{x}_0 is a $m \times n$ matrix and denoted by $\mathbf{J}_f(\mathbf{x}_0)$ and is defined as follow:

$$\mathbf{J}_f(\mathbf{x}_0) = \begin{bmatrix} \left. \frac{\partial y_1}{\partial x_1} \right|_{\mathbf{x}_0} & \cdots & \left. \frac{\partial y_1}{\partial x_n} \right|_{\mathbf{x}_0} \\ \vdots & \ddots & \vdots \\ \left. \frac{\partial y_m}{\partial x_1} \right|_{\mathbf{x}_0} & \cdots & \left. \frac{\partial y_m}{\partial x_n} \right|_{\mathbf{x}_0} \end{bmatrix}$$

Jacobian

For a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, such that $(y_1, \dots, y_m) = f(x_1, \dots, x_n)$, the **Jacobian** of f at \mathbf{x}_0 is a $m \times n$ matrix and denoted by $\mathbf{J}_f(\mathbf{x}_0)$ and is defined as follow:

$$\mathbf{J}_f(\mathbf{x}_0) = \begin{bmatrix} \left. \frac{\partial y_1}{\partial x_1} \right|_{\mathbf{x}_0} & \cdots & \left. \frac{\partial y_1}{\partial x_n} \right|_{\mathbf{x}_0} \\ \vdots & \ddots & \vdots \\ \left. \frac{\partial y_m}{\partial x_1} \right|_{\mathbf{x}_0} & \cdots & \left. \frac{\partial y_m}{\partial x_n} \right|_{\mathbf{x}_0} \end{bmatrix}$$

another notation,

Jacobian

For a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, such that $(y_1, \dots, y_m) = f(x_1, \dots, x_n)$, the **Jacobian** of f at \mathbf{x}_0 is a $m \times n$ matrix and denoted by $\mathbf{J}_f(\mathbf{x}_0)$ and is defined as follow:

$$\mathbf{J}_f(\mathbf{x}_0) = \begin{bmatrix} \left. \frac{\partial y_1}{\partial x_1} \right|_{\mathbf{x}_0} & \cdots & \left. \frac{\partial y_1}{\partial x_n} \right|_{\mathbf{x}_0} \\ \vdots & \ddots & \vdots \\ \left. \frac{\partial y_m}{\partial x_1} \right|_{\mathbf{x}_0} & \cdots & \left. \frac{\partial y_m}{\partial x_n} \right|_{\mathbf{x}_0} \end{bmatrix}$$

another notation,

$$\mathbf{J}_f(\mathbf{x}_0) =$$

Jacobian

For a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, such that $(y_1, \dots, y_m) = f(x_1, \dots, x_n)$, the **Jacobian** of f at \mathbf{x}_0 is a $m \times n$ matrix and denoted by $\mathbf{J}_f(\mathbf{x}_0)$ and is defined as follow:

$$\mathbf{J}_f(\mathbf{x}_0) = \begin{bmatrix} \left. \frac{\partial y_1}{\partial x_1} \right|_{\mathbf{x}_0} & \cdots & \left. \frac{\partial y_1}{\partial x_n} \right|_{\mathbf{x}_0} \\ \vdots & \ddots & \vdots \\ \left. \frac{\partial y_m}{\partial x_1} \right|_{\mathbf{x}_0} & \cdots & \left. \frac{\partial y_m}{\partial x_n} \right|_{\mathbf{x}_0} \end{bmatrix}$$

another notation,

$$\mathbf{J}_f(\mathbf{x}_0) = \left. \frac{\partial(y_1, \dots, y_m)}{\partial(x_1, \dots, x_n)} \right|_{\mathbf{x}_0}$$

When $m = 1$ then the Jacobian of $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is a row vector and its transpos is the gradient of f , i.e.,

When $m = 1$ then the Jacobian of $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is a row vector and its transpos is the gradient of f , i.e.,

$$\mathbf{J}_f(\mathbf{x}) = (\nabla f(\mathbf{x}))^T$$

Let $f(x_1, x_2) = (x_1x_2, \sin(x_1), \cos(x_2))$. What is $\mathbf{J}_f(2, 3)$?

Let $f(x_1, x_2) = (x_1x_2, \sin(x_1), \cos(x_2))$. What is $\mathbf{J}_f(2, 3)$?

$$\mathbf{J}_f(2, 3) =$$

Let $f(x_1, x_2) = (x_1x_2, \sin(x_1), \cos(x_2))$. What is $\mathbf{J}_f(2, 3)$?

$$\mathbf{J}_f(2, 3) = \begin{bmatrix} \left. \frac{\partial(x_1x_2)}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(x_1x_2)}{\partial x_2} \right|_{(2,3)} \end{bmatrix}$$

Let $f(x_1, x_2) = (x_1x_2, \sin(x_1), \cos(x_2))$. What is $\mathbf{J}_f(2, 3)$?

$$\mathbf{J}_f(2, 3) = \begin{bmatrix} \left. \frac{\partial(x_1x_2)}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(x_1x_2)}{\partial x_2} \right|_{(2,3)} \\ \left. \frac{\partial(\sin(x_1))}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(\sin(x_1))}{\partial x_2} \right|_{(2,3)} \end{bmatrix}$$

Let $f(x_1, x_2) = (x_1x_2, \sin(x_1), \cos(x_2))$. What is $\mathbf{J}_f(2, 3)$?

$$\mathbf{J}_f(2, 3) = \begin{bmatrix} \left. \frac{\partial(x_1x_2)}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(x_1x_2)}{\partial x_2} \right|_{(2,3)} \\ \left. \frac{\partial(\sin(x_1))}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(\sin(x_1))}{\partial x_2} \right|_{(2,3)} \\ \left. \frac{\partial(\cos(x_2))}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(\cos(x_2))}{\partial x_2} \right|_{(2,3)} \end{bmatrix}$$

Let $f(x_1, x_2) = (x_1x_2, \sin(x_1), \cos(x_2))$. What is $\mathbf{J}_f(2, 3)$?

$$\begin{aligned}\mathbf{J}_f(2, 3) &= \begin{bmatrix} \left. \frac{\partial(x_1x_2)}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(x_1x_2)}{\partial x_2} \right|_{(2,3)} \\ \left. \frac{\partial(\sin(x_1))}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(\sin(x_1))}{\partial x_2} \right|_{(2,3)} \\ \left. \frac{\partial(\cos(x_2))}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(\cos(x_2))}{\partial x_2} \right|_{(2,3)} \end{bmatrix} \\ &= \begin{bmatrix} x_2|_{(2,3)} & x_1|_{(2,3)} \\ 0 & 0 \\ 0 & 0 \end{bmatrix}\end{aligned}$$

Let $f(x_1, x_2) = (x_1x_2, \sin(x_1), \cos(x_2))$. What is $\mathbf{J}_f(2, 3)$?

$$\begin{aligned} \mathbf{J}_f(2, 3) &= \begin{bmatrix} \left. \frac{\partial(x_1x_2)}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(x_1x_2)}{\partial x_2} \right|_{(2,3)} \\ \left. \frac{\partial(\sin(x_1))}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(\sin(x_1))}{\partial x_2} \right|_{(2,3)} \\ \left. \frac{\partial(\cos(x_2))}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(\cos(x_2))}{\partial x_2} \right|_{(2,3)} \end{bmatrix} \\ &= \begin{bmatrix} x_2|_{(2,3)} & x_1|_{(2,3)} \\ \cos(x_1)|_{(2,3)} & 0|_{(2,3)} \end{bmatrix} \end{aligned}$$

Let $f(x_1, x_2) = (x_1x_2, \sin(x_1), \cos(x_2))$. What is $\mathbf{J}_f(2, 3)$?

$$\begin{aligned}\mathbf{J}_f(2, 3) &= \begin{bmatrix} \left. \frac{\partial(x_1x_2)}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(x_1x_2)}{\partial x_2} \right|_{(2,3)} \\ \left. \frac{\partial(\sin(x_1))}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(\sin(x_1))}{\partial x_2} \right|_{(2,3)} \\ \left. \frac{\partial(\cos(x_2))}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(\cos(x_2))}{\partial x_2} \right|_{(2,3)} \end{bmatrix} \\ &= \begin{bmatrix} x_2|_{(2,3)} & x_1|_{(2,3)} \\ \cos(x_1)|_{(2,3)} & 0|_{(2,3)} \\ 0|_{(2,3)} & -\sin(x_2)|_{(2,3)} \end{bmatrix}\end{aligned}$$

Let $f(x_1, x_2) = (x_1x_2, \sin(x_1), \cos(x_2))$. What is $J_f(2, 3)$?

$$\begin{aligned}
 J_f(2, 3) &= \begin{bmatrix} \left. \frac{\partial(x_1x_2)}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(x_1x_2)}{\partial x_2} \right|_{(2,3)} \\ \left. \frac{\partial(\sin(x_1))}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(\sin(x_1))}{\partial x_2} \right|_{(2,3)} \\ \left. \frac{\partial(\cos(x_2))}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(\cos(x_2))}{\partial x_2} \right|_{(2,3)} \end{bmatrix} \\
 &= \begin{bmatrix} x_2|_{(2,3)} & x_1|_{(2,3)} \\ \cos(x_1)|_{(2,3)} & 0|_{(2,3)} \\ 0|_{(2,3)} & -\sin(x_2)|_{(2,3)} \end{bmatrix} \\
 &= \begin{bmatrix} 3 & 2 \\ \cos(2) & 0 \\ 0 & -\sin(3) \end{bmatrix}
 \end{aligned}$$

Let $f(x_1, x_2) = (x_1x_2, \sin(x_1), \cos(x_2))$. What is $\mathbf{J}_f(2, 3)$?

$$\begin{aligned}
 \mathbf{J}_f(2, 3) &= \begin{bmatrix} \left. \frac{\partial(x_1x_2)}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(x_1x_2)}{\partial x_2} \right|_{(2,3)} \\ \left. \frac{\partial(\sin(x_1))}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(\sin(x_1))}{\partial x_2} \right|_{(2,3)} \\ \left. \frac{\partial(\cos(x_2))}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(\cos(x_2))}{\partial x_2} \right|_{(2,3)} \end{bmatrix} \\
 &= \begin{bmatrix} x_2|_{(2,3)} & x_1|_{(2,3)} \\ \cos(x_1)|_{(2,3)} & 0|_{(2,3)} \\ 0|_{(2,3)} & -\sin(x_2)|_{(2,3)} \end{bmatrix} \\
 &= \begin{bmatrix} 3 & 2 \\ 0.99 & 0 \end{bmatrix}
 \end{aligned}$$

Let $f(x_1, x_2) = (x_1x_2, \sin(x_1), \cos(x_2))$. What is $J_f(2, 3)$?

$$\begin{aligned}
 J_f(2, 3) &= \begin{bmatrix} \left. \frac{\partial(x_1x_2)}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(x_1x_2)}{\partial x_2} \right|_{(2,3)} \\ \left. \frac{\partial(\sin(x_1))}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(\sin(x_1))}{\partial x_2} \right|_{(2,3)} \\ \left. \frac{\partial(\cos(x_2))}{\partial x_1} \right|_{(2,3)} & \left. \frac{\partial(\cos(x_2))}{\partial x_2} \right|_{(2,3)} \end{bmatrix} \\
 &= \begin{bmatrix} x_2|_{(2,3)} & x_1|_{(2,3)} \\ \cos(x_1)|_{(2,3)} & 0|_{(2,3)} \\ 0|_{(2,3)} & -\sin(x_2)|_{(2,3)} \end{bmatrix} \\
 &= \begin{bmatrix} 3 & 2 \\ 0.99 & 0 \\ 0 & -0.05 \end{bmatrix}
 \end{aligned}$$

Let define $f(\mathbf{x}) = \mathbf{x}^T \mathbf{a}$, where $\mathbf{a} : \mathbb{R}^n$ is a constant vector. Then $f : \mathbb{R}^n \rightarrow \mathbb{R}$.
What is the Jacobian of f at \mathbf{x}_0 ?

Let define $f(\mathbf{x}) = \mathbf{x}^T \mathbf{a}$, where $\mathbf{a} : \mathbb{R}^n$ is a constant vector. Then $f : \mathbb{R}^n \rightarrow \mathbb{R}$.
What is the Jacobian of f at \mathbf{x}_0 ?

$$f(\mathbf{x}_0) =$$

Let define $f(\mathbf{x}) = \mathbf{x}^T \mathbf{a}$, where $\mathbf{a} : \mathbb{R}^n$ is a constant vector. Then $f : \mathbb{R}^n \rightarrow \mathbb{R}$.
What is the Jacobian of f at \mathbf{x}_0 ?

$$f(\mathbf{x}_0) = [x_1 \quad \dots \quad x_n]$$

Let define $f(\mathbf{x}) = \mathbf{x}^T \mathbf{a}$, where $\mathbf{a} : \mathbb{R}^n$ is a constant vector. Then $f : \mathbb{R}^n \rightarrow \mathbb{R}$.
What is the Jacobian of f at \mathbf{x}_0 ?

$$f(\mathbf{x}_0) = \begin{bmatrix} x_1 & \dots & x_n \end{bmatrix} \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix}$$

Let define $f(\mathbf{x}) = \mathbf{x}^T \mathbf{a}$, where $\mathbf{a} : \mathbb{R}^n$ is a constant vector. Then $f : \mathbb{R}^n \rightarrow \mathbb{R}$.
What is the Jacobian of f at \mathbf{x}_0 ?

$$\begin{aligned} f(\mathbf{x}_0) &= \begin{bmatrix} x_1 & \dots & x_n \end{bmatrix} \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} \\ &= x_1 a_1 + \dots + x_n a_n \end{aligned}$$

$$\mathbf{J}_f(\mathbf{x}_0) = \left[\frac{\partial(x_1 a_1 + \cdots + x_n a_n)}{\partial x_1} \quad \cdots \quad \frac{\partial(x_1 a_1 + \cdots + x_n a_n)}{\partial x_n} \right]$$

$$\begin{aligned}\mathbf{J}_f(\mathbf{x}_0) &= \left[\frac{\partial(x_1 a_1 + \dots + x_n a_n)}{\partial x_1} \quad \dots \quad \frac{\partial(x_1 a_1 + \dots + x_n a_n)}{\partial x_n} \right] \\ &= [a_1 \quad \dots \quad a_n]\end{aligned}$$

$$\begin{aligned}
 \mathbf{J}_f(\mathbf{x}_0) &= \left[\frac{\partial(x_1 a_1 + \dots + x_n a_n)}{\partial x_1} \quad \dots \quad \frac{\partial(x_1 a_1 + \dots + x_n a_n)}{\partial x_n} \right] \\
 &= [a_1 \quad \dots \quad a_n] \\
 &= \mathbf{a}^T
 \end{aligned}$$

Let define $f(\mathbf{x}) = \mathbf{a}^T \mathbf{x}$, where $\mathbf{a} : \mathbb{R}^n$ is a constant vector. Then $f : \mathbb{R}^n \rightarrow \mathbb{R}$.
What is the Jacobian of f at \mathbf{x}_0 ?

Let define $f(\mathbf{x}) = \mathbf{a}^T \mathbf{x}$, where $\mathbf{a} : \mathbb{R}^n$ is a constant vector. Then $f : \mathbb{R}^n \rightarrow \mathbb{R}$.
What is the Jacobian of f at \mathbf{x}_0 ?

$$f(\mathbf{x}_0) =$$

Let define $f(\mathbf{x}) = \mathbf{a}^T \mathbf{x}$, where $\mathbf{a} : \mathbb{R}^n$ is a constant vector. Then $f : \mathbb{R}^n \rightarrow \mathbb{R}$.
What is the Jacobian of f at \mathbf{x}_0 ?

$$f(\mathbf{x}_0) = [a_1 \quad \dots \quad a_n]$$

Let define $f(\mathbf{x}) = \mathbf{a}^T \mathbf{x}$, where $\mathbf{a} : \mathbb{R}^n$ is a constant vector. Then $f : \mathbb{R}^n \rightarrow \mathbb{R}$.
What is the Jacobian of f at \mathbf{x}_0 ?

$$f(\mathbf{x}_0) = \begin{bmatrix} a_1 & \dots & a_n \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$$

Let define $f(\mathbf{x}) = \mathbf{a}^T \mathbf{x}$, where $\mathbf{a} : \mathbb{R}^n$ is a constant vector. Then $f : \mathbb{R}^n \rightarrow \mathbb{R}$. What is the Jacobian of f at \mathbf{x}_0 ?

$$\begin{aligned} f(\mathbf{x}_0) &= [a_1 \quad \dots \quad a_n] \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} \\ &= x_1 a_1 + \dots + x_n a_n \end{aligned}$$

$$\mathbf{J}_f(\mathbf{x}_0) = \left[\frac{\partial(x_1a_1+\cdots+x_na_n)}{\partial x_1} \quad \cdots \quad \frac{\partial(x_1a_1+\cdots+x_na_n)}{\partial x_n} \right]$$

$$\begin{aligned}\mathbf{J}_f(\mathbf{x}_0) &= \left[\frac{\partial(x_1 a_1 + \dots + x_n a_n)}{\partial x_1} \quad \dots \quad \frac{\partial(x_1 a_1 + \dots + x_n a_n)}{\partial x_n} \right] \\ &= [a_1 \quad \dots \quad a_n]\end{aligned}$$

$$\begin{aligned}
 \mathbf{J}_f(\mathbf{x}_0) &= \left[\frac{\partial(x_1 a_1 + \dots + x_n a_n)}{\partial x_1} \quad \dots \quad \frac{\partial(x_1 a_1 + \dots + x_n a_n)}{\partial x_n} \right] \\
 &= [a_1 \quad \dots \quad a_n] \\
 &= \mathbf{a}^T
 \end{aligned}$$

Let $\mathbf{W}_{m \times n}$ be a constant matrix and let define $f(\mathbf{x}) = \mathbf{W}\mathbf{x}$. Then $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$.
What is the Jacobian of f at \mathbf{x}_0 ?

$$f(\mathbf{x}) =$$

Let $\mathbf{W}_{m \times n}$ be a constant matrix and let define $f(\mathbf{x}) = \mathbf{W}\mathbf{x}$. Then $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$. What is the Jacobian of f at \mathbf{x}_0 ?

$$f(\mathbf{x}) = \begin{bmatrix} w_{11} & \dots & w_{1n} \\ \vdots & & \vdots \\ w_{m1} & \dots & w_{mn} \end{bmatrix}$$

Let $\mathbf{W}_{m \times n}$ be a constant matrix and let define $f(\mathbf{x}) = \mathbf{W}\mathbf{x}$. Then $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$. What is the Jacobian of f at \mathbf{x}_0 ?

$$f(\mathbf{x}) = \begin{bmatrix} w_{11} & \dots & w_{1n} \\ \vdots & \ddots & \vdots \end{bmatrix}$$

Let $\mathbf{W}_{m \times n}$ be a constant matrix and let define $f(\mathbf{x}) = \mathbf{W}\mathbf{x}$. Then $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$. What is the Jacobian of f at \mathbf{x}_0 ?

$$f(\mathbf{x}) = \begin{bmatrix} w_{11} & \dots & w_{1n} \\ \vdots & \ddots & \vdots \\ w_{m1} & \dots & w_{mn} \end{bmatrix}$$

Let $\mathbf{W}_{m \times n}$ be a constant matrix and let define $f(\mathbf{x}) = \mathbf{W}\mathbf{x}$. Then $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$. What is the Jacobian of f at \mathbf{x}_0 ?

$$f(\mathbf{x}) = \begin{bmatrix} w_{11} & \dots & w_{1n} \\ \vdots & \ddots & \vdots \\ w_{m1} & \dots & w_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$$

Let $\mathbf{W}_{m \times n}$ be a constant matrix and let define $f(\mathbf{x}) = \mathbf{W}\mathbf{x}$. Then $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$. What is the Jacobian of f at \mathbf{x}_0 ?

$$f(\mathbf{x}) = \begin{bmatrix} w_{11} & \dots & w_{1n} \\ \vdots & \ddots & \vdots \\ w_{m1} & \dots & w_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \end{bmatrix}$$

Let $\mathbf{W}_{m \times n}$ be a constant matrix and let define $f(\mathbf{x}) = \mathbf{W}\mathbf{x}$. Then $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$. What is the Jacobian of f at \mathbf{x}_0 ?

$$f(\mathbf{x}) = \begin{bmatrix} w_{11} & \dots & w_{1n} \\ \vdots & \ddots & \vdots \\ w_{m1} & \dots & w_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$$

Let $\mathbf{W}_{m \times n}$ be a constant matrix and let define $f(\mathbf{x}) = \mathbf{W}\mathbf{x}$. Then $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$. What is the Jacobian of f at \mathbf{x}_0 ?

$$\begin{aligned} f(\mathbf{x}) &= \begin{bmatrix} w_{11} & \dots & w_{1n} \\ \vdots & \ddots & \vdots \\ w_{m1} & \dots & w_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} \\ &= \begin{bmatrix} w_{11}x_1 + \dots + w_{1n}x_n \\ \vdots \\ w_{m1}x_1 + \dots + w_{mn}x_n \end{bmatrix} \end{aligned}$$

Let $\mathbf{W}_{m \times n}$ be a constant matrix and let define $f(\mathbf{x}) = \mathbf{W}\mathbf{x}$. Then $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$. What is the Jacobian of f at \mathbf{x}_0 ?

$$\begin{aligned} f(\mathbf{x}) &= \begin{bmatrix} w_{11} & \dots & w_{1n} \\ \vdots & \ddots & \vdots \\ w_{m1} & \dots & w_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} \\ &= \begin{bmatrix} w_{11}x_1 + \dots + w_{1n}x_n \\ \vdots \end{bmatrix} \end{aligned}$$

Let $\mathbf{W}_{m \times n}$ be a constant matrix and let define $f(\mathbf{x}) = \mathbf{W}\mathbf{x}$. Then $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$. What is the Jacobian of f at \mathbf{x}_0 ?

$$\begin{aligned} f(\mathbf{x}) &= \begin{bmatrix} w_{11} & \dots & w_{1n} \\ \vdots & \ddots & \vdots \\ w_{m1} & \dots & w_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} \\ &= \begin{bmatrix} w_{11}x_1 + \dots + w_{1n}x_n \\ \vdots \\ w_{m1}x_1 + \dots + w_{mn}x_n \end{bmatrix} \end{aligned}$$

$$\mathbf{J}_f(\mathbf{x}_0) = \begin{bmatrix} \frac{\partial(w_{11}x_1 + \dots + w_{1n}x_n)}{\partial x_1} & \dots & \frac{\partial(w_{11}x_1 + \dots + w_{1n}x_n)}{\partial x_n} \end{bmatrix}$$

$$\mathbf{J}_f(\mathbf{x}_0) = \begin{bmatrix} \frac{\partial(w_{11}x_1 + \dots + w_{1n}x_n)}{\partial x_1} & \dots & \frac{\partial(w_{11}x_1 + \dots + w_{1n}x_n)}{\partial x_n} \\ \vdots & \ddots & \vdots \end{bmatrix}$$

$$\mathbf{J}_f(\mathbf{x}_0) = \begin{bmatrix} \frac{\partial(w_{11}x_1 + \dots + w_{1n}x_n)}{\partial x_1} & \dots & \frac{\partial(w_{11}x_1 + \dots + w_{1n}x_n)}{\partial x_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial(w_{m1}x_1 + \dots + w_{mn}x_n)}{\partial x_1} & \dots & \frac{\partial(w_{m1}x_1 + \dots + w_{mn}x_n)}{\partial x_n} \end{bmatrix}$$

$$\begin{aligned}
 \mathbf{J}_f(\mathbf{x}_0) &= \begin{bmatrix} \frac{\partial(w_{11}x_1+\cdots+w_{1n}x_n)}{\partial x_1} & \cdots & \frac{\partial(w_{11}x_1+\cdots+w_{1n}x_n)}{\partial x_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial(w_{m1}x_1+\cdots+w_{mn}x_n)}{\partial x_1} & \cdots & \frac{\partial(w_{m1}x_1+\cdots+w_{mn}x_n)}{\partial x_n} \end{bmatrix} \\
 &= \begin{bmatrix} w_{11} & \cdots & w_{1n} \\ \vdots & \ddots & \vdots \\ w_{m1} & \cdots & w_{mn} \end{bmatrix}
 \end{aligned}$$

$$\begin{aligned}
 \mathbf{J}_f(\mathbf{x}_0) &= \begin{bmatrix} \frac{\partial(w_{11}x_1+\cdots+w_{1n}x_n)}{\partial x_1} & \cdots & \frac{\partial(w_{11}x_1+\cdots+w_{1n}x_n)}{\partial x_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial(w_{m1}x_1+\cdots+w_{mn}x_n)}{\partial x_1} & \cdots & \frac{\partial(w_{m1}x_1+\cdots+w_{mn}x_n)}{\partial x_n} \end{bmatrix} \\
 &= \begin{bmatrix} w_{11} & \cdots & w_{1n} \\ \vdots & \ddots & \vdots \end{bmatrix}
 \end{aligned}$$

$$\begin{aligned}
 \mathbf{J}_f(\mathbf{x}_0) &= \begin{bmatrix} \frac{\partial(w_{11}x_1+\cdots+w_{1n}x_n)}{\partial x_1} & \cdots & \frac{\partial(w_{11}x_1+\cdots+w_{1n}x_n)}{\partial x_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial(w_{m1}x_1+\cdots+w_{mn}x_n)}{\partial x_1} & \cdots & \frac{\partial(w_{m1}x_1+\cdots+w_{mn}x_n)}{\partial x_n} \end{bmatrix} \\
 &= \begin{bmatrix} w_{11} & \cdots & w_{1n} \\ \vdots & \ddots & \vdots \\ w_{m1} & \cdots & w_{mn} \end{bmatrix}
 \end{aligned}$$

$$\begin{aligned}
 \mathbf{J}_f(\mathbf{x}_0) &= \begin{bmatrix} \frac{\partial(w_{11}x_1+\dots+w_{1n}x_n)}{\partial x_1} & \dots & \frac{\partial(w_{11}x_1+\dots+w_{1n}x_n)}{\partial x_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial(w_{m1}x_1+\dots+w_{mn}x_n)}{\partial x_1} & \dots & \frac{\partial(w_{m1}x_1+\dots+w_{mn}x_n)}{\partial x_n} \end{bmatrix} \\
 &= \begin{bmatrix} w_{11} & \dots & w_{1n} \\ \vdots & \ddots & \vdots \\ w_{m1} & \dots & w_{mn} \end{bmatrix} \\
 &= \mathbf{W}
 \end{aligned}$$

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a quadratic map defined by $f(\mathbf{x}) = \mathbf{x}^T \mathbf{A} \mathbf{x}$, where \mathbf{A} is a symmetric matrix of dimension $(n \times n)$.

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a quadratic map defined by $f(\mathbf{x}) = \mathbf{x}^T \mathbf{A} \mathbf{x}$, where \mathbf{A} is a symmetric matrix of dimension $(n \times n)$. Since matrix multiplication is associative we can see f as the multiplication of two matrix \mathbf{x}^T and $\mathbf{y}(\mathbf{x}) = \mathbf{A} \mathbf{x}$.

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a quadratic map defined by $f(\mathbf{x}) = \mathbf{x}^T \mathbf{A} \mathbf{x}$, where \mathbf{A} is a symmetric matrix of dimension $(n \times n)$. Since matrix multiplication is associative we can see f as the multiplication of two matrix \mathbf{x}^T and $\mathbf{y}(\mathbf{x}) = \mathbf{A} \mathbf{x}$. Note that the matrix multiplication is exactly the function composition!

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a quadratic map defined by $f(\mathbf{x}) = \mathbf{x}^T \mathbf{A} \mathbf{x}$, where \mathbf{A} is a symmetric matrix of dimension $(n \times n)$. Since matrix multiplication is associative we can see f as the multiplication of two matrix \mathbf{x}^T and $\mathbf{y}(\mathbf{x}) = \mathbf{A} \mathbf{x}$. Note that the matrix multiplication is exactly the function composition! So $f(\mathbf{x}) = h(\mathbf{x}, \mathbf{y}) = \mathbf{x}^T \mathbf{y}$ and if we apply the chaine rule to h we have:

h

h

x

h

x


y


h


x


y


x




















$$\frac{\partial f(\mathbf{x})}{\partial \mathbf{x}} = \frac{\partial h(\mathbf{x}, \mathbf{y})}{\partial \mathbf{x}}$$

$$\begin{aligned}\frac{\partial f(\mathbf{x})}{\partial \mathbf{x}} &= \frac{\partial h(\mathbf{x}, \mathbf{y})}{\partial \mathbf{x}} \\ &= \frac{\partial h}{\partial \mathbf{x}} \frac{\partial \mathbf{x}}{\partial \mathbf{x}} + \frac{\partial h}{\partial \mathbf{y}} \frac{\partial \mathbf{y}}{\partial \mathbf{x}}\end{aligned}$$

$$\begin{aligned}
 \frac{\partial f(\mathbf{x})}{\partial \mathbf{x}} &= \frac{\partial h(\mathbf{x}, \mathbf{y})}{\partial \mathbf{x}} \\
 &= \frac{\partial h}{\partial \mathbf{x}} \frac{\partial \mathbf{x}}{\partial \mathbf{x}} + \frac{\partial h}{\partial \mathbf{y}} \frac{\partial \mathbf{y}}{\partial \mathbf{x}} \\
 &= \mathbf{y}^T + \mathbf{x}^T \mathbf{A}
 \end{aligned}$$

$$\begin{aligned}
 \frac{\partial f(\mathbf{x})}{\partial \mathbf{x}} &= \frac{\partial h(\mathbf{x}, \mathbf{y})}{\partial \mathbf{x}} \\
 &= \frac{\partial h}{\partial \mathbf{x}} \frac{\partial \mathbf{x}}{\partial \mathbf{x}} + \frac{\partial h}{\partial \mathbf{y}} \frac{\partial \mathbf{y}}{\partial \mathbf{x}} \\
 &= \mathbf{y}^T + \mathbf{x}^T \mathbf{A} \\
 &= (\mathbf{Ax})^T + \mathbf{x}^T \mathbf{A}
 \end{aligned}$$

$$\begin{aligned}
 \frac{\partial f(\mathbf{x})}{\partial \mathbf{x}} &= \frac{\partial h(\mathbf{x}, \mathbf{y})}{\partial \mathbf{x}} \\
 &= \frac{\partial h}{\partial \mathbf{x}} \frac{\partial \mathbf{x}}{\partial \mathbf{x}} + \frac{\partial h}{\partial \mathbf{y}} \frac{\partial \mathbf{y}}{\partial \mathbf{x}} \\
 &= \mathbf{y}^T + \mathbf{x}^T \mathbf{A} \\
 &= (\mathbf{Ax})^T + \mathbf{x}^T \mathbf{A} \\
 &= \mathbf{x}^T \mathbf{A}^T + \mathbf{x}^T \mathbf{A}
 \end{aligned}$$

$$\begin{aligned}
 \frac{\partial f(\mathbf{x})}{\partial \mathbf{x}} &= \frac{\partial h(\mathbf{x}, \mathbf{y})}{\partial \mathbf{x}} \\
 &= \frac{\partial h}{\partial \mathbf{x}} \frac{\partial \mathbf{x}}{\partial \mathbf{x}} + \frac{\partial h}{\partial \mathbf{y}} \frac{\partial \mathbf{y}}{\partial \mathbf{x}} \\
 &= \mathbf{y}^T + \mathbf{x}^T \mathbf{A} \\
 &= (\mathbf{Ax})^T + \mathbf{x}^T \mathbf{A} \\
 &= \mathbf{x}^T \mathbf{A}^T + \mathbf{x}^T \mathbf{A} \\
 &= \mathbf{x}^T (\mathbf{A}^T + \mathbf{A})
 \end{aligned}$$

$$\begin{aligned}
\frac{\partial f(\mathbf{x})}{\partial \mathbf{x}} &= \frac{\partial h(\mathbf{x}, \mathbf{y})}{\partial \mathbf{x}} \\
&= \frac{\partial h}{\partial \mathbf{x}} \frac{\partial \mathbf{x}}{\partial \mathbf{x}} + \frac{\partial h}{\partial \mathbf{y}} \frac{\partial \mathbf{y}}{\partial \mathbf{x}} \\
&= \mathbf{y}^T + \mathbf{x}^T \mathbf{A} \\
&= (\mathbf{Ax})^T + \mathbf{x}^T \mathbf{A} \\
&= \mathbf{x}^T \mathbf{A}^T + \mathbf{x}^T \mathbf{A} \\
&= \mathbf{x}^T (\mathbf{A}^T + \mathbf{A}) \\
&= 2\mathbf{x}^T \mathbf{A}
\end{aligned}$$

The last equality holds because \mathbf{A} is symmetric, i.e., $\mathbf{A}^T = \mathbf{A}$

Let $f : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}^n$ with $f(\mathbf{W}) = \mathbf{W}\mathbf{x}$ for a fixed $\mathbf{x} : \mathbb{R}^n$. Then,

$$f(\mathbf{W}) =$$

Let $f : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}^n$ with $f(\mathbf{W}) = \mathbf{W}\mathbf{x}$ for a fixed $\mathbf{x} : \mathbb{R}^n$. Then,

$$f(\mathbf{W}) = \begin{bmatrix} w_{11} & \dots & w_{1n} \\ \vdots & & \vdots \\ w_{m1} & \dots & w_{mn} \end{bmatrix}$$

Let $f : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}^n$ with $f(\mathbf{W}) = \mathbf{W}\mathbf{x}$ for a fixed $\mathbf{x} : \mathbb{R}^n$. Then,

$$f(\mathbf{W}) = \begin{bmatrix} w_{11} & \dots & w_{1n} \\ \vdots & \ddots & \vdots \end{bmatrix}$$

Let $f : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}^n$ with $f(\mathbf{W}) = \mathbf{W}\mathbf{x}$ for a fixed $\mathbf{x} : \mathbb{R}^n$. Then,

$$f(\mathbf{W}) = \begin{bmatrix} w_{11} & \dots & w_{1n} \\ \vdots & \ddots & \vdots \\ w_{m1} & \dots & w_{mn} \end{bmatrix}$$

Let $f : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}^n$ with $f(\mathbf{W}) = \mathbf{W}\mathbf{x}$ for a fixed $\mathbf{x} : \mathbb{R}^n$. Then,

$$f(\mathbf{W}) = \begin{bmatrix} w_{11} & \dots & w_{1n} \\ \vdots & \ddots & \vdots \\ w_{m1} & \dots & w_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$$

Let $f : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}^n$ with $f(\mathbf{W}) = \mathbf{W}\mathbf{x}$ for a fixed $\mathbf{x} : \mathbb{R}^n$. Then,

$$f(\mathbf{W}) = \begin{bmatrix} w_{11} & \dots & w_{1n} \\ \vdots & \ddots & \vdots \\ w_{m1} & \dots & w_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \end{bmatrix}$$

Let $f : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}^n$ with $f(\mathbf{W}) = \mathbf{W}\mathbf{x}$ for a fixed $\mathbf{x} : \mathbb{R}^n$. Then,

$$f(\mathbf{W}) = \begin{bmatrix} w_{11} & \dots & w_{1n} \\ \vdots & \ddots & \vdots \\ w_{m1} & \dots & w_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$$

Let $f : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}^n$ with $f(\mathbf{W}) = \mathbf{W}\mathbf{x}$ for a fixed $\mathbf{x} : \mathbb{R}^n$. Then,

$$\begin{aligned} f(\mathbf{W}) &= \begin{bmatrix} w_{11} & \dots & w_{1n} \\ \vdots & \ddots & \vdots \\ w_{m1} & \dots & w_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} \\ &= \begin{bmatrix} w_{11}x_1 + \dots + w_{1n}x_n \\ \vdots \\ w_{m1}x_1 + \dots + w_{mn}x_n \end{bmatrix} \end{aligned}$$

Let $f : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}^n$ with $f(\mathbf{W}) = \mathbf{W}\mathbf{x}$ for a fixed $\mathbf{x} : \mathbb{R}^n$. Then,

$$\begin{aligned} f(\mathbf{W}) &= \begin{bmatrix} w_{11} & \dots & w_{1n} \\ \vdots & \ddots & \vdots \\ w_{m1} & \dots & w_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} \\ &= \begin{bmatrix} w_{11}x_1 + \dots + w_{1n}x_n \\ \vdots \end{bmatrix} \end{aligned}$$

Let $f : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}^n$ with $f(\mathbf{W}) = \mathbf{W}\mathbf{x}$ for a fixed $\mathbf{x} : \mathbb{R}^n$. Then,

$$\begin{aligned} f(\mathbf{W}) &= \begin{bmatrix} w_{11} & \dots & w_{1n} \\ \vdots & \ddots & \vdots \\ w_{m1} & \dots & w_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} \\ &= \begin{bmatrix} w_{11}x_1 + \dots + w_{1n}x_n \\ \vdots \\ w_{m1}x_1 + \dots + w_{mn}x_n \end{bmatrix} \end{aligned}$$

$$\mathbf{J}_f(\mathbf{W}_0) = \begin{bmatrix} \frac{\partial(w_{11}x_1 + \dots + w_{1n}x_n)}{\partial w_{11}} & \dots & \frac{\partial(w_{11}x_1 + \dots + w_{1n}x_n)}{\partial w_{mn}} \end{bmatrix}$$

$$\mathbf{J}_f(\mathbf{W}_0) = \begin{bmatrix} \frac{\partial(w_{11}x_1 + \dots + w_{1n}x_n)}{\partial w_{11}} & \dots & \frac{\partial(w_{11}x_1 + \dots + w_{1n}x_n)}{\partial w_{mn}} \\ \vdots & \ddots & \vdots \end{bmatrix}$$

$$\mathbf{J}_f(\mathbf{W}_0) = \begin{bmatrix} \frac{\partial(w_{11}x_1 + \dots + w_{1n}x_n)}{\partial w_{11}} & \dots & \frac{\partial(w_{11}x_1 + \dots + w_{1n}x_n)}{\partial w_{mn}} \\ \vdots & \ddots & \vdots \\ \frac{\partial(w_{m1}x_1 + \dots + w_{mn}x_n)}{\partial w_{11}} & \dots & \frac{\partial(w_{m1}x_1 + \dots + w_{mn}x_n)}{\partial w_{mn}} \end{bmatrix}$$

$$\begin{aligned}
 \mathbf{J}_f(\mathbf{W}_0) &= \begin{bmatrix} \frac{\partial(w_{11}x_1 + \dots + w_{1n}x_n)}{\partial w_{11}} & \dots & \frac{\partial(w_{11}x_1 + \dots + w_{1n}x_n)}{\partial w_{mn}} \\ \vdots & \ddots & \vdots \\ \frac{\partial(w_{m1}x_1 + \dots + w_{mn}x_n)}{\partial w_{11}} & \dots & \frac{\partial(w_{m1}x_1 + \dots + w_{mn}x_n)}{\partial w_{mn}} \end{bmatrix} \\
 &= \begin{bmatrix} x_1 & \dots & x_n & 0 & \dots & 0 & \dots & 0 & \dots & 0 \end{bmatrix}
 \end{aligned}$$

$$\begin{aligned}
 \mathbf{J}_f(\mathbf{W}_0) &= \begin{bmatrix} \frac{\partial(w_{11}x_1 + \dots + w_{1n}x_n)}{\partial w_{11}} & \dots & \frac{\partial(w_{11}x_1 + \dots + w_{1n}x_n)}{\partial w_{mn}} \\ \vdots & \ddots & \vdots \\ \frac{\partial(w_{m1}x_1 + \dots + w_{mn}x_n)}{\partial w_{11}} & \dots & \frac{\partial(w_{m1}x_1 + \dots + w_{mn}x_n)}{\partial w_{mn}} \end{bmatrix} \\
 &= \begin{bmatrix} x_1 & \dots & x_n & 0 & \dots & 0 & \dots & 0 & \dots & 0 \\ 0 & \dots & 0 & x_1 & \dots & x_n & \dots & 0 & \dots & 0 \end{bmatrix}
 \end{aligned}$$

$$\begin{aligned}
 \mathbf{J}_f(\mathbf{W}_0) &= \begin{bmatrix} \frac{\partial(w_{11}x_1 + \dots + w_{1n}x_n)}{\partial w_{11}} & \dots & \frac{\partial(w_{11}x_1 + \dots + w_{1n}x_n)}{\partial w_{mn}} \\ \vdots & \ddots & \vdots \\ \frac{\partial(w_{m1}x_1 + \dots + w_{mn}x_n)}{\partial w_{11}} & \dots & \frac{\partial(w_{m1}x_1 + \dots + w_{mn}x_n)}{\partial w_{mn}} \end{bmatrix} \\
 &= \begin{bmatrix} x_1 & \dots & x_n & 0 & \dots & 0 & \dots & 0 & \dots & 0 \\ 0 & \dots & 0 & x_1 & \dots & x_n & \dots & 0 & \dots & 0 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \ddots & \vdots & \ddots & \vdots \\ 0 & \dots & 0 & 0 & \dots & 0 & \dots & x_1 & \dots & x_n \end{bmatrix}
 \end{aligned}$$

$$\begin{aligned}
\mathbf{J}_f(\mathbf{W}_0) &= \begin{bmatrix} \frac{\partial(w_{11}x_1 + \dots + w_{1n}x_n)}{\partial w_{11}} & \dots & \frac{\partial(w_{11}x_1 + \dots + w_{1n}x_n)}{\partial w_{mn}} \\ \vdots & \ddots & \vdots \\ \frac{\partial(w_{m1}x_1 + \dots + w_{mn}x_n)}{\partial w_{11}} & \dots & \frac{\partial(w_{m1}x_1 + \dots + w_{mn}x_n)}{\partial w_{mn}} \end{bmatrix} \\
&= \begin{bmatrix} x_1 & \dots & x_n & 0 & \dots & 0 & \dots & 0 & \dots & 0 \\ 0 & \dots & 0 & x_1 & \dots & x_n & \dots & 0 & \dots & 0 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \ddots & \vdots & \ddots & \vdots \\ 0 & \dots & 0 & 0 & \dots & 0 & \dots & x_1 & \dots & x_n \end{bmatrix} \\
&= [x_1 \ \dots \ x_n] \otimes \mathbf{I}_m
\end{aligned}$$

$$\begin{aligned}
\mathbf{J}_f(\mathbf{W}_0) &= \begin{bmatrix} \frac{\partial(w_{11}x_1 + \dots + w_{1n}x_n)}{\partial w_{11}} & \dots & \frac{\partial(w_{11}x_1 + \dots + w_{1n}x_n)}{\partial w_{mn}} \\ \vdots & \ddots & \vdots \\ \frac{\partial(w_{m1}x_1 + \dots + w_{mn}x_n)}{\partial w_{11}} & \dots & \frac{\partial(w_{m1}x_1 + \dots + w_{mn}x_n)}{\partial w_{mn}} \end{bmatrix} \\
&= \begin{bmatrix} x_1 & \dots & x_n & 0 & \dots & 0 & \dots & 0 & \dots & 0 \\ 0 & \dots & 0 & x_1 & \dots & x_n & \dots & 0 & \dots & 0 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \ddots & \vdots & \ddots & \vdots \\ 0 & \dots & 0 & 0 & \dots & 0 & \dots & x_1 & \dots & x_n \end{bmatrix} \\
&= [x_1 \ \dots \ x_n] \otimes \mathbf{I}_m \\
&= \mathbf{x} \otimes \mathbf{I}_m
\end{aligned}$$

Thank You