

# Deep Learning

## Lecture 1: Linear Algebra

**Dr. Mehrdad Maleki**

# Sets

- ▶ **Set** is a collection of objects represented as a unit.

# Sets

- ▶ **Set** is a collection of objects represented as a unit.
- ▶ A set with no elements is called the **empty set** and is denoted by  $\emptyset$ . So  $\emptyset = \{ \}$ .

# Sets

- ▶ **Set** is a collection of objects represented as a unit.
- ▶ A set with no elements is called the **empty set** and is denoted by  $\emptyset$ . So  $\emptyset = \{ \}$ .
- ▶ Set  $A = \{a, b, c\}$  is the same as the set  $B = \{b, c, a\}$ .

# Sets

- ▶ **Set** is a collection of objects represented as a unit.
- ▶ A set with no elements is called the **empty set** and is denoted by  $\emptyset$ . So  $\emptyset = \{ \}$ .
- ▶ Set  $A = \{a, b, c\}$  is the same as the set  $B = \{b, c, a\}$ .
- ▶ There is no duplicate element in a set. So  $\{a, b, b, c\}$  is the same as  $\{a, b, c\}$ .

# Basic Concepts

- ▶ If  $x$  is an element of a set  $A$  we write  $x \in A$ .

# Basic Concepts

- ▶ If  $x$  is an element of a set  $A$  we write  $x \in A$ .
- ▶ If  $x$  is not an element of a set  $A$  we write  $x \notin A$ .

# Basic Concepts

- ▶ If  $x$  is an element of a set  $A$  we write  $x \in A$ .
- ▶ If  $x$  is not an element of a set  $A$  we write  $x \notin A$ .
- ▶ If every element of a set  $A$  is also an element of set  $B$  we call  $A$  a **subset** of  $B$  and write  $A \subseteq B$ .



# Important Sets

- ▶ **Natural Numbers:**  $\mathbb{N} = \{1, 2, 3, \dots\}$

# Important Sets

- ▶ **Natural Numbers:**  $\mathbb{N} = \{1, 2, 3, \dots\}$
- ▶ **Integers:**  $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

# Important Sets

- ▶ **Natural Numbers:**  $\mathbb{N} = \{1, 2, 3, \dots\}$
- ▶ **Integers:**  $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$
- ▶ **Rational Numbers:**  $\mathbb{Q} = \{\frac{a}{b} : a, b \in \mathbb{Z}, b \neq 0\}$

# Important Sets

- ▶ **Natural Numbers:**  $\mathbb{N} = \{1, 2, 3, \dots\}$
- ▶ **Integers:**  $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$
- ▶ **Rational Numbers:**  $\mathbb{Q} = \{\frac{a}{b} : a, b \in \mathbb{Z}, b \neq 0\}$
- ▶ **Irrational Numbers:**  $\mathbb{Q}^c = \{x : x \text{ is not rational}\}$

# Important Sets

- ▶ **Natural Numbers:**  $\mathbb{N} = \{1, 2, 3, \dots\}$
- ▶ **Integers:**  $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$
- ▶ **Rational Numbers:**  $\mathbb{Q} = \{\frac{a}{b} : a, b \in \mathbb{Z}, b \neq 0\}$
- ▶ **Irrational Numbers:**  $\mathbb{Q}^c = \{x : x \text{ is not rational}\}$
- ▶ **Real Numbers:**  $\mathbb{R} = \mathbb{Q} \cup \mathbb{Q}^c$

# Hierarchy of Sets

$$\mathbb{N} \subseteq \mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{R} \subseteq \mathbb{C}$$

# Construction

- ▶ Looking for the roots of the equation like  $x(x + 3) = 0$  leads to invention of  $\mathbb{Z}$ .

# Construction

- ▶ Looking for the roots of the equation like  $x(x + 3) = 0$  leads to invention of  $\mathbb{Z}$ .
- ▶ Looking for the roots of the equation like  $2x + 3 = 4$  leads to invention of  $\mathbb{Q}$ .



# Construction

- ▶ Looking for the roots of the equation like  $x(x + 3) = 0$  leads to invention of  $\mathbb{Z}$ .
- ▶ Looking for the roots of the equation like  $2x + 3 = 4$  leads to invention of  $\mathbb{Q}$ .
- ▶ Looking for the roots of the equation like  $x^2 = 2$  leads to invention of  $\mathbb{Q}^c$ .

# Construction

- ▶ Looking for the roots of the equation like  $x(x + 3) = 0$  leads to invention of  $\mathbb{Z}$ .
- ▶ Looking for the roots of the equation like  $2x + 3 = 4$  leads to invention of  $\mathbb{Q}$ .
- ▶ Looking for the roots of the equation like  $x^2 = 2$  leads to invention of  $\mathbb{Q}^c$ .
- ▶ Looking for the roots of the equation like  $x^2 + 1 = 0$  leads to invention of  $\mathbb{C}$ .

# Operations on Sets

► **Union:**  $A \cup B = \{x : x \in A \text{ or } x \in B\}$

# Operations on Sets

- ▶ **Union:**  $A \cup B = \{x : x \in A \text{ or } x \in B\}$
- ▶ **Intersection:**  $A \cap B = \{x : x \in A \text{ and } x \in B\}$

# Operations on Sets

- ▶ **Union:**  $A \cup B = \{x : x \in A \text{ or } x \in B\}$
- ▶ **Intersection:**  $A \cap B = \{x : x \in A \text{ and } x \in B\}$
- ▶ **Complement:**  $\bar{A} = \{x : x \notin A\}$

# Operations on Sets

- ▶ **Union:**  $A \cup B = \{x : x \in A \text{ or } x \in B\}$
- ▶ **Intersection:**  $A \cap B = \{x : x \in A \text{ and } x \in B\}$
- ▶ **Complement:**  $\bar{A} = \{x : x \notin A\}$
- ▶ **Set Minus:**  $A \setminus B = \{x : x \in A \text{ and } x \notin B\}$
- ▶ Definition of complement is depend on a reference set.

# Venn Diagram $A \cup B$




Figure: Union  $A \cup B$

# Venn Diagram $A \cap B$




Figure: Intersection  $A \cap B$



# Venn Diagram $A \setminus B$




Figure: Set minus  $A \setminus B$

# Some Useful Properties

- ▶  $\emptyset \subseteq A$
- ▶  $A \cup \emptyset = A$  and  $A \cap \emptyset = \emptyset$
- ▶ **De Morgan's laws:**

$$\overline{(A \cup B)} = \bar{A} \cap \bar{B}$$

$$\overline{(A \cap B)} = \bar{A} \cup \bar{B}$$

$$\overline{\bar{A}} = A$$

# Set in Python

```
1 empty_set=set()
```

```
1 x={'a','b','c'}
```

```
1 y={'c','d','d','e','f'}
```

```
1 y
```

```
{'c', 'd', 'e', 'f'}
```

# Set in Python

```
1 x.union(y)
```

```
{'a', 'b', 'c', 'd', 'e', 'f'}
```

```
1 x.intersection(y)
```

```
{'c'}
```

```
1 x.difference(y)
```

```
{'a', 'b'}
```

# Set in Python

```
1 'a' in x
```

True

```
1 'a' not in x
```

False

```
1 x.issubset(y)
```

False

# Cartesian Product

- ▶ The **Cartesian Product** of sets  $A$  and  $B$ , denoted  $A \times B$ , is the set of all ordered pairs  $(a, b)$  such that  $a \in A$  and  $b \in B$ , i.e.,

# Cartesian Product

- ▶ The **Cartesian Product** of sets  $A$  and  $B$ , denoted  $A \times B$ , is the set of all ordered pairs  $(a, b)$  such that  $a \in A$  and  $b \in B$ , i.e.,

$$A \times B = \{(a, b) : a \in A \text{ and } b \in B\}$$

# Cartesian Product

- ▶ The **Cartesian Product** of sets  $A$  and  $B$ , denoted  $A \times B$ , is the set of all ordered pairs  $(a, b)$  such that  $a \in A$  and  $b \in B$ , i.e.,

$$A \times B = \{(a, b) : a \in A \text{ and } b \in B\}$$

- ▶ If  $A = \{Joe, Philip\}$  and  $B = \{Daniel, Peter, Joe\}$  then:



# Cartesian Product

- ▶ The **Cartesian Product** of sets  $A$  and  $B$ , denoted  $A \times B$ , is the set of all ordered pairs  $(a, b)$  such that  $a \in A$  and  $b \in B$ , i.e.,

$$A \times B = \{(a, b) : a \in A \text{ and } b \in B\}$$

- ▶ If  $A = \{Joe, Philip\}$  and  $B = \{Daniel, Peter, Joe\}$  then:

$$A \times B = \{(Joe, Daniel), (Joe, Peter), (Joe, Joe), \\ (Philip, Daniel), (Philip, Peter), (Philip, Joe)\}$$

# Cartesian Product

- ▶ The **Cartesian Product** of sets  $A$  and  $B$ , denoted  $A \times B$ , is the set of all ordered pairs  $(a, b)$  such that  $a \in A$  and  $b \in B$ , i.e.,

$$A \times B = \{(a, b) : a \in A \text{ and } b \in B\}$$

- ▶ If  $A = \{Joe, Philip\}$  and  $B = \{Daniel, Peter, Joe\}$  then:

$$A \times B = \{(Joe, Daniel), (Joe, Peter), (Joe, Joe), \\ (Philip, Daniel), (Philip, Peter), (Philip, Joe)\}$$

- ▶  $A \times B \neq B \times A$

# Cartesian Product

- ▶ The **Cartesian Product** of sets  $A$  and  $B$ , denoted  $A \times B$ , is the set of all ordered pairs  $(a, b)$  such that  $a \in A$  and  $b \in B$ , i.e.,

$$A \times B = \{(a, b) : a \in A \text{ and } b \in B\}$$

- ▶ If  $A = \{Joe, Philip\}$  and  $B = \{Daniel, Peter, Joe\}$  then:

$$A \times B = \{(Joe, Daniel), (Joe, Peter), (Joe, Joe), \\ (Philip, Daniel), (Philip, Peter), (Philip, Joe)\}$$

- ▶  $A \times B \neq B \times A$

- ▶ A **relation** between  $A$  and  $B$  is a subset of  $A \times B$ .

# Functions

- ▶ A **function** is a relation between two sets such that each element of the first set is related to at most one element of the second set.

# Functions

- ▶ A **function** is a relation between two sets such that each element of the first set is related to at most one element of the second set.
- ▶ Temperature of a cup of coffee is a function of time.
- ▶ Stock price with respect to time.
- ▶ Area of a circle with respect to its radius.

# Formal Definition

- ▶ A function  $f$  from set  $A$  to set  $B$ , denoted  $f : A \rightarrow B$ , is a subset of  $A \times B$  such that:

# Formal Definition

- ▶ A function  $f$  from set  $A$  to set  $B$ , denoted  $f : A \rightarrow B$ , is a subset of  $A \times B$  such that:

$$(a_1, b_1), (a_2, b_2) \in f \text{ and } a_1 = a_2 \implies b_1 = b_2$$

# Formal Definition

- ▶ A function  $f$  from set  $A$  to set  $B$ , denoted  $f : A \rightarrow B$ , is a subset of  $A \times B$  such that:

$$(a_1, b_1), (a_2, b_2) \in f \text{ and } a_1 = a_2 \implies b_1 = b_2$$

- ▶ If  $(a, b) \in f$  we write  $f(a) = b$ .



# Formal Definition

- ▶ A function  $f$  from set  $A$  to set  $B$ , denoted  $f : A \rightarrow B$ , is a subset of  $A \times B$  such that:

$$(a_1, b_1), (a_2, b_2) \in f \text{ and } a_1 = a_2 \implies b_1 = b_2$$

- ▶ If  $(a, b) \in f$  we write  $f(a) = b$ .
- ▶ So if  $a_1 = a_2$  then  $f(a_1) = f(a_2)$ .


An vector is a mathematical object which has a **magnitude** and a **direction**:

An vector is a mathematical object which has a **magnitude** and a **direction**:




In this case we refer to this vector as  $\overrightarrow{AB}$ .

An vector is a mathematical object which has a **magnitude** and a **direction**:




In this case we refer to this vector as  $\overrightarrow{AB}$ .

An vector is a mathematical object which has a **magnitude** and a **direction**:



In this case we refer to this vector as  $\overrightarrow{AB}$ .

An vector is a mathematical object which has a **magnitude** and a **direction**:



In this case we refer to this vector as  $\overrightarrow{AB}$ .

We can add two vectors:

We can add two vectors:






We can add two vectors:



We can add two vectors:



We can add two vectors:




—  $u$ :

$-u:$



$-u$ :




$$v - u = v + (-u) = w:$$

$$v - u = v + (-u) = w:$$







$$v - u = v + (-u) = w:$$




$$v - u = v + (-u) = w:$$




$$v - u = v + (-u) = w:$$



$$v - u = v + (-u) = w:$$





$$v - u = v + (-u) = w:$$




We can consider a reference and put the starting point of all vectors in this reference point, call it **O**. So in two dimensions we will write a vector as:


$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

















# Algebra of Vectors

$$\mathbf{0} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

# Algebra of Vectors

$$\mathbf{0} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} x_1 + x'_1 \\ x_2 + x'_2 \end{bmatrix}$$

# Algebra of Vectors

$$\mathbf{0} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} x_1 + x'_1 \\ x_2 + x'_2 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} - \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} x_1 - x'_1 \\ x_2 - x'_2 \end{bmatrix}$$

# Algebra of Vectors

$$\mathbf{0} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} x_1 + x'_1 \\ x_2 + x'_2 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} - \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} x_1 - x'_1 \\ x_2 - x'_2 \end{bmatrix}$$

$$c \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} cx_1 \\ cx_2 \end{bmatrix}$$



# $n$ -dimensional vectors

The set of all  $n$ -dimensional vectors,

# n-dimensional vectors

The set of all  $n$ -dimensional vectors,

$$\begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$$

# n-dimensional vectors

The set of all  $n$ -dimensional vectors,

$$\begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$$

is denoted by  $\mathbb{R}^n$ .

# Euclidean Length of Vectors

The Euclidean length of a vector,

# Euclidean Length of Vectors

The Euclidean length of a vector,

$$u = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$$

# Euclidean Length of Vectors

The Euclidean length of a vector,

$$u = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$$

is denoted by  $\|u\|$  and is a real number that represents the length of  $u$  and is calculated by,

# Euclidean Length of Vectors

The Euclidean length of a vector,

$$u = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$$

is denoted by  $\|u\|$  and is a real number that represents the length of  $u$  and is calculated by,

$$\|u\| = \sqrt{x_1^2 + \cdots + x_n^2}$$

# Inner Product

The **inner product** of two vectors  $\mathbf{u}$  and  $\mathbf{v}$  is a real number defined by the following formula,



# Inner Product

The **inner product** of two vectors  $\mathbf{u}$  and  $\mathbf{v}$  is a real number defined by the following formula,


$$\mathbf{u} \cdot \mathbf{v} = \|\mathbf{u}\| \|\mathbf{v}\| \cos(\theta)$$

# Inner Product

The **inner product** of two vectors  $\mathbf{u}$  and  $\mathbf{v}$  is a real number defined by the following formula,

$$\mathbf{u} \cdot \mathbf{v} = \|\mathbf{u}\| \|\mathbf{v}\| \cos(\theta)$$

where  $\theta$  is the angel between  $\mathbf{u}$  and  $\mathbf{v}$ .



If,

If,

$$\mathbf{u} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}, \mathbf{v} = \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix}$$

If,

$$\mathbf{u} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}, \mathbf{v} = \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix}$$

then

If,

$$\mathbf{u} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}, \mathbf{v} = \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix}$$

then

$$\mathbf{u} \cdot \mathbf{v} = x_1 y_1 + \cdots + x_n y_n$$

If  $\mathbf{u} \cdot \mathbf{v} = 0$  then  $\mathbf{u}$  is perpendicular with  $\mathbf{v}$ .

If  $\mathbf{u} \cdot \mathbf{v} = 0$  then  $\mathbf{u}$  is perpendicular with  $\mathbf{v}$ .






Let  $\|\mathbf{u}\| = 1$  then  $\mathbf{u} \cdot \mathbf{v}$  is the length of projection of  $\mathbf{v}$  in the direction of  $\mathbf{u}$ ,


Let  $\|\mathbf{u}\| = 1$  then  $\mathbf{u} \cdot \mathbf{v}$  is the length of projection of  $\mathbf{v}$  in the direction of  $\mathbf{u}$ ,




Let  $\|\mathbf{u}\| = 1$  then  $\mathbf{u} \cdot \mathbf{v}$  is the length of projection of  $\mathbf{v}$  in the direction of  $\mathbf{u}$ ,




Let  $\|\mathbf{u}\| = 1$  then  $\mathbf{u} \cdot \mathbf{v}$  is the length of projection of  $\mathbf{v}$  in the direction of  $\mathbf{u}$ ,




Let  $\|\mathbf{u}\| = 1$  then  $\mathbf{u} \cdot \mathbf{v}$  is the length of projection of  $\mathbf{v}$  in the direction of  $\mathbf{u}$ ,




Let  $\|\mathbf{u}\| = 1$  then  $\mathbf{u} \cdot \mathbf{v}$  is the length of projection of  $\mathbf{v}$  in the direction of  $\mathbf{u}$ ,



Let  $\|\mathbf{u}\| = 1$  then  $\mathbf{u} \cdot \mathbf{v}$  is the length of projection of  $\mathbf{v}$  in the direction of  $\mathbf{u}$ ,




Let  $\|\mathbf{u}\| = 1$  then  $\mathbf{u} \cdot \mathbf{v}$  is the length of projection of  $\mathbf{v}$  in the direction of  $\mathbf{u}$ ,







Let  $\|\mathbf{u}\| = 1$  then  $\mathbf{u} \cdot \mathbf{v}$  is the length of projection of  $\mathbf{v}$  in the direction of  $\mathbf{u}$ ,




Let  $\|\mathbf{u}\| = 1$  then  $\mathbf{u} \cdot \mathbf{v}$  is the length of projection of  $\mathbf{v}$  in the direction of  $\mathbf{u}$ ,



Let  $\|\mathbf{u}\| = 1$  then  $\mathbf{u} \cdot \mathbf{v}$  is the length of projection of  $\mathbf{v}$  in the direction of  $\mathbf{u}$ ,



Let  $\|\mathbf{u}\| = 1$  then  $\mathbf{u} \cdot \mathbf{v}$  is the length of projection of  $\mathbf{v}$  in the direction of  $\mathbf{u}$ ,



So,  $\mathbf{u} \cdot \mathbf{v}$  is maximum if  $\mathbf{v}$  is in the same direction as  $\mathbf{u}$ .

So,  $\mathbf{u} \cdot \mathbf{v}$  is maximum if  $\mathbf{v}$  is in the same direction as  $\mathbf{u}$ .



# Vectors in Python

```
1 u=np.array([1,2,3])
```

```
1 v=np.array([4,5,6])
```

```
1 np.dot(u,v)
```

```
32
```

```
1 np.linalg.norm(u)
```

```
3.7416573867739413
```

# Vectors in Python

```
1 t=np.inner(u,v)/(np.linalg.norm(u)*np.linalg.norm(v))
```

```
1 np.degrees(np.arccos(t))
```

```
12.933154491899135
```



# Matrices

Matrix is a two-dimensional array-like data structure that represents a common quantity of two categories. For example, if you want to represent the number of students by gender in three different class of Math, CS, and Physic then you should use matrix representation as follow;

	<i>Math</i>	<i>CS</i>	<i>Physic</i>
<i>Men</i>	23	37	18
<i>Women</i>	28	14	16

If a matrix has  $m$  rows and  $n$  columns then we say the **order** of the matrix is  $m \times n$ . So an  $n$ -dimensional vector is a special case of a matrix of order  $n \times 1$ .

A **square matrix** is a matrix that have the same number of rows and columns, i.e.,  $m = n$ .

A **square matrix** is a matrix that have the same number of rows and columns, i.e.,  $m = n$ .

An **identity matrix** is a square matrix with 1 in the main diagonal and 0 elsewhere. We denote the identity matrix of order  $n$  by  $\mathbf{I}_n$ .

A **square matrix** is a matrix that have the same number of rows and columns, i.e.,  $m = n$ .

An **identity matrix** is a square matrix with 1 in the main diagonal and 0 elsewhere. We denote the identity matrix of order  $n$  by  $\mathbf{I}_n$ .

$$\mathbf{I}_n = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{bmatrix}$$

The **zero matrix** is a matrix such that every entries is 0. We denote the zero matrix of order  $m \times n$  by  $\mathbf{O}_{m \times n}$ .

The **zero matrix** is a matrix such that every entries is 0. We denote the zero matrix of order  $m \times n$  by  $\mathbf{O}_{m \times n}$ .

$$\mathbf{O}_{m \times n} = \begin{bmatrix} 0 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 \end{bmatrix}$$

Transpose of an  $m \times n$  matrix is an  $n \times m$  matrix. The transpose of the matrix  $\mathbf{A}$  is denoted by  $\mathbf{A}^T$ .



Transpose of an  $m \times n$  matrix is an  $n \times m$  matrix. The transpose of the matrix  $\mathbf{A}$  is denoted by  $\mathbf{A}^T$ .

$$\mathbf{A} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$$

Transpose of an  $m \times n$  matrix is an  $n \times m$  matrix. The transpose of the matrix  $\mathbf{A}$  is denoted by  $\mathbf{A}^T$ .

$$\mathbf{A} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$$

$$\mathbf{A}^T = \begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$$

We could add or subtract two matrices of the same order by adding or subtracting the corresponding entries together.

We could add or subtract two matrices of the same order by adding or subtracting the corresponding entries together.

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} +$$

We could add or subtract two matrices of the same order by adding or subtracting the corresponding entries together.

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} + \begin{bmatrix} 3 & 0 & 1 \\ 2 & -1 & 2 \end{bmatrix} =$$

We could add or subtract two matrices of the same order by adding or subtracting the corresponding entries together.

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} + \begin{bmatrix} 3 & 0 & 1 \\ 2 & -1 & 2 \end{bmatrix} = \begin{bmatrix} 1+3 & 2+0 & 3+1 \\ 4+2 & 5+(-1) & 6+2 \end{bmatrix}$$

If  $\mathbf{A}_{1 \times n}$  and  $\mathbf{B}_{n \times 1}$  then we could define  $\mathbf{AB}$  as follow,

If  $\mathbf{A}_{1 \times n}$  and  $\mathbf{B}_{n \times 1}$  then we could define  $\mathbf{AB}$  as follow,

$$\begin{bmatrix} a_1 & \dots & a_n \end{bmatrix} \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix} = a_1 b_1 + \dots + a_n b_n$$

So if  $\mathbf{u}$  and  $\mathbf{v}$  are  $n$ -dimensional vectors then,



If  $\mathbf{A}_{1 \times n}$  and  $\mathbf{B}_{n \times 1}$  then we could define  $\mathbf{AB}$  as follow,

$$\begin{bmatrix} a_1 & \dots & a_n \end{bmatrix} \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix} = a_1 b_1 + \dots + a_n b_n$$

So if  $\mathbf{u}$  and  $\mathbf{v}$  are  $n$ -dimensional vectors then,

$$\mathbf{u} \cdot \mathbf{v} = \mathbf{u}^T \mathbf{v}$$

If  $\mathbf{A}_{m \times n}$  and  $\mathbf{B}_{n \times k}$  then we could define  $\mathbf{AB}$  as the following  $m \times k$  matrix,

$$\begin{bmatrix} a_{11} & \dots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{m1} & \dots & a_{mn} \end{bmatrix} \begin{bmatrix} b_{11} & \dots & b_{1k} \\ \vdots & \ddots & \vdots \\ b_{n1} & \dots & b_{nk} \end{bmatrix} = \begin{bmatrix} \vdots & & & \\ \dots & [a_{i1} & \dots & a_{in}] & \begin{bmatrix} b_{1j} \\ \vdots \\ b_{nj} \end{bmatrix} & \dots \\ \vdots & & & \end{bmatrix}$$

# Inverse of Matrix

If for the square matrix **A** there exists another square matrix **B** such that,

$$\mathbf{AB} = \mathbf{BA} = \mathbf{I}$$

then we say **A** is invertible and its inverse is **B**. We denote **B** by  $\mathbf{A}^{-1}$ .

If **A** is invertible then the system of equations,

$$\mathbf{Ax} = \mathbf{b}$$

has a unique solution, i.e.,

$$\mathbf{x} = \mathbf{A}^{-1}\mathbf{b}$$

$$\mathbf{IA} = \mathbf{AI} = \mathbf{A}$$

$$\mathbf{I}\mathbf{A} = \mathbf{A}\mathbf{I} = \mathbf{A}$$

$$(\mathbf{A} + \mathbf{B})^T = \mathbf{A}^T + \mathbf{B}^T$$

$$\mathbf{I}\mathbf{A} = \mathbf{A}\mathbf{I} = \mathbf{A}$$

$$(\mathbf{A} + \mathbf{B})^T = \mathbf{A}^T + \mathbf{B}^T$$

$$(\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T$$

$$\mathbf{I}\mathbf{A} = \mathbf{A}\mathbf{I} = \mathbf{A}$$

$$(\mathbf{A} + \mathbf{B})^T = \mathbf{A}^T + \mathbf{B}^T$$

$$(\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T$$

A symmetric matrix is a square matrix such that  $\mathbf{A}^T = \mathbf{A}$ .





# Tensors

**Tensors** are multi-dimensional array with a uniform type. A tensor of order  $m \times n \times k$  (or shape  $(m, n, k)$ ) is actually  $m$  matrices of order  $n \times k$ .

$$\begin{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \\ \begin{bmatrix} 7 & 8 & 9 \\ 10 & 11 & 12 \\ 13 & 14 & 15 \\ 16 & 17 & 18 \\ 19 & 20 & 21 \\ 22 & 23 & 24 \end{bmatrix} \end{bmatrix}$$

is a tensor of order  $4 \times 2 \times 3$

$3 \times 2 \times 3$  tensor.



# Kronecker(Tensor) Product

The **Kronecker product** of matrices  $\mathbf{A}_{m \times n}$  and  $\mathbf{B}_{k \times l}$  is a matrix of order  $mk \times nl$  and define as follow,

$$\mathbf{A} \otimes \mathbf{B} = \begin{bmatrix} a_{11}\mathbf{B} & \dots & a_{1n}\mathbf{B} \\ \vdots & \ddots & \vdots \\ a_{m1}\mathbf{B} & \dots & a_{mn}\mathbf{B} \end{bmatrix}$$

$$\begin{aligned}
 \begin{bmatrix} x_1 & x_2 \end{bmatrix} \otimes \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} &= \begin{bmatrix} x_1 \mathbf{I} & x_2 \mathbf{I} \end{bmatrix} \\
 &= \begin{bmatrix} x_1 & 0 & x_2 & 0 \\ 0 & x_1 & 0 & x_2 \end{bmatrix}
 \end{aligned}$$

# Eigenvalue

Let  $\mathbf{A}$  be an square  $n \times n$  matrix. If there exists  $n$ -dimensional vector  $\mathbf{v}$  and a real number  $\lambda$  such that,

$$\mathbf{A}\mathbf{v} = \lambda\mathbf{v}$$

then we say  $\mathbf{v}$  is an eigenvector and the corresponding eigenvalue is  $\lambda$ .

To obtain the eigenvalues of a matrix you need to find the roots of the following polynomial equation of order  $n$ ,

To obtain the eigenvalues of a matrix you need to find the roots of the following polynomial equation of order  $n$ ,

$$\det(\lambda \mathbf{I} - \mathbf{A}) = 0$$



To obtain the eigenvalues of a matrix you need to find the roots of the following polynomial equation of order  $n$ ,

$$\det(\lambda \mathbf{I} - \mathbf{A}) = 0$$

eigenvalue could be complex number, i.e., a matrix may don't have any real valued eigenvalue but

To obtain the eigenvalues of a matrix you need to find the roots of the following polynomial equation of order  $n$ ,

$$\det(\lambda \mathbf{I} - \mathbf{A}) = 0$$

eigenvalue could be complex number, i.e., a matrix may don't have any real valued eigenvalue but all eigenvalues of a symmetric matrix are real numbers.

# Rotation Matrix

$$\mathbf{R}_\theta = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

# Rotation Matrix

$$\mathbf{R}_\theta = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$



# Rotation Matrix


$$\mathbf{R}_\theta = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$




$\mathbf{R}_{\frac{\pi}{2}}$  has no eigenvalue but  $\mathbf{R}_{\frac{\pi}{2}}^2$  has  $-1$  as the only real eigenvalue.



$\mathbf{R}_{\frac{\pi}{2}}$  has no eigenvalue but  $\mathbf{R}_{\frac{\pi}{2}}^2$  has  $-1$  as the only real eigenvalue.



$\mathbf{R}_{\frac{\pi}{2}}$  has no eigenvalue but  $\mathbf{R}_{\frac{\pi}{2}}^2$  has  $-1$  as the only real eigenvalue.





# Matrices in Python

```
1 A=np.array([[1,2,3],  
2             [4,5,6]])
```

```
1 A.shape
```

```
(2, 3)
```

```
1 A.T
```

```
array([[1, 4],  
       [2, 5],  
       [3, 6]])
```

# Matrices in Python

```
1 A=np.array([[1,2,3],  
2             [4,5,6]])
```

```
1 B=np.array([[1,0,1],  
2             [0,1,0],  
3             [1,1,1]])
```

```
1 np.dot(A,B).shape
```

```
(2, 3)
```

```
1 np.dot(A,B)
```

```
array([[ 4,  5,  4],  
       [10, 11, 10]])
```

```
1 np.matmul(A,B)
```

```
array([[ 4,  5,  4],  
       [10, 11, 10]])
```

# Matrices in Python

```
1 np.identity(3)
```

```
array([[1., 0., 0.],  
       [0., 1., 0.],  
       [0., 0., 1.]])
```

```
1 np.zeros((3,4))
```

```
array([[0., 0., 0., 0.],  
       [0., 0., 0., 0.],  
       [0., 0., 0., 0.]])
```

# Matrices in Python

```
1 Z=np.array([[1,1,0],  
2             [0,1,1],  
3             [1,0,1]])
```

```
1 np.linalg.inv(Z)
```

```
array([[ 0.5, -0.5,  0.5],  
       [ 0.5,  0.5, -0.5],  
       [-0.5,  0.5,  0.5]])
```

```
1 np.linalg.det(Z)
```

```
2.0
```

# Matrices in Python

```
1 X=np.array([[2,3]])  
2 Y=np.identity(2)
```

```
1 np.kron(X,Y)
```

```
array([[2., 0., 3., 0.],  
       [0., 2., 0., 3.]])
```

# Matrices in Python

```
1 np.kron(np.identity(3),np.zeros((4,5))+3)
```

```
array([[3., 3., 3., 3., 3., 0., 0., 0., 0., 0., 0., 0., 0., 0., 0.],  
       [3., 3., 3., 3., 3., 0., 0., 0., 0., 0., 0., 0., 0., 0., 0.],  
       [3., 3., 3., 3., 3., 0., 0., 0., 0., 0., 0., 0., 0., 0., 0.],  
       [3., 3., 3., 3., 3., 0., 0., 0., 0., 0., 0., 0., 0., 0., 0.],  
       [0., 0., 0., 0., 0., 3., 3., 3., 3., 3., 0., 0., 0., 0., 0.],  
       [0., 0., 0., 0., 0., 3., 3., 3., 3., 3., 0., 0., 0., 0., 0.],  
       [0., 0., 0., 0., 0., 3., 3., 3., 3., 3., 0., 0., 0., 0., 0.],  
       [0., 0., 0., 0., 0., 3., 3., 3., 3., 3., 0., 0., 0., 0., 0.],  
       [0., 0., 0., 0., 0., 0., 0., 0., 0., 0., 0., 3., 3., 3., 3.],  
       [0., 0., 0., 0., 0., 0., 0., 0., 0., 0., 0., 3., 3., 3., 3.],  
       [0., 0., 0., 0., 0., 0., 0., 0., 0., 0., 0., 3., 3., 3., 3.],  
       [0., 0., 0., 0., 0., 0., 0., 0., 0., 0., 0., 3., 3., 3., 3.]])
```

# Matrices in Python

```
1 A=np.array([[0,1,1],
2             [1,0,1],
3             [1,1,0]])
```

```
1 np.linalg.eigvals(A)
```

```
array([-1.,  2., -1.])
```

```
1 np.linalg.eig(A)
```

```
(array([-1.,  2., -1.]), array([[-0.81649658,  0.57735027,  0.19219669],
  [ 0.40824829,  0.57735027, -0.7833358 ],
  [ 0.40824829,  0.57735027,  0.59113912]]))
```

**eigenvalues**

**eigenvectors**

*Thank You*