

Release Note : This work describes an approach to covid-19 recognizing that severity in old age as an important clue to the disease process. Known and suspected disease processes are considered in terms of the literature on aging leading to a set of nutritional interventions that may effect both conditions. Initial personal experiences optimizing dog diets have demonstrated some encouraging results but organizing the literature has become an endless task. In the meantime, progress against covid-19 using conventional approaches appears to have reached a dead end. Strikingly absent from recent literature is any mention of vitamin K even when discussing TAM [120]. Tests of speculative ideas on aging may be easier and more productively performed against covid-19 so I decided to release this early "notes" version as a basis for discussion. The more controversial ideas are mostly the result of re-evaluation of some old ideas in light of new findings. Continued literature searches seem to confirm the plausibility of the motivation but it is difficult to come up with a convincing argument for the more controversial issues without empirical trial and error. A goal of the work is to find simple informative and safe tests at this point. Two speculative ideas about SARS-Cov-2 and aging converged on leaking plasma membranes- that is, loss of tryptophan and high infidelity translation products that leak working with the action of NEDD4 on ACE2 to enhance viral entry in an attempt to regulate cytoplasmic ion concentrations. These ideas may not have merit but the similarities between conditions may have different reasons for existing while still pointing to same nutrients. This work should present enough information for the formulation of informative experiments to explore the ideas put forth.

On the age distribution of SARS-Cov-2 Patients

Mike Marchywka*

306 Charles Cox, Canton GA 30115

(Dated: July 1, 2020)

The covid-19 disease caused by SARS-Cov-2 is often fatal, difficult to treat, and difficult to control without disruptive social actions such as distancing and lockdown which have detrimental effects of their own. The search for a better immediate solution has generated speculation on severity modulators among existing anti-viral and host directed approaches based on experience with similar diseases. One robust feature of the epidemic is a preference for older people leading to interest in informative theories of aging that may suggest virus mitigation strategies. SARS-Cov-2 is therefore a good test case for nascent theories on aging such as my ongoing work involving nutrient limitations. In normal aging, nutrient issues may be largely due to primary GI problems usually described as "asymptomatic" but reducing amino acid, cation, and fat soluble vitamin absorption. In the diseased state, host reactions place additional demands on existing nutrients while others may be programmatically destroyed suggesting that covid-19 patients with lower reserves will have worse outcomes. Many of the intervention ideas derived from my aging theory differ from conventional expectations but rely mostly on nutrient manipulation making them reasonably safe and easy for discriminating tests. Despite all the literature on SARS-Cov-2 some important gaps remain related to nutrient issues. ACE2 dynamics related to B0AT and NEDD4 seem to be ignored but are mentioned here. Coagulopathies are reported but neutrophil extracellular traps and protein C have been almost ignored. Protein C will be considered here while discussing a role for vitamin K.

The nutrient risks are both absolute deficiency and imbalance due to a prolonged disease oriented metabolism. I describe candidates for the most pathogenic virus related deficiencies among amino acids (tryptophan, tyrosine, and arginine), metals (mostly copper) and miscellaneous but critical nutrients such as vitamin K and citric acid.

Contents

1. Introduction	3
2. SARS-Cov-2 Relevant features, epidemiology, possible nutritional involvement based on similar diseases	4
2.1. General features	4
2.2. Nutritional issues in covid-19 and related conditions	5
3. Limitations of conventional analyses	6

*Electronic address: marchywka@hotmail.com; to cite or credit this work, see bibtex in Appendix E

4. Overview of my "Old Age as Starvation" work	7
5. Discussion - Approaching SARS-Cov-2 as a nutritional disease	8
5.1. ACE2 and age related conditions	9
5.2. Citric acid	9
5.3. Amino acids	10
5.4. Trace nutrients normally associated with protein	12
5.5. Metals, atomic cations	12
5.6. Fat solubles : A,D,E,K	13
5.7. Integration	14
6. Conclusions	16
7. Supplemental information	16
7.1. Computer code	16
8. Bibliography	16
References	16
Acknowledgments	30
A. Statement of conflicts	31
B. About the author	31
C. Symbols, abbreviations and colloquialisms	31
D. General caveats and disclaimer	31
E. Citing this as a tech report or white paper	32

1. INTRODUCTION

The recently emerged coronavirus, SARS-Cov-2, targets old people preferentially beyond what is seen in similar viral diseases [204]. While some SARS-Cov-2 related conditions have emerged targeting children [212], their relation to the virus remains unclear. With no particularly promising cure in the known pipelines, it may be helpful to understand the characteristics of old age that lead to this distribution. In an ongoing work, I outline some characteristics of old age in dogs and humans analyzed as nutrient issues [141]. The features described in [141] generally relate to effective malnutrition that may cause the same age-related conditions that are prognostic for SARS-Cov-2 virulence with old age as a persistent predictor [226]. The nutrients highlighted in [141] are known to the virology and immunology communities but roles may be unappreciated by conventional thinking. The different predictions regarding nutrient supplements make a good test of my theory and may help resolve some current limitations in progress against SARS-Cov-2. Infectious disease provokes sudden loss of select nutrients, the aging theory addresses a more gradual loss of a wider range of nutrients at varying rates. So, different effects will be observed even if the causes are related and tests need to consider this among other factors. The effective malnutrition in both cases may not resemble classical forms such as food deprivation but that can be rationalized by the nature of the defects and their time evolution.

Several scientific publications have come out with nutritional approaches to covid-19 and many likely exist in other literature sources but none have been proven successful and none appear identical to this work. At least one group has come up with a dietary strategy for covid-19 but methodology and evaluation are unclear from the abstract [214] and there is also a more recent suggestion for a plant-based immunomodulatory diet [50]. One recent work discussed a link between aging and SARS-Cov-2 looking for pathways of interest, even mentioning mTOR [179], but did not apparently recommend the supplements suggested here. Nicotinamide or niacin supplementation was considered during an analysis of NAD fate in SARS-Cov-2 cases [81]. One work does specifically suggest copper for SARS-Cov-2 patients while warning about zinc [172] following some of the reasoning here but somewhat drawing on disinfectant properties and apparently not mentioning clotting issues and nutritional immunity. Another work describes a possible benefit from copper and N-acetylcysteine based on a literature search [10]. The present work is almost orthogonal to works such as [234] that promote vitamins D and E to the exclusion of those noted here. In fact, vitamin E is considered potentially deleterious.

While metabolic changes in the disease state always make nutrition a concern, some recent reports should further motivate an interest in nutritional status for SARS-Cov-2 infected patients. One study of gene expression in bronchoalveolar lavage fluid from 3 groups of people demonstrated maximal down regulation of a group called "GPCR-Mediate Nutrient Sensing in Enteroendocrine Cells" in SARS-Cov-2 patients [237] (see for example ¹). Another report observed that ileal absorptive cells express ACE2 and appear receptive to infection [238] suggesting some sudden loss of nutrient uptake was possible. In common with other coronaviruses, SARS-Cov-2 uses ACE2 for entry but as ACE2 is also a chaperone for an amino acid transporter that transports tryptophan [230] anything that modulates its dynamics may relate to nutrient flow. Also specific to SARS-Cov-2 was the finding of unusual alterations in amino acid metabolism and down regulated platelet factor 4 along with lipid and steroid changes [196].

Currently, the SARS-Cov-2 literature does not address several specific features that may help discriminate among nutritional treatment candidates. In particular, absent is much about NEDD4 and ACE2 dynamics despite NEDD4's involvement in processes related to some prognostic conditions. NEDD4 senses intracellular ions and can internalize ACE2 [160] although the literature on SARS-Cov-2 [193] and earlier coronaviruses [217] does not mention it. As the endogenous ligands may not provoke internalization (internalization in response to Angiotensin-II appears to be mediated by another receptor [39]) it is unclear if the virus itself can do that without help. NEDD4 is an attractive speculative explanation as it senses a common attribute of many predisposing conditions. A very recent comment mentions TAM and protein S but does not discuss vitamin K much [120]. Literature on protein C and the much of vitamin K dependent components are absent despite coagulopathies, the prognostic significance of protein C in ARDS [144] and the cutaneous manifestations possibly suggestive of their relevance. Nor is there much about NETs despite recurring issues with hyperactivated neutrophils (or EET's and depleted eosinophils) and coagulopathies [145] and correspondingly no mention of a role for DNA-ase [240] [98]. In the absence of more information like this, or direct measures of local reduced vitamin K or tryptophan, empirically testing nutrient therapies may be worthwhile due to simplicity and informative value.

The work is organized a bit redundantly to make each section coherent but not complete in itself. The situations in disease and old age are outlined prior to a discussion of nutrients they motivate for covid-19 management. Note that many of the steps in going from observations to logical interventions are empirical. The hypothesized details need

¹ <https://targetexplorer.ingenuity.com/pathway/ING/ING:crzcc#1/api/rest/v1/client/searchPathwayNodes?pathwayId=ING:crzcc&rows=0&facetLimit=5000&responseType=default>

not be entirely accurate for the approach to be informative and even effective.

2. SARS-COV-2 RELEVANT FEATURES, EPIDEMIOLOGY, POSSIBLE NUTRITIONAL INVOLVEMENT BASED ON SIMILAR DISEASES

2.1. General features

Generally studies have found increased severity or mortality in older people or younger people mostly with identifiable risk factors similar to diseases of old age. One work concluded that an age of 60 was a bad prognostic factor [129]. One study of Japanese patients conclude various metrics were worst for people between 50-69 years old [150]. A recent CDC report broke down statistics by age and state showing hospitalizations per capita increasing with age from about 2 per 100,000 in the 18-49 group and almost 8 in the 50-64 approaching 17 for the oldest [61]. Conditions predisposing to severe disease include diabetes [102], cardiovascular diseases [23], hypertension and kidney disease [216]. Males may get more severe disease than females [191] and are generally known to have shorter life expectancies. Smoking is thought to reduce the risk leading to interest in nicotine [51] although physical barriers to infection may need to be considered.

Skin conditions have been variously described in the popular press as rashes or "covid toes". Some consider that urticaria with pyrexia may be a common early symptom [209]. Most reports link to Kawasaki disease (for example, [212]) This does not seem unique to SARS-Cov-2 as a novel coronavirus , New Haven coronavirus, was linked to Kawasaki Disease in 2005 [49]. While no obvious literature exists on vitamin K involvement, as discussed later there is reason to suspect it. It is interesting to note that SARS-CoV nsp10 interacts with NADH4L [125] which speculatively may allow it to mutate to interact with VKORC or other proteins with vaguely similar functions. Another mystery seems to be post-infection olfactory dysfunction [219]. Interestingly, the VKDP's Gas 6 and protein S appear to play a regulatory role in the SVZ and in particular olfactory bulb neurogenesis [64].

Common hemotological observations worth noting include elevated d-dimers with mild increase in prothrombin time and mild thrombocytopenia [122] , eosinopenia [126] , lymphopenia commonly with eosinophilia [128] , and leukopenia mostly in younger or asymptomatic patients [235]. Higher neutrophils [60] or neutrophil-to-lymphocyte ratio [130] was found to be predictive of worse outcomes. One recent review found leukocytosis and leukopenia to correlate with severe disease [90].

The causes for these results are unclear although leukocyte apoptosis is a possible contributor to leukopenia. The fluctuating nature of excess and depletion of some cell types may reflect different disease trajectories of simply that the production and consumption rates vary without correlation to each other. Highly activated neutrophils are commonly noted for their damage although explicit reference to NET's and calprotectin was absent until recently [241] and more a request for clinical information on them [152] . One publication did finally find extracellular material including DNA suggestive of NET's [56] . D-dimer may not be too surprising given coagulopathies but interpretation may be open. It has, for example, been considered a biomarker for antihistamine resistant chronic urticaria [12] .

A lot of literature has discussed the affinity of SARS-Cov-2 for ACE2 as a means of cell entry. Lab measurements put the SARS-Cov-2 receptor binding domain affinity for ACE2 in the "low nanomolar" range [115]. Some works have explored increases or decreases of some measures of ACE2 expression. Once translated and trafficked to the plasma membrane it can engage in several processes before meeting one of two fates- internalization or cleavage. Figures such as transcription or translation rate may not be as important as internalization rate which describes how quickly it can turnover and bring in virus. while deployed on the cell surface, ACE2 can hydrolyse AngII to Ang-(1-7) , bind integrin $\beta 1$, associate with neutral amino acid transporter B0AT and be modified with calmodulin to reduce cleavage [34]. These features do not suggest a need for ligand to provoke internalization.

At least one work predicts lower SARS-Cov-2 case counts but higher severity for blacks due to lower ACE2 levels [213] but the dynamics of creation, cleavage, and internalization is complex. Measurable levels of protein may be lower because internalization is faster and the internalization of virus loaded receptors may be the important step. In any case, the current epidemiological data it difficult to sort out due to methodology issues and confounding factors like social policies.

Apparently lacking is any literature linking covid-19 to things like NEDD4. The entry of ACE2 relates to NEDD4 [160] competing with extracellular cleavage to release a soluble form although as late as 2014 details of the competing fates remained obscure [84] .

2.2. Nutritional issues in covid-19 and related conditions

There is not a lot of literature specifically about SARS-Cov-2 nutritional issues just because it has not been around that long but at least one specific motivation for interest in nutrition arises from the recent observation that ileal absorptive cells express ACE2 and appear receptive to infection [238]. One recent report [196] suggested increased arginine and tryptophan metabolism during infection along with "massive suppression of amino acid metabolism." This report also found noteworthy lipid and steroid metabolic changes as well as suppression of platelet factor 4 which has a confusing relationship to protein C [170] [111] .

Nutritional issues for related diseases have been assumed to exist for covid-19 and trendy ideas have been advanced even if historically not effective against these other diseases. For example, at least passing interest was expressed in vitamin C for usage in a clinical trial [29]. Vitamin D status has gotten a lot of attention too, for example [46]. However, these seem to be based on the remaining anti-oxidant hysteria of the last few decades or association studies in the case of vitamin D. Association studies are ambiguous and tend to promote confirmation bias. These can also be interpreted as pointing to other nutrient issues that tend to segregate with vitamin D status. Vitamin D may covary somewhat with vitamin K among others especially when limited by GI issues , for example [192] .

Nutritional immunity, the host's attempt to deprive a pathogen of nutrients, is a well known response to infection. One notable feature common to many infections is the increased activity of IDO to degrade tryptophan. Its overall role remains a topic of research and exceptional status with this virus has not been determined. Tryptophan is observed to be depleted in various conditions such as HIV but supplementation is not always pursued. Instead, IDO inhibition is favored due to the belief that the breakdown products are the bigger problem [153]. It is also known, however, that tryptophan can overcome inhibition of proliferation by IDO products in T-cells and natural killers (see figure 3 for example [58]). Tryptophan depletion per se, regardless of other metabolites is an apoptosis sensitizer [117] and supplementation may prevent apoptosis in the presence of these other metabolites considered deleterious.

Besides tryptophan, phenylalanine metabolism and more directly tyrosine deficiency may be an issue. Dysregulation of tryptophan and phenylalanine metabolism has been studied in HIV [70]. Phenylalanine to tyrosine ratio was observed to increase in HIV-1 patients without treatment suggesting PAH reduction [233]. It has recently been observed that PAH is likely inhibited during several pathological states and these effects are likely to be effects common to many disease states. For example, both have been observed to be depleted during resolution of contact dermatitis [239]. Further interest in tyrosine depletion may be suggested from age related issues with thyroid hormone [19] [105] , which has been observed in covid-19 patients [218], or even graying hair as the latter has been related to innate immune activation and viral infection [75]. PKU comorbidities include a large range of conditions that may not entirely relate to excess phenylalanine [207] but rather low tyrosine.

Arginine may be consumed in the generation of NO and exogenous NO in breathing atmosphere is being explored [143] perhaps more so than arginine status. The interest in nitric oxide against coronavirus dates back at least to 2005 and SARS [7]. Arginine has been considered in some detail in various infections but question of good or bad remains complex [27] . Interestingly, lysine stimulates HIV growth but this may be largely a virus specific issue [28] as in Feline Herpesvirus it was observed that arginine promotes while lysine retards growth [135]. As lysine and arginine may compete for transporters, there is always a concern when these are a bottleneck.

The coagulopathies possibly relate to NET nucleation and a dysregulated coagulation cascade. Calcium and vitamin K are two important components of this system. One reason for pathological clotting and bleeding may be sudden erratic depletion of fully reduced vitamin K. Some evidence suggests an increase in ischemic strokes for AFIB patients when starting warfarin [106] prior to achieving intended anticoagulation consistent with depletion of protein C and S occurring early on. For other conditions, there is finally controversy over supplementation versus antagonism of vitamin K [24]. There is little doubt that vitamin K contributes to regulated coagulation as well as other functions such as maturation of Gas6, an apoptosis preventor [76], even if the clotting may be the most well known. It has a nuclear receptor like other lipid soluble vitamins [69] and may have wide ranging effects along with its role in gamma carboxylation of calcium binding areas. The vitamin K dependent TAM system - TYRO3, AXL, and MERTK relies on Gas6 and protein S to aid resolution of inflammation [175]. Sometimes VKOR antagonists are cited as anti-fibrotic or having other benefits but some such as 3-acetyl-5-methyltetronic acid, like other small molecules, may have many effects although they are worth consideration [208] .

One case report series did show protein S deficiency related to Varicella infection,"... six cases of protein S deficiency secondary to varicella. Five cases were complicated by thrombotic and vascular events, namely purpura fulminans and necrotic vasculitis, deep vein thrombosis and stroke. " [158]

Other interesting features of vitamin K exist that may point to relevance in disease and old age. Vitamin K2 was shown to reduce inflammation and ROS production while improving mitochondria performance in the presence of rotenone [222] suggesting at least possible support for some disease related conditions. Lifetime low-phyloquinone diet was associated with cognitive problems in old but not young rats [30].

Possibly the covid-19 skin conditions relate to acute depletion of vitamin K creating both clotting and clot regulation

problems as has been observed with other diseases.

In the diseased state, use of energy and citric acid skeletons is enhanced and depletion of TCA components is a real concern although it may be programmed as part of nutritional immunity. There is some indication that under hypoxic conditions TCA cycle intermediates are regulated as part of a nutritional immune program [78],

Citrate, the first intermediate of the Krebs cycle, plays an important role in immunity. For instance, citrate is required for fatty acid biosynthesis, which is involved in mounting an appropriate immune response (Moon et al., 2015). Moreover, citrate can be converted via cis-aconitate to itaconate, which possesses antimicrobial and anti-inflammatory potential (Michelucci et al., 2013). Thus, citrate is thought to form a building block for the production of inflammatory mediators and antimicrobial molecules. However, high levels of citrate may also entertain the proliferation of *C. burnetii* (Omsland et al., 2008), which is likely to counteract the positive effects of citrate on microbial defense. Citrate is not only important in immunity but also in cancer, as it provides an important source of carbon (Hatzivassiliou et al., 2005; Mycielska et al., 2018). Of note, several cancers are characterized by HIF1a stabilization, which impedes citrate production through oxidative metabolism. Thus, there are remarkable parallels in the regulation of cellular metabolism in immunity and cancer with respect to citrate limitation. However, tumor cells may escape citrate limitation by the use of the reductive carboxylation pathway (Wise et al., 2011);

Citric acid depletion is thought to contribute to disease as it relates to the arginine paradox, [42],

Therefore, export of intermediates of the cycle is indispensable for anabolic purposes, but also subtracts intermediates that would be recycled to the oxaloacetate, indispensable for condensation with acetate and for maintaining the cycle fully working by forming new citrate. Insufficient oxaloacetate availability into the cycle is followed both by a reduction of energy production and, subsequently, by a reduction of the synthesis of nonessential AA from intermediates of the cycle. This process, however, also triggers the entry of AA into mitochondria to refuel the cycle of oxaloacetate precursors, mostly at the level of succinate or -ketoglutarate. Such phenomena explain the pathophysiological mechanism underlying the sarcopenia/cachexia syndrome occurring in all conditions of peripheral insulin resistance.

Citrate has under lab conditions reduced mobility of calcium and magnesium [162] which may be useful in the presence of leaky membranes.

Calcium status is perhaps not appreciated but just as with citrate in stored blood, regulation with citric acid may be biologically relevant.

3. LIMITATIONS OF CONVENTIONAL ANALYSES

These results differ from others due to being somewhat more speculative but also due to failure of systems analysis based too much on dominant effects. Even though much of the analysis here leads to simply replacing what seems to be missing, it is important to remember thoughts like this [42],

In this review, conditions influencing the low arginine concentrations found in plasma will be reviewed, revising the paradigm that simple replenishment of what is lacking will always produce beneficial consequences. [emphasis added]

Part of the problem is determining what meaningful things are missing after compensatory mechanisms disguise a primary cause. There is plenty of precedent for these issues. For example, one well known confusion occurs with the interaction between folate and B-12 in the elderly [100]. In the current situation, older people generally have a lot that is missing but some of that may be adaptive or based on reference values of little diagnostic value.

Several of the ideas here are probably contrary to conventional wisdom simply due to a more complete exploration of cause and effect relationships in light of failures of conventional wisdom to produce useful results or more recent research modifying interpretation of prior observations. In most of the cases described here, the conventional wisdom suggests limiting some nutrient- for example, tryptophan or vitamin K- in a diseased state due to a few prominent (often the first discovered) effects. Vitamin K is known for its role in clot formation but it also contributes to clot regulation on different time scales. So, you could stop clotting by eliminating vitamin K or, better, regulate it with more vitamin K within some range of other parameters. Tryptophan is destroyed by IDO and the metabolites are often thought to be the problem but these may correct with more tryptophan and eventually depletion per se is a problem anyway. The motivation may make sense- indeed nutritional immunity is based on beneficial nutrient loss - but empirically the overall system dynamics may favor the opposite intervention. With age largely predisposing to nutrient loss, anything that effects old people should elicit a concern about nutritional dependencies.

Once a molecule is categorized as a clotting factor or some other trendy thing like antioxidant it tends to become good or evil and new observations may be rationalized in terms of these first known effects. Or, for lack of a better catch phrase, "chemical X is a Z so it can't do Y" [139]. In the case of vitamin K, clinical success antagonising it makes it difficult to accept evidence that more may be even better under many conditions.

There is also a lot of interest in easy to analyze solutions as, by definition, they are easy to understand - one drug, one target for example. Small molecules are often problematic due to off target effects and more selective antibodies are desirable due to more specific location of action. However, for nutritional interventions the target is often a well regulated metabolic network and trying to find general rules like "more is better" is difficult. Monotonic dose-response curves robust against other factors may not be very common. Further, a single point of failure may require multiple nutrients for remediation before much benefit is observed. It may not be possible to even vary "one thing at a time" to get an unambiguous error signal let alone a cure. Before concluding much based on one effect, it is probably better to actively hunt for compensating mechanisms that create a time or metabolite differential or ratio or something similar.

Food association studies may be another source of difference. These tend to create confirmation bias around something trendy like vitamin C. Much of this analysis was explored after rejecting assumptions about anti-oxidants as the dominant nutrients of value. Earlier a case was made against tocopherols [140] that better agrees with known clinical data. And as is often the case, inconsistent association studies may point to more complicated causal relationships or none at all if the independent variable happens to have a misleading correlation. Another interesting observation is the apparent benefit of citrate against cancer [173] which could explain many food related health benefits often ascribed to vitamin C. . Anti-oxidants as a class have been over rated and ROS signalling is now well known but the myths persist.

Correlations may also occur in shared pathways and a trendy vitamin's associations may mask a more important one with shared statistics. One trendy vitamin, vitamin D for example, could be confused with vitamin K in some observational studies and even intervention where it provides a partial solution. Vitamin D stimulates secretion of vitamin K dependent proteins and is even suggested to work by synergy with vitamin K [89] but presumably does not replace vitamin K for gamma carboxylation.

4. OVERVIEW OF MY "OLD AGE AS STARVATION" WORK

In my theory of aging notes [141], I try to motivate the case for aging being largely the result of GI problems and nutritional issues but not the ones commonly considered - or in essence starvation. The relationship to usual types of starvation is not obvious because the dynamics of age related nutrient deprivation are not typical of classical forms of starvation as compensatory systems can do a good job moving the problem around. Initial results attempting to optimize dog diets have guided the selection of important nutrients.

There is recognition that poor nutrition plays some role in aging but most of the literature assumes it is an effect of conditions such as loss of taste or a stroke [121] rather than a cause. Others concentrate on a trendy, incomplete set of nutrients. Possibly the age related damage is part of a positive feedback loop and malnutrition creates more damage that makes nutrient uptake even more difficult. In some cases, correction of the right nutrients in huge non-competing doses may allow for passive diffusion in place of normal damaged mechanisms. Considering low acid and enzymes as part of a positive feedback loop that continues to decrease the supply of things they absorb, it is worthwhile to consider the known nutrients likely to be effected and determine that supplementation is complete enough to fix identifiable problems. Much literature seems to state as fact that some subset of nutrient is responsible for some effects but the evidence is not clear.

One feature of old age in this scenario is amino acid starvation with muscle wasting being largely due to attempts to scavenge amino acids. The loss is likely not uniform and chronic stimulation of IDO may slowly deplete tryptophan preferentially as seen in diseased state. In essence the body is in a state of chronic low level disease combined with a specific absorption problem. One interesting work found mutation in an amino acid transporter, SLC7A8, that effects age related hearing loss [72]. This may suggest an imbalance of amino acids could occur in old age although replacement of any particular one may not be the solution.

Generally, lower proton(acid), chloride and enzyme levels can be expected in old age and broad areas of concerns are metals, amino acids, things attached to proteins, and fat soluble vitamins. Depending on the pH preferences of the enzymes, acid alone may be a big factor and oral enzyme replacement of limited value. One common reason in old age may be "atrophic gastritis" which would not be obviously related to an acute disease. Adding acid to meals does apparently help folate and B-12 absorption [177]. This source then goes on to consider absorption of vitamin D decrease with age as only an effect and not so much a cause. It also singles out iron with no comments on other cations. "Things attached to proteins" could include biotin or the better known issue with B-12 Literature on amino acid supplements for humans has had mixed results but this is probably attributable to inability to balance the amino

acid pool, lack of availability of hydrophobics like tryptophan and valine, and lack of associated components like citric acid for metabolic flexibility.

In cases where its impact on nutrition is considered, analysis may just be confined to trendy vitamins. One recent article suggested vitamin C was needed for serotonin and norepinephrine production as well as collagen [32] but neglected stoichiometric amino acid requirements and copper for crosslinking. Correcting the suspects discussed here did seem to help several older dogs although several other additions such as B vitamins were included too (unpublished observations). In the case of covid-19, only those that seemed to help the dogs and appear relevant to the SARS-Cov-2 are included.

Low stomach acid and enzymes would be expected to deplete amino acids and things attached to proteins as well as cations with lower solubility. In some documented cases, significant GI disruptions including bacterial overgrowth can impair all fat soluble vitamin levels but leave vitamin K as the most important [163]. Also note that nutrient absorption is not the same as fat absorption. Initial results feeding these nutrients to old dogs have been encouraging [141]. Recent work suggesting utility of vitamin K for age related conditions [199] appears to be the less common view. The symptoms of old age empirically suggest vitamin K deficiency is likely a big contributor but there seems to be a general concern about giving vitamin K to old animals or people. In particular, there has been no indication of severe problems with vitamin K in older dogs. Recently up to 10mg/day K1 and 15mg / day K2 have produced no obvious problems and may be contributing to recovery in 5-30kg dogs. With a background of amino acids, copper, citric acid but little zinc even in the presence of heartworm vitamin K appeared safe in one dog although a therapeutic role could not be determined and coughing symptoms did worsen coincident with the addition of aromatic amino acids [138]. While some age related conditions like osteoporosis may appear to be issues with cations like calcium, those may in fact be more related to amino acids, vitamin K, citric acid, and nutrient signalling. That is, to form the proper calcium minerals, collagen scaffold needs to exist using the required amino acids and conditions need to encourage appetite nucleation and growth. Added calcium may help density but may not create normal strength bone and rather precipitate calcium all over.

Tryptophan depletion seems to correlate with aging [210] [211] although it is not clear if tryptophan depletion per se is an issue and generally there is a focus on disabling TDO and IDO. Metabolic changes may occur with many nutrients and sorting out cause and effect may be difficult. In particular, dog breed longevity appears to correlate with tryptophan metabolism [87]. In extreme depletion, protein synthesis will suffer and uncharged tRNA can accumulate [232]. Low fidelity translation can occur due to lack of one amino acid or the mischarging due to amino acid analogues competing with the limited supply of accurate ones. This can become toxic in many ways but one possible problem is replacement of a tryptophan with something smaller in a transmembrane region one result of this is ion leak which may cause many age related diseases. Amino acid supplements for old people demonstrate some reports of success [157] but results do not appear to be consistent as the varying details may prevent consistent success [224] [93]. This may be due partially to getting a complete mix of components that solve existing problems without creating more of them. This work is not based on any of these but rather an analysis of what is likely to be missing. Originally it would have been easy to dismiss tyrosine as non-essential except for some quick literature search and feeding experience with dogs suggests it as an important contributor to stressful conditions like aging or disease.

Copper has gained attention lately as being important to cardiovascular disease, specifically ischaemic heart disease [41] which may be a concern with exaggerated coagulopathies. It has been well known to facilitate killing by macrophages [55]. While the theory suggests many cations like amino acids will be lacking, copper is likely the most important to address. Pharmacological supplements of zinc and copper fed to newly weaned piglets reversibly modified microbiota and improved intestinal health [156] which may be important in older or diseased people.

Many of these issues would benefit from acid consumed with meals for which citric acid may be a good choice. The citrate has other benefits that may be important for calcium control and metabolic flexibility.

5. DISCUSSION - APPROACHING SARS-COV-2 AS A NUTRITIONAL DISEASE

If the age dependence of SARS-Cov-2 severity can be explained in terms of nutrients involved in my aging theory, then it should be possible to reduce both disease severity and typical age related diseases by modifying the status of these nutrients. The observation of a SARS-Cov-2 related syndrome in children with prolonged or second disease course may be rationalized as nutrient depletion similar to that theorized to occur in old people. Young children need protein synthesis to grow more than middle aged although they both presumably have well functioning GI tracts. If the disease can damage GI tract and deplete nutrients eventually similar nutritional situations may arise.

The above considerations lead to suggestions that depart from most published notions about related diseases and aging. The controversial issues largely involve benefits from tryptophan and vitamin K while adding possible benefits from tyrosine, copper and citric acid while agreeing with some ideas related to ambiguity about arginine or nitric

oxide. There is no obvious involvement of antioxidants or other trendy nutrients for most people eating a Western diet although as noted , especially in old age, broad systems could be damaged requiring a range of nutrients until function is restored. The ones mentioned here would likely be the most important and able to show benefit on their own in many cases.

While a specific protocol is not yet defined, adding these nutrients incrementally and observing initial results may be useful for formulating a specific therapy. Any approach should be designed initially to create data not hypothesis test. Hopefully the details are not that critical however some patients may have idiosyncratic issues such as protein S deficiencies and evaluation of the more controversial items may benefit from better monitoring.

5.1. ACE2 and age related conditions

ACE2 received a lot of attention related to SARS-Cov-2 and is known to be part of the blood pressure issues common in older people. However, apparently lacking is any literature linking covid-19 to things like NEDD4. The entry of ACE2 relates to NEDD4 [160] competing with extracellular cleavage to release a soluble form although as late as 2014 details of the competing fates remained obscure [84] . Reiterating the earlier point, NEDD4 is an attractive speculative internalization rate controller as it senses a common attribute of many predisposing conditions. The old age theory suggests ion leakage due to defective membrane spanning proteins among other reasons and this would presumably increase NEDD4 activity.

Expression of ACE2 was recently shown to be interferon stimulated in humans [238] leading to the suggestion that infection should increase the net amount of ACE2 on the surface. NEDD4 is downregulated by ISG15 in response to interferon stimulated by many virus infections such as " influenza, herpes, Sindbis, and HIV " and apparently can reduce Ebola VLP release [136]. Both of these may allow for more surface expression and cleavage. NEDD4 has many recognized functions [91] but as a cytosolic sodium sensor, culling sodium channels to maintain intracellular sodium levels, it could be expected to be more active in cases of sodium leak or intracellular excess. NEDD4 also responds to calcium [215] which may be a bigger issue than sodium as maintaining submicromolar concentrations could be more difficult. NEDD4 has been shown to use, " 1)ion channels and transporters, 2)membrane receptors, 3)tumor suppressors, and 4) endocytic regulation proteins" as substrates [215]. Anything which causes upregulation may result in culling of many things suggesting intracellular starvation would be possible under the same conditions that enhance viral entry.

Interestingly, ADAM17, the protease creating the soluble cleavage product, appears to be stimulated by thrombin [132] which may reduce infectivity suggesting a benefit for the virus coagulation inhibition.

Artificially introduced mRNA for ACE2 was found to benefit from codon optimization which at least partially works by switching to a more abundant synonym [187]. This does not address amino acid starvation explicitly but rather the abundance of charged tRNA which directly is impacted by amino acid starvation.

5.2. Citric acid

Citric acid is expected to have benefits by controlling calcium and replenishing TCA skeletons in the presence of increased metabolic loads. Some evidence exists for immunomodulation and related effects [5]. It may also be metabolized to bicarbonate . The transfusion literature is full of interesting observations (for example [88]) It may also be instructive to look at earlier observations for its "obscure " role in pH regulation noted since at least 1933 [114] [188] .

However, it seems to be ignored in food association studies that look for vitamin C or flavonoid effects. For example, in one study most of the beneficial foods [96] have high amounts of citric acid or other organic acids . This is despite it being the central skeleton in the citric acid cycle which can be depleted during heavy metabolic demands. It may turn out that many benefits attributed to vitamin C are really due to citric acid and prove to be another food association misinterpretation. There is some indication that 3grams/day can reduce fatigue [203]

Calcium control can influence both signalling and precipitation. Accumulated precipitates may be common in old age and virally induced whereas inappropriate signalling may occur due to leaky membranes or in various calcium dependent processes such as coagulation. As a metal chelator, it may help distribute calcium and other metals. Dietary organic acids including citric acid have been explored for thrombus management with some indication of effect [99].

The TCA turnover may increase in any "stressful" situation such as SARS-Cov-2 infection or common conditions of old age. Fever in children has been associated with increased heat production of 11.3% per C and increased nitrogen excretion [18]. Protein-calorie malnutrition is common in the critically ill at least in part due to increased resting energy expenditure [189]. It has been generally known that infection increases demands for amino acids [165] as part

of a larger metabolic turnover. These may all seem unrelated but the metabolism of unbalanced amino acids often proceeds through the TCA with significant effect on energy flow [17].

While fever is a well known common result of SARS-Cov-2 infection, it is notable that there is no indication yet of any calcifications related to SARS-Cov-2 [97] [77] although they may take time to grow to discernable size. Intracranial calcifications detectable with CT can be caused by several viruses [178] and appear to be best known for fetal infections. Mineralized avian influenza virus has been recently observed and the minerals thought to contribute to stability [236]. Ion permeability increases including for calcium are well known for HIV to encourage its survival and translation of the viral mRNA [36] but literature on coronavirus was not found.

Citrate may be best known in blood preservation and kidney stone literature but similar principles operate in *in vivo* blood but with different scales.

5.3. Amino acids

Amino acid nutrition may be relevant due to metabolic changes, GI digestive efficiency, increased protein synthesis, or changes in transporter distribution. In old age, the GI problems are thought to be primary but in the disease state the others may be most important. The common themes in disease and aging point mostly to tryptophan, tyrosine, and arginine with maybe some importance for others like valine. As they may compete and shift metabolism, benefits from essential or broad supplementation may not exceed those of a selective approach which can balance the pools in light of excess metabolism or altered transport in the disease state. At least with IBD, broad spectrum supplementation at normal levels is recognized as a possible problem compared to targeted high-dose supplements [65]. One notable issue with extreme imbalance is the amino acid pool available for charging tRNA and the effects on protein synthesis rate and fidelity as well as signalling by uncharged tRNA and charging synthases.

Thinking aloud

mischarging with amino acid analogs would depend on the existence of such analogs which could be of microbial origin or maybe host. These may not compete until the parent amino acid is in short supply.

The latter is an emerging field with roles being eluci-

dated in nutrient sensing and cellular adaptations to starvation [79] [25] [166] Amino acid considerations in the disease state have been explored before, in one case leading to a formulation called CATHON but also pointing to depletion of TCA skeletons [42]. It is likely to be easier to make non-essential amino acids if the TCA is fed with citric acid. In the aged state chronic low level infections may be a problem but the primary GI issues may favor a more comprehensive amino acid mixture while still favoring these beyond normal abundances. Lists of well known functions exist, such as [124], and supplement modifications could be designed in response to outcome. After taking care of these, there may be more issues that become apparent as, besides the B0AT transporter associated with ACE2, amino acid transporters come up in more and more contexts often related to mTOR or immunity. For example, SLC15A4 [108] just came up in the context of endolysosomal transport related to SLE [82] .

After making this list of amino acids, it became apparent that the amino acids most prominently highlighted all are in pathways using tetrahydrobiopterin (BH4) [155] which, along with neopterin, reacts to disease [146] [47]. One work from 1961 that only knew of PAH as a possible pteridine indicated the conversion to tyrosine was slowed with methotrexate [67] and it is interesting to speculate on nutritional immunity playing a role via folate or look for follow up work. One recent survey of cytokine storm markers in SARS-Cov-2 patients lists neopterin as "not assessed" [83]. Interestingly, neopterin levels were found to inversely correlate with survival in elderly hip fracture patients [116] suggestive of related processes occurring in this group. Phenylalanine stimulates the activity of tetrahydrobiopterin [124] and PAH downregulation appears to be common in the diseased state. If Phe stimulation does not overcome other sources of downregulation, this leads to concerns about the essentiality of tyrosine but also Phe toxicity and reasons to be suspicious of supplementation.

Tryptophan is the well known target of IDO and TDO. Earlier works on metabolites demonstrated some activity of the metabolites suggesting IDO inhibitors would be more important than tryptophan supplements but at least one work's data showed tryptophan prevented the bad things. This protection may be quite diverse. For example, yeast exposed to SDS are more resistant with tryptophan [186] and as described below there may be detergent like effects from membrane spanning proteins mistranslated at tryptophan sites.

Being rare to begin and a precursor to consumables , it is easy to speculate that it gets depleted quickly and could get to the point of making high fidelity protein translation difficult leading to non-specific problems and futile cycling of defective proteins. Specifically, with a role in sealing membrane bound proteins, it is a good suspect for possible ion leaks at low fidelity products that make it to the plasma membrane. This would likely be mostly sodium and calcium inbound. Sodium would likely lead to swelling commonly observed in infection and calcium signalling would be chronically stimulated and eventually could lead to precipitation [137] . Further speculation suggests citric acid

could minimize calcium leakage and help maintain sub micromolar cytosolic levels. This leakage is likely a natural component of many old age diseases including high blood pressure.

Any amino acid pool imbalance may stall ribosomes and produce defective proteins for both host and virus. Tryptophan and tyrosine appear to be preferred at membrane interfaces likely due to large rigid planar structure [229].

Certainly with GI issues as may occur with old age and probably with SARS-Cov-2 , improving solubility may help. Wetting of tryptophan powder for example appears to improve with calcium pantothenate powder although reactions and degradation of both may need to be explored. In general, it is good to remember these things are materials not just nutrients [139].

Interestingly, tryptophan to niacin conversion appears to be decreased in women due to hormonal control [197]. Women do tend to live longer and have less severe covid-19 although statistics on various groups of women and men with varied hormone status would be needed to demonstrate an important correlation.

Tyrosine I originally dismissed and there may be more surprises but a recent in silico drug discovery effort for SARS-Cov-2 turned up targets including several carboxylases and homogentisate 1,2 dioxygenase [31] which is part of the tyrosine degradation pathway. It did appear to be beneficial in old dogs and maybe my grey hair(unpublished observation) . Observations of increased Phe/Tyr in HIV patients cited earlier suggest it may be decreased. Depression related to interferon-alpha may be due to tyrosine decrease [54]. Other work suggests biopterin remains intact during ifn-alpha treatment [68]. The need for MAOI's to make even high doses of tryptophan effective against depression [151] further suggests a role for tyrosine in some cases(for an interesting discussion of misconceptions see [85]) .

Tyrosine is not normally considered essential, although see the CATHON paper [42]. , and often nutrition assays quote tyrosine plus phenylalanine [62] as if they are interconvertible . The prior reference also points out that unsuccessful supplements for kidney patients lack tyrosine, even though reduced conversion is observed [109] , and anecdotal evidence supports addition of tyrosine to achieve positive nitrogen balance.

If tyrosine is a virulence limiting molecule, then people with increased demands due to highly pigmented hair or skin may be at additional risk or have metabolic flow already somewhat adjusted.

For SARS-Cov-2 intervention relationship to phenylalanine may need consideration. Phenylalanine or tyrosine dietary restriction was sufficient to induce lymphocytopenia in mice and modulate natural killer activity [147]. One reason for the observed low cell counts with SARS-Cov-2 could be reasonable nutrient depletion. However, phenylalanine may spontaneously be hydroxylated to an m- or o- tyrosine isomer and the latter two may produce contaminant resistance among tumors due to tRNA mischarging [176]. Interestingly, phenylalanine and histidine appear to minimize this resistance although direct relevance to SARS-Cov-2 is unclear. Tyrosine mistranslation as phenylalanine or histidine is often observed in CHO antibody production and it can be corrected with supplemental tyrosine [52] although this does not prove it is ever significant in mammals it does establish plausibility. In any case, selective supplementation avoiding phenylalanine and histidine may be worth exploration.

Thinking aloud

An interesting example of wide ranging effects of amino acid depletion is the case of poisoning by the common food contaminant Ochratoxin A. A lethal dose for mice can be overcome with a similar weight of phenylalanine [37]. The mischarging of Phe-tRNA creates kidney and liver problems that may be difficult to identify in the case of lower levels of multiple problems.

There are various claims about arginine anti-aging effects [59] Arginine is somewhat controversial as it has shown bad results in some clinical trials [223] and is not considered essential, maybe semi-essential , for healthy humans but its interactions with copper and lysine may need to be considered.

At least one study with aging rats suggested that age related declines in NO were not likely to be due to arginine synthesis declines [149] although age related declines in absorption may still be an issue as endogenous production may not be sufficient with increased demands. A similar ambiguity may exist with tyrosine which can be produced but also supplied through a normally functioning GI tract.

One interesting work demonstrated teratogenicity due to low copper and nitric oxide [228] demonstrating its signalling role and interaction with copper. This is in addition to NO and copper as they relate to vasodilation [66].

Another study found good reason to look for food association with improved health in the elderly but found none [164]. Food association studies can be confounded and in the case of the elderly multiple nutrient deficiencies may need to be corrected to see any clinical improvements.

Nitric oxide was identified previously as suppressing an earlier SARS in vitro through unexpected mechanisms of decreased RNA production and spike protein post translational modification reduction [8]. Clinical trials involving NO inhalation are being designed for these reasons [119].

Arginine could be depleted due to beneficial NO production. However, there have been some concerns raised with the dogs a few suspicious results even with lysine supplementation. One trial with many people over 60 and ST-segment elevation MI demonstrated excess deaths in the arginine arm leading to trial stop [190]. Arginine has been observed to improve cellular immune response [92].

Valine is possible good as some publications have found effects on virus behavior. There is some indication it can improve DC maturation in normal and HCV patients [101] and supplemented with interferon appeared to benefit and HCV infection [103]. Interestingly, turnip yellow mosaic virus apparently contains a sequence that can esterify valine or tRNA like behavior [231] although general effects of virus on amino acid or tRNA's remain to be explored.

5.4. Trace nutrients normally associated with protein

In the presence of a protein digestion issue it is likely trace nutrients associated with proteins will be deficient too. Biotin for example and all SMVT substrates in rotation. However, there is no indication the disease state increases consumption of these significantly.

5.5. Metals, atomic cations

The aging theory suggests many trace metals will be in short supply especially with popular emphasis on calcium and iron crowding the others out. Among the possible victims are copper, zinc, magnesium, and manganese. The symptoms of old age and SARS-Cov-2 disease suggest copper is a dominant problem in both.

Thinking aloud

In some informal feeding experience with old dogs, the copper and zinc seemed most important although magnesium and manganese were included. Copper without zinc eventually created some feeding hesitancy which resolved with zinc. Incidents of diarrhea seemed to decrease but otherwise not sure. Other work suggests copper would probably be the most relevant forgotten nutrient related to SARS-Cov-2.

Early results with mice demonstrated an indication that copper deficiency could prolong clotting times and, at least in female mice, promote thrombotic lesions [133]. By 2000, it was a recognized problem with heart and vascular disease from a Western diet [107]. Some of the features of old age would likely manifest slowly such as the lack of crosslinking. In the acute disease setting, different effect may matter.

One issue with SARS-Cov-2 treatment is QTc related to hydroxychloroquine. While not discussed much in modern literature, earlier arthritis work uses this along with ACE inhibitor captopril as examples of possibly effective treatments due to copper manipulation [22]. As it is also used to treat prophyrria cutanea tarda, an iron related disease [200], and has retinal side effects similar to deferoxamine [40]. Chloroquine is often described as a zinc ionophore with little relationship to iron or copper [225].

One study found measurements of QT interval such as QTc tended to get worse with low copper rather than low zinc [94]. While copper can shift voltage dependence of potassium channels, a concern for QTc elongation, deficiency was seen in 1999 to have an effect on calcium tolerance that is not correctable with calcium channel blockers [171]. Literature on cardiac involvement seems to be lacking but a recent result summarizing metal ion modulation of ion channels demonstrated an important interaction between copper and a calcium channel in the retina [134] which is likely to represent a common interaction.

One other work [92] suggests including copper in a diet directed at SARS-Cov-2.

Like with the other nutrients, the sequestration of copper could leave the host cells with an effective deficiency. Clotting and fibrinolysis both appear to be modulated by copper but in non-obvious ways presumably to allow robust responses in most situations. Some of the known regulatory pathways suggest it would be impossible to say in isolation if more or less copper would be better or worse but generally the disease and old age symptoms make more sense as insufficient copper. One interesting case is the effect of copper on PAI-1 which can either increase or decrease stability depending on the presence of vitronectin [26].

Copper is generally thought to be a target for nutritional immunity along with iron and zinc but the extent of this may not be well determined [20]. Calprotectin, a calcium modulated zinc binder composed of S100 [221] alarmin members, was just found in 2018 to bind copper with subpicomolar affinity [21] (also nickel [131]). COMMD1/Murr1 is the means of biliary copper loss and also involved in ion homeostasis elsewhere and a known HIV-1 restriction factor [205]. Interestingly, it also works with NEDD4-2 to remove epithelial sodium channels [104] in an activity which

presumably increases in the presence of sodium leak which this theory predicts increases due to misincorporation at tryptophan codons. For a recent review of ion channel regulation see for example [161] .

Interestingly, copper apparently can inhibit activated protein C's anticoagulant activity but not with human serum albumin present [16] .

Copper is generally added to the phagosome under infection conditions [73] . Indeed, copper has been observed to reduce susceptibility to coxsackievirus although the mechanisms remain unknown [86]. In the presence of possible ADE, any additional neutralization may help if the virus can be exposed to copper prior to emerging.

For some reason, zinc has attracted a lot of popular attention as an antiviral and specifically is thought to work well with hydroxychloroquine. The latter likely is a significant broad spectrum chelator as well as an ERGIC inhibitor. Even in this combination, copper may at minimum reduce side effects.

Some literature suggests that copper is more likely to cause inflammation or other pathological immune responses [184] ZZ .

Recent literature and experience have pointed to a larger role for copper that would likely be especially true with zinc supplements. Copper has recently been appreciated for its role in killing or modulating the virulence of bacteria. Many seem surprised it is an essential nutrient even if they know about zinc.

In general the metals compete with each other. In terms of reaction to the virus, while many people may benefit from most metals, nutritional immunity may be favored with excess copper and zinc depletion.

The coronavirus envelope protein, a viroporin, was compared to the influenza M2 [185] which can be inhibited with copper. It is not clear if SARS-Cov-2 has any proteins significantly effected by excess copper although that may compete with zinc.

While not strictly a nutrient, another metal , bismuth could also be suggested for inhibition of viral helicase.

Bismuth is also considered beneficial here for virus specific reasons although its reputed broader antibacterial roles may benefit the GI tract.

5.6. Fat solubles : A,D,E,K

The pattern of cutaneous symptoms and general blood parameters (elevated d-dimers, mild increase in prothrombin time, mild thrombocytopenia[122]) is not inconsistent with depletion of proteins S and C and coagulant factors as would be expected from a fast loss of reduced vitamin K. The covid-19 skin features may relate to acute depletion of vitamin K creating both clotting and clot regulation problems as has been observed with other diseases. Elucidated roles for protein C, protein S, and Gas6 [76] all suggest a useful role against the pathological immune response similar to that observed in covid-19 patients. Notably Gas6 tends to suppress apoptosis which may be a cotributor to disease progression. Very recently, reduced functional vitamin K status inferred from increased uncarboxylated matrix Gla protein was correlated with elastin degradation and COPD ([168]) was correlated with worse outcome in covid-19 [43]. Direct measurements are lacking at this point but vitamin K deficiency is often ignored and at least one report described misdiagnosis as DIC [9]. One case report abstract noted that infection associated skin lesions can resemble warfarin induced leading to detection of protein C and S deficiencies in their patient [44] but it does not sound common to suspect this. Another case report of infant suggests differential diagnosis of purpuric skin lesions can include suspects including Kawasaki disease and association with various infections [11]. It is also interesting to note that SARS-CoV nsp10 interacts with NADH4L [125] which speculatively may allow it to mutate to interact with VKORC or other proteins with vaguely similar functions.

Protein S and C defects or deficiencies are known or suspected to play many roles in coagulopathies [127] although involvement of vitamin K is not always addressed. Decreased protein S was observed in children with varicella infection [113]. Increased D-dimer and reduced protein S can occur during pregnancy [110] and in children with HIV [169]. With liver involvement, many result are possible as coagulant and anticoagulant proteins suffer as well as clearance in the case of d-dimers [159]. Note also possibility of hypocitremia under disease conditions leading to lack of calcium control in the clotting system.

Loss of reduced vitamin K forms would benefit the virus in several ways. While host factors would be considered, there is some indication of a virus specific reasons. Vitamin K3 exhibited an IC-50 of about $8\mu M$ against SARS-Cov-2 Mpro [80] encoded from nsp5 [13]. This does appear to motivate a survival benefit from inhibiting availability of vitamin K and perhaps the epoxide form is not suitable for viral inhibition. This may be accomplished if nsp10 has mutated to inhibit VKOR locally. Menadione may not be a good dietary choice but phylloquinone can be converted to it [195]. It is probably worth noting that menadione has also been observed to inhibit IDO to a recognizable degree[112] and is also a carbonic anhydrase inhibitor [33]. Loss of vitamin K may then explain reduced blood CO_2 if carbonic anhydrases are limited by menadione.

Host factors could include anything that modulates protein synthesis or immune defenses such as clotting. Precedents with other viruses are known. Receptors Tyro3, Axl, and Mertk with immunomodulating properties and

interaction with viral pathogens have vitamin K dependent ligands [198].

Among host factors, clotting is the most obvious but there are many other vitamin K modulated systems. One overlooked observation is the role for vitamin K1 in embryogenesis. Observed in chicks in 1997 [183] it appears to be programmed suggesting physiological mechanism for reduced vitamin K activity. This may be due to a change in programming to that may facilitate virus replication. Note that this alone may not be a dominant feature as neonates do not appear at high risk for severe disease although there may be few cases of infection.

A 2019 review on aging issues involving vitamin K list several modes of action including on androgen receptors with a pro-cancer effect [14]. The sex distribution in SARS-Cov-2 patients is skewed towards males and this may be a negative factor. i

A few years ago it was noted that warfarin appeared to decrease survival in idiopathic pulmonary fibrosis patients not needing anticoagulation [220] highlighting the possibility of vitamin K contribution to healing problems associated with SARS-Cov-2 infection.

Fully reduced vitamin K is needed to gamma carboxylate and form calcium binding sites. There are two obvious ways to create an effective depletion of vitamin K dependent proteins. One is to make the vitamin K unavailable either by sequestering, or eliminating. Another is to prevent it from being reduced by VKORC. Vitamin E apparently effects supply as α tocopherol selectively depletes extra-hepatic phyloquinone (but not menaquinone) [74]. Interestingly, quinone impurities in vitamin E can inhibit vitamin K carboxylation in very small quantities [45]. The above source also points out that reductions in ischemic stroke by vitamin E seem to be "balanced" by hemorrhagic stroke suggesting something like copper or citrate may be a more accurate approach for clotting problems.

If cycling is being inhibited, there is some suggestion vitamin C could help [95] but there is only inconsistent indication it interferes with warfarin some reports suggesting no clinical relevance [53] while the prescribing information does list high-dose vitamin C as a interaction concern ² as would be expected if it had clinical benefit on VKOR cycling. At least one case report exists in which a patient on warfarin and vitamin C experienced various symptoms due to clotting which resolved with vitamin C withdrawal [181]. Possibly the vitamin C effects the clotting part more than the other functions as both apparently can be selectively targeted somewhat.

It is worth discussing the less suspicious fat soluble vitamins because attention to them may be due to confusion with vitamin K as some evidence does not discriminate well. Currently, vitamin D gets most of the popular press attention due to relationship to calcium and prior to that vitamin E because it could be called an "anti-oxidant." However, vitamin K is the most important based on this analysis .There is some indication that vitamins D and K may associate somewhat whereas E and A can antagonize K. In any case, the interactions may confuse discovery of the primary causes of disease and current interest in D may be more appropriately directed at K. For one recent review on vitamin K, see [194].

Vitamin D is known to effect NEDD4 in two ways. In an indirect way, the increase of serum calcium could increase cytosolic calcium due to leaks and activation of NEDD4. More directly however, the active form of vitamin D (minority in the blood) $1,25(OH)_2D_3$, at least at high doses

Thinking outloud

(blot looks like control smaller than low dose)

was able to reduce NEDD4 expression at least in VSMC's [118] . It is worth noting in passing that [118] also described a good mechanism for general infection stimulated mineralization which I think leads to other age related diseases of putative infectious origin including Alzheimer's.

As a known inhibitor of vitamin K, vitamin E and certainly the more confusing mixed tocopherols should be avoided. There is at least passing concern about them in this context [71] aside from deleterious clinical trial results. I outlined a larger issue with overuse of antioxidants and in particular tocopherols earlier [142].

Vitamin A has not been considered.

5.7. Integration

If SARS-Cov-2 provokes similar nutritional problems as related viruses and if my aging theories have merit, the combination of amino acids, copper, citric acid, and vitamin K may be helpful in both conditions. Some of these work together but most could provide some testable benefits individually. There may be concern about vitamin K producing pathological clotting among those with protein S or C problems, elevated calcium, or those without sufficient copper or citric acid. Any vitamin's benefit can be limited by others and idiosyncratic deficiencies may need additional consideration. More widespread deficiencies may be more likely in the chronic setting that includes old age or a prolonged infection but not as much for an acute disease.

² https://www.accessdata.fda.gov/drugsatfda_docs/label/2010/009218s1081b1.pdf

The most important amino acids appear to be tryptophan, tyrosine and maybe arginine or valine (not discussed in detail). It is not clear if other amino acids would help although some such as phenylalanine and histidine may be deleterious. One realistic concern is not just an absolute deficiency but not enough of one nutrient to allow others to be used properly. Balancing the amino acid pool for fast high fidelity translation and appropriate signalling from the tRNA assembly pathway is one concern. Adaptations can occur too as the interferon gamma of the disease state may divert more tryptophan to protein synthesis, at least in cancer cells, [6] but there is a lot less of it. These amino acids were highlighted due to them being consumed preferentially by disease processes.

Among the metals, copper is likely the most important to consider and zinc may be important but overemphasized to the detriment of copper repletion. It often appears to be the metal that limits use of other nutrients at least under lab conditions. For example, issues with fructose and NAFLD [202] which also may be relevant as obesity, or more inclusively fat to muscle ratio, appears to be a covid-19 risk factor [182]. I tend to rotate metals for the dogs, working in magnesium and manganese. For a while, the vitamin K was given with a zinc deficient diet which may have had some impact. Bismuth may have some utility against SARS-Cov-2 but as part of a virus directed therapy [227] and not an age related nutrient issue. Overall metal balance may be effected by dietary acid or sulfur sources. Taurine or garlic was commonly included in the dog diets with some early indication of a benefit from garlic.

Vitamin K and citric acid may also be useful despite the concern about coagulopathy and the almost placebo-like attitude towards citric acid. The disease severity correlation with old age and vitamin K status suggests that vitamin K levels may be enough to overcome any VKOR inhibition if it exists at all. The citric acid should provide metabolic flexibility in the presence of imbalanced amino acid levels and better control over calcium and metals.

A complete formulation must recognize interactions with each other and patient as well as the materials properties and chemistries of these components and their environments. Lecithin and pantothenate may act as emulsifiers for the more hydrophobic components although reactions and degradation need to be considered. Both valine and tryptophan appear to wet and suspend better with pantothenate while the lecithin does not appear as effective(unpublished observations). It is not clear if any of the emulsifiers aid absorption although these details may explain inconsistent or confusing results in past literature. It is never really possible to test "one thing" at a time as the drug being tested is effected by every other detail.

The limited experience with self and dogs provided some specific results but have not yet been organized. Diets were more complex than the concepts presented here but the isolated nutrients may capture much of the virus specific requirements. The vitamin K used was typically K2 as mixture of MK-4 and MK-7 in 15mg capsules[4]. Addition of bulk K1 powder in dextrin[1] appeared to have some additional benefits but this change was not made in isolation. Copper was usually obtained as citrate [3] or glycinate[2]. Any food quality citric acid powder is likely equivalent and various diet or energy drinks may contain substantial amounts.

Therapeutic or counterproductive effects may be quick to assess. If the cutaneous issues with SARS-Cov-2 are due to coagulopathies, additional vitamin K or even copper or citric acid may create visible changes within days although disease trajectory likely varies on similar time scales requiring better control and statistical analysis. Citric acid's impact on metabolism could occur within hours, indeed impact on calcium could be within minutes, and any fatigue reduction could be on similar time scales. All clinical signs would likely be subtle. If it helps dissolve clots, clean up metals, and replenish TCA skeletons effects on various times scales may occur.

The contributions of amino acids may be more difficult to assess. Tryptophan depletion in the case of aging will likely create leaky membranes leading to sodium and calcium influx activating NEDD4 and promoting low level swelling. In the acute case, recently translated proteins including antibodies will suffer and signalling related to uncharged tRNA will occur . Apoptosis may be a problem especially in effector cells. Response to repletion may be notable in days. Tyrosine may have some similar issues if PAH shuts down. Both are precursors to neurotransmitters and thought to produce mood disorders when deficient [123] although controversy exists [167] . As an alternative to tryptophan, possibly niacin supplementation would reduce the loss of tryptophan (discussed in terms of NAD etc in [81], observed in HIV patients [154]) as it can be a significant sink (and also note the ability of another essential amino acid, leucine, to make this worse) [15]. Pellagra response to treatment typically occurs within days or weeks and presumably similar less obvious issues would correct quickly with tryptophan. In the US, where B vitamins are routinely supplemented, it is possible to have tryptophan deficiency without niacin deficiency. Copper effects due to enzyme mis-metalization or mitochondrial deficiency could be vague and of unknown delay.

Table I shows some dosing ideas for nominally healthy adult humans based on literature values, theory, and scaled amounts from dog feeding.

Material	RDA	Human NOAEL	Amount/Dose	Dosing	Comments
L-Tryptophan	250-425mg/day [174]	4.5g/d[57]	200mg	2-3/day	powder floats, mix with oil or emulsifier
L-Tyrosine	100mg/kg/day [63]		400mg	2-3/day	may be reactive, avoid with copper, may need more with dark hair or skin
L-Arginine	4-6g/day [201]	30g/day [38]	400mg	2-3/day	for PAH deficiency
K1			5-10mg	3/day	may also need lysine hours apart
K2, MK-4 + MK-7			15mg	daily	1% powder, avoid light, include emulsifier
Vitamin K	0.065-.12mg/day [180]	10mg/day [206]	5-10mg[148]	daily	15mg capsules, avoid light , include emulsifier
Copper	.9-1.2mg/day [35]				all sources of vitamin K
Citric acid		1200mg/kg/day [48]	3g/day ^a	multiple	citrate or glycinate, may also require zinc or have other issues with chronic higher doses
					available in bulk as a spice online, in diet drinks and juice. May demineralize teeth at low pH, consider swallowed capsules.

^a this is total for the day as opposed to other quantities

TABLE I: Baseline human dosing estimates based on limited personal experience with self and dogs scaled roughly by weight. NOAEL's are literature values for healthy adults from the indicated source. RDA's change with time and vary among sources. Tryptophan powder will not wet easily, mix with emulsifier pantothenate although there is reaction potential. For home use, note that a quarter teaspoon is about 1.25 cc and ingredient densities may vary between about .2-2 gram/cc or so. Calibration of volume measures with an accurate milligram scale may be required for bulk products.

6. CONCLUSIONS

The lack of an obvious or proven safe and effective approach to the covid-19 suggests that some exploratory work in less accepted areas may be informative if not useful. Currently published analyses have ignored things like protein C and neopterin as well as larger issues with nutrient status creating much uncertainty about the disease. The age distribution suggests that the virus is aided by attributes that segregate with age. My current work points to specific nutritional features of old age that make good candidates for testable and informative interventions. Most of these could be easy to explore safely even in an amateur informal setting. Some may be considered reasonably safe but others like vitamin K may be more controversial. Any information obtained on these in the disease setting may be informative about the SARS-Cov-2 and aging vulnerabilities too. It is not just the lack of nutrient intake that may need to be corrected but also the distribution depending on ultimate bottlenecks in defective or competing transporters or metabolic pathways. Details matter and just mixing together nutrients is no assurance that any benefits will be realized. Physical properties and interactions need to be considered and some nutrients may even need to be limited to see benefits from those of interest. If these approaches work against the SARS-Cov-2 with more careful control , they may be modified to also mitigate age related diseases.

7. SUPPLEMENTAL INFORMATION

7.1. Computer code

8. BIBLIOGRAPHY

-
- [1] 1 percent vitamin k1 in dextrin. Bulk Supplements Website. URL: <https://www.bulksupplements.com/products/vitamin-k1-1-powder?variant=29386494115951>.
- [2] Copper chelate 5mg. Carlson Labs website. URL: <https://carlsonlabs.com/chelated-copper/>.

- [3] Copper citrate 2mg. Pure Encapsulations website. URL: <http://www.pureencapsulations.com/copper-citrate.html>.
- [4] Vitamin k2 mk-4 15mg plus mk-7 60mcg. Relentless Improvement website. URL: <https://supplements.relentlessimprovement.com/vitamin-k2-mk-4-15mg-plus-mk-7-60mcg-p395.aspx>.
- [5] OM Abdel-Salam, ER Youness, NA Mohammed, SMY Morsy, EA Omara, and AA Sleem. Citric acid effects on brain and liver oxidative stress in lipopolysaccharide-treated mice. *Journal of Medicinal Food*, 17(5):588–98, 2014. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4026104/>.
- [6] Isabell Adam, Dyah L. Dewi, Joram Mooiweer, Ahmed Sadik, Soumya R. Mohapatra, Bianca Berdel, Melanie Keil, Jana K. Sonner, Kathrin Thedieck, Adam J. Rose, Michael Platten, Ines Heiland, Saskia Trump, and Christiane A. Opitz. Upregulation of tryptophanyl-trna synthetase adapts human cancer cells to nutritional stress caused by tryptophan degradation. *Oncot Immunology*, 7(12):e1486353, 2018. PMID: 30524887. URL: <https://doi.org/10.1080/2162402X.2018.1486353>, [arXiv:https://doi.org/10.1080/2162402X.2018.1486353](https://arxiv.org/abs/https://doi.org/10.1080/2162402X.2018.1486353), [doi:10.1080/2162402X.2018.1486353](https://doi.org/10.1080/2162402X.2018.1486353).
- [7] Sara Åkerström, Mehrdad Mousavi-Jazi, Jonas Klingström, Mikael Leijon, Åke Lundkvist, and Ali Mirazimi. Nitric oxide inhibits the replication cycle of severe acute respiratory syndrome coronavirus. *Journal of Virology*, 79(3):1966–1969, 2005. URL: <https://jvi.asm.org/content/79/3/1966>, [arXiv:https://jvi.asm.org/content/79/3/1966.full.pdf](https://arxiv.org/abs/https://jvi.asm.org/content/79/3/1966.full.pdf), [doi:10.1128/JVI.79.3.1966-1969.2005](https://doi.org/10.1128/JVI.79.3.1966-1969.2005).
- [8] Sara Åkerström, Vithiagarun Gunalan, Choong Tat Keng, Yee-Joo Tan, and Ali Mirazimi. Dual effect of nitric oxide on SARS-CoV replication: Viral RNA production and palmitoylation of the s protein are affected. *Virology*, 395(1):1–9, dec 2009. URL: <https://doi.org/10.1016/j.virol.2009.09.007>, [doi:10.1016/j.virol.2009.09.007](https://doi.org/10.1016/j.virol.2009.09.007).
- [9] Jack B. Alperin. Coagulopathy caused by vitamin k deficiency in critically III, hospitalized patients. *JAMA: The Journal of the American Medical Association*, 258(14):1916, oct 1987. URL: <https://doi.org/10.1001/jama.1987.03400140078028>, [doi:10.1001/jama.1987.03400140078028](https://doi.org/10.1001/jama.1987.03400140078028).
- [10] ANDRI ANDREOU, SOFIA TRANTZA, DEMETRIOS FILIPPOU, NIKOLAOS SIPSAS, and SOTIRIOS TSIO-DRAS. Covid-19: The potential role of copper and n-acetylcysteine (nac) in a combination of candidate antiviral treatments against sars-cov-2. *In Vivo*, 34(3 suppl):1567–1588, 2020. URL: http://iv.iiarjournals.org/content/34/3_suppl/1567.abstract, [arXiv:http://iv.iiarjournals.org/content/34/3_suppl/1567.full.pdf+html](https://arxiv.org/abs/http://iv.iiarjournals.org/content/34/3_suppl/1567.full.pdf+html), [doi:10.21873/invivo.11946](https://doi.org/10.21873/invivo.11946).
- [11] Ioanna Argyri, Anastasia Korona, Katerina Mougkou, Olga Vougiouka, Maria Tsolia, and Nikos Spyridis. An Infant With Purpuric Rash and Edema. *Clinical Infectious Diseases*, 61(10):1624–1625, 10 2015. URL: <https://doi.org/10.1093/cid/civ550>, [arXiv:https://academic.oup.com/cid/article-pdf/61/10/1624/855209/civ550.pdf](https://arxiv.org/abs/https://academic.oup.com/cid/article-pdf/61/10/1624/855209/civ550.pdf), [doi:10.1093/cid/civ550](https://doi.org/10.1093/cid/civ550).
- [12] Riccardo Asero. D-dimer: A biomarker for antihistamine-resistant chronic urticaria. *Journal of Allergy and Clinical Immunology*, 132(4):983–986, oct 2013. URL: <https://doi.org/10.1016/j.jaci.2013.04.037>, [doi:10.1016/j.jaci.2013.04.037](https://doi.org/10.1016/j.jaci.2013.04.037).
- [13] I Astuti and Ysrafil. Severe acute respiratory syndrome coronavirus 2 (sars-cov-2): An overview of viral structure and host response. *Diabetes & Metabolic Syndrome*, 14(4):407–12, 2020. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC7165108/>.
- [14] Kotaro Azuma and Satoshi Inoue. Multiple modes of vitamin k actions in aging-related musculoskeletal disorders. *International Journal of Molecular Sciences*, 20(11), 2019. URL: <https://www.mdpi.com/1422-0067/20/11/2844>, [doi:10.3390/ijms20112844](https://doi.org/10.3390/ijms20112844).
- [15] AAB Badawy, SL Lake, and DM Dougherty. Mechanisms of the pellagrogenic effect of leucine: Stimulation of hepatic tryptophan oxidation by administration of branched-chain amino acids to healthy human volunteers and the role of plasma free tryptophan and total kynurenines. *International Journal of Tryptophan Research : IJTR*, 7:23–32, 2014. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4259507/>.
- [16] David Bar-Or, Leonard T. Rael, James V. Winkler, Richard L. Yukl, Gregory W. Thomas, and Richard P. Shimonkevitz. Copper inhibits activated protein c: Protective effect of human albumin and an analogue of its high-affinity copper-binding site, d-DAHK. *Biochemical and Biophysical Research Communications*, 290(5):1388–1392, feb 2002. URL: <https://doi.org/10.1006/bbrc.2002.6363>, [doi:10.1006/bbrc.2002.6363](https://doi.org/10.1006/bbrc.2002.6363).
- [17] David A. Bender. The metabolism of surplus amino acids. *British Journal of Nutrition*, 108(S2):S113S121, 2012. URL: <https://www.cambridge.org/core/journals/british-journal-of-nutrition/article/metabolism-of-surplus-amino-acids/83AA18D847252C78AF8EA42958D50101/core-reader>, [doi:10.1017/S0007114512002292](https://doi.org/10.1017/S0007114512002292).
- [18] M. Benhariz, O. Goulet, J. Salas, V. Colomb, and C. Ricour. Energy cost of fever in children on total parenteral nutrition. *Clinical Nutrition*, 16(5):251–255, oct 1997. URL: [https://doi.org/10.1016/s0261-5614\(97\)80037-4](https://doi.org/10.1016/s0261-5614(97)80037-4), [doi:10.1016/s0261-5614\(97\)80037-4](https://doi.org/10.1016/s0261-5614(97)80037-4).
- [19] IM Bensenor, RD Olmos, and PA Lotufo. Hypothyroidism in the elderly: diagnosis and management. *Clinical Interventions in Aging*, 7:97–111, 2012. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3340110/>.
- [20] AN Besold, EM Culbertson, and VC Culotta. The yin and yang of copper during infection. *of Biological Inorganic Chemistry*, 21(2):137–44, 2016. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC5535265/>.
- [21] Angelique N. Besold, Benjamin A. Gilston, Jana N. Radin, Christian Ramsoomair, Edward M. Culbertson, Cissy X. Li, Brendan P. Cormack, Walter J. Chazin, Thomas E. Kehl-Fie, and Valeria C. Culotta. Role of calprotectin in withholding zinc and copper from candida albicans. *Infection and Immunity*, 86(2), 2018. URL: <https://iai.asm.org/content/86/2/e00779-17>, [arXiv:https://iai.asm.org/content/86/2/e00779-17.full.pdf](https://arxiv.org/abs/https://iai.asm.org/content/86/2/e00779-17.full.pdf), [doi:10.1128/IAI.00779-17](https://doi.org/10.1128/IAI.00779-17).
- [22] H. A. Bird. Heavy metals and arthritis. *Annals of the Rheumatic Diseases*, 42(4):474–475, aug 1983. URL: <https://doi.org/10.1136/ard.42.4.474>, [doi:10.1136/ard.42.4.474](https://doi.org/10.1136/ard.42.4.474).

- [23] M Boukhris, A Hillani, F Moroni, MS Annabi, F Addad, MH Ribeiro, S Mansour, X Zhao, LF Ybarra, A Abbate, LM Vilca, and L Azzalini. Cardiovascular implications of the covid-19 pandemic: a global perspective. *The Canadian Journal of Cardiology*, 20200516. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC7229739/>.
- [24] Bart De Brouwer, Ianthe Piscaer, Jan H. Von Der Thusen, Jan C Grutters, Roger EG. Schutgens, Emiel FM. Wouters, and Rob Janssen. Should vitamin k be supplemented instead of antagonised in patients with idiopathic pulmonary fibrosis? *Expert Review of Respiratory Medicine*, 12(3):169–175, 2018. PMID: 29303380. URL: <https://doi.org/10.1080/17476348.2018.1424544>, [arXiv:https://doi.org/10.1080/17476348.2018.1424544](https://doi.org/10.1080/17476348.2018.1424544), doi:10.1080/17476348.2018.1424544.
- [25] S Brer and A Brer. Amino acid homeostasis and signalling in mammalian cells and organisms. *Biochemical Journal*, 474(12):1935–63, 2017. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC5444488/>.
- [26] Joel C. Bucci, Carlee S. McClintock, Yuzhuo Chu, Gregory L. Ware, Kayla D. McConnell, Joseph P. Emerson, and Cynthia B. Peterson. Resolving distinct molecular origins for copper effects on pai-1. *JBIC Journal of Biological Inorganic Chemistry*, 22(7):1123–1135, Oct 2017. URL: <https://doi.org/10.1007/s00775-017-1489-5>, doi:10.1007/s00775-017-1489-5.
- [27] Kristina S. Burrack and Thomas E. Morrison. The role of myeloid cell activation and arginine metabolism in the pathogenesis of virus-induced diseases. *Frontiers in Immunology*, 5, sep 2014. URL: <https://doi.org/10.3389/fimmu.2014.00428>, doi:10.3389/fimmu.2014.00428.
- [28] Evgeny Vlad Butorov. Influence of l-lysine amino acid on the HIV-1 RNA replication in vitro. *Antiviral Chemistry and Chemotherapy*, 24(1):39–46, feb 2015. URL: <https://doi.org/10.1177/2040206614566582>, doi:10.1177/2040206614566582.
- [29] AC Carr. A new clinical trial to test high-dose vitamin c in patients with covid-19. *Critical Care*, 24, 2020. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC7137406/>.
- [30] Isabelle Carri, Elisabeth Blanger, Jacques Portoukalian, Joseph Rochford, and Guylaine Ferland. Lifelong Low-Phylloquinone Intake Is Associated with Cognitive Impairments in Old Rats. *The Journal of Nutrition*, 141(8):1495–1501, 06 2011. URL: <https://doi.org/10.3945/jn.110.137638>, [arXiv:https://academic.oup.com/jn/article-pdf/141/8/1495/30016388/1495.pdf](https://academic.oup.com/jn/article-pdf/141/8/1495/30016388/1495.pdf), doi:10.3945/jn.110.137638.
- [31] Claudia Cava, Gloria Bertoli, and Isabella Castiglioni. In silico discovery of candidate drugs against covid-19. *Viruses*, 12(4):404, apr 2020. URL: <https://doi.org/10.3390/v12040404>, doi:10.3390/v12040404.
- [32] F Cavalcoli, A Zilli, D Conte, and S Massironi. Micronutrient deficiencies in patients with chronic atrophic autoimmune gastritis: A review. *World Journal of Gastroenterology*, 23(4):563–72, 2017. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC5292330/>.
- [33] Ahmet Cetin, Taha Abdulkadir Coban, Murat Cankaya, Umit Nayki, and Mehmet Kuzucu. Investigation of some vitamin type inhibition on human cord blood carbonic anhydrase i and II. *Journal of Academic Research in Medicine*, 3(2):79–83, sep 2013. URL: <https://doi.org/10.5152/jarem.2013.20>, doi:10.5152/jarem.2013.20.
- [34] NE Clarke, MJ Fisher, KE Porter, DW Lambert, and AJ Turner. Angiotensin converting enzyme (ace) and ace2 bind integrins and ace2 regulates integrin signalling. *PLoS ONE*, 7(4), 2012. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3327712/>.
- [35] James F. Collins and Leslie M. Klevay. Copper. *Advances in Nutrition*, 2(6):520–522, 11 2011. URL: <https://doi.org/10.3945/an.111.001222>, [arXiv:https://academic.oup.com/advances/article-pdf/2/6/520/23735153/520.pdf](https://academic.oup.com/advances/article-pdf/2/6/520/23735153/520.pdf), doi:10.3945/an.111.001222.
- [36] JM Costin. Cytopathic mechanisms of hiv-1. *Virology Journal*, 4:100, 2007. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2174939/>.
- [37] Edmond E. Creppy, Monique Schlegel, Robert Rschenthaler, and Guy Dirheimer. Phenylalanine prevents acute poisoning by ochratoxin-a in mice. *Toxicology Letters*, 6(2):77–80, jul 1980. URL: [https://doi.org/10.1016/0378-4274\(80\)90171-x](https://doi.org/10.1016/0378-4274(80)90171-x), doi:10.1016/0378-4274(80)90171-x.
- [38] Luc Cynober, Dennis M Bier, Motoni Kadowaki, Jr. Morris, Sidney M, Rajavel Elango, and Miro Smriga. Proposals for Upper Limits of Safe Intake for Arginine and Tryptophan in Young Adults and an Upper Limit of Safe Intake for Leucine in the Elderly. *The Journal of Nutrition*, 146(12):2652S–2654S, 11 2016. URL: <https://doi.org/10.3945/jn.115.228478>, [arXiv:https://academic.oup.com/jn/article-pdf/146/12/2652S/30020788/jn228478.pdf](https://academic.oup.com/jn/article-pdf/146/12/2652S/30020788/jn228478.pdf), doi:10.3945/jn.115.228478.
- [39] MR Deshotels, H Xia, S Sriramula, E Lazartigues, and CM Filipeanu. Angiotensin-ii mediates ace2 internalization and degradation through an angiotensin-ii type i receptor-dependent mechanism. *Hypertension (Dallas, Tex. : 1979)*, 64(6):1368–75, 2014. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4231883/>.
- [40] M. Dettoraki and M.M. Moschos. The role of multifocal electroretinography in the assessment of drug-induced retinopathy: A review of the literature. *Ophthalmic Res*, 56(4), 2016. URL: <https://www.karger.com/DOI/10.1159/000446321>.
- [41] James J DiNicolantonio, Dennis Mangan, and James H O’Keefe. Copper deficiency may be a leading cause of ischaemic heart disease. *Open Heart*, 5(2):e000784, oct 2018. URL: <https://doi.org/10.1136/openhrt-2018-000784>, doi:10.1136/openhrt-2018-000784.
- [42] Francesco Saverio DiGuardi. To give or not to give? lessons from the arginine paradox. *Journal of Nutrigenetics and Nutrigenomics*, 4(2):90–98, 2011. URL: <https://doi.org/10.1159/000327777>, doi:10.1159/000327777.
- [43] Anton S.M. Dofferhoff, Ianthe Piscaer, Leon J. Schurgers, Jona Walk, Jody M.W. van den Ouweland, Tilman M. Hackeng, Petra Lux, Cecile Maassen, Esther G.A. Karssemeijer, Emiel F.M. Wouters, and Rob Janssen. Reduced vitamin k status as a potentially modifiable prognostic risk factor in COVID-19. apr 2020. URL: <https://doi.org/10.20944/preprints202004.0457.v1>, doi:10.20944/preprints202004.0457.v1.

- [44] Andrea Dominey. Purpura fulminans and transient protein c and s deficiency. *Archives of Dermatology*, 124(9):1442, sep 1988. URL: <https://doi.org/10.1001/archderm.1988.01670090084024>, doi:10.1001/archderm.1988.01670090084024.
- [45] P Dowd and ZB Zheng. On the mechanism of the anticlotting action of vitamin e quinone. *Proceedings of the National Academy of Sciences of the United States of America*, 92(18):8171–5, 1995. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC41118/>.
- [46] M Ebadi and AJ Montano-Loza. Perspective: improving vitamin d status in the management of covid-19. *European Journal of Clinical Nutrition*, pages 1–4, 20200512. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC7216123/>.
- [47] M Eisenhut. Neopterin in diagnosis and monitoring of infectious diseases. *Journal of Biomarkers*, 2013, 2013. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4437389/>.
- [48] MuawiaI.A.Abdelgadir (Elrrigieg, Mahmood.M. Hassan, Murwan Sabahelkhier, OmerF Idris, and MohamedA Adris. Liver dysfunction of wistar rats influenced by citric acid as food additive. *International Journal of Advanced Research*, 4:2085–2089, 06 2016. URL: https://www.researchgate.net/publication/305422860_LIVER_DYSFUNCTION_OF_WISTAR_RATS_INFLUENCED_BY_CITRIC_ACID_AS_FOOD_ADDITIVE, doi:10.21474/IJAR01/817.
- [49] Frank Esper, Eugene D. Shapiro, Carla Weibel, David Ferguson, Marie L. Landry, and Jeffrey S. Kahn. Association between a Novel Human Coronavirus and Kawasaki Disease. *The Journal of Infectious Diseases*, 191(4):499–502, 02 2005. URL: <https://doi.org/10.1086/428291>, arXiv:<https://academic.oup.com/jid/article-pdf/191/4/499/2761847/191-4-499.pdf>, doi:10.1086/428291.
- [50] Yang Fan, Yue Zhang, Akash Tariq, Xiaolan Jiang, Zeeshan Ahamd, Zhang Zhihao, Muhammad Idrees, Azizullah Azizullah, Muhammad Adnan, and Rainer W. Bussmann. Food as medicine: a possible preventive measure against coronavirus disease (covid-19). *Phytotherapy Research*, n/a(n/a). URL: <https://onlinelibrary.wiley.com/doi/abs/10.1002/ptr.6770>, arXiv:<https://onlinelibrary.wiley.com/doi/pdf/10.1002/ptr.6770>, doi:10.1002/ptr.6770.
- [51] K Farsalinos, A Barbouni, and R Niaura. Systematic review of the prevalence of current smoking among hospitalized covid-19 patients in china: could nicotine be a therapeutic option? *Internal and Emergency Medicine*, pages 1–8, 20200509. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC7210099/>.
- [52] Lauren Feeney, Veronica Carvalhal, Christopher Yu, Betty Chan, David Michels, Jennifer Wang, Amy Shen, Jan Ressler, Brendon Dusel, and Michael Laird. Eliminating tyrosine sequence variants in cho cell lines producing recombinant monoclonal antibodies. *Biotechnology and bioengineering*, 110, 04 2013. URL: https://www.researchgate.net/publication/232738679_Eliminating_tyrosine_sequence_variants_in_CHO_cell_lines_producing_recombinant_monoclonal_antibodies, doi:10.1002/bit.24759.
- [53] Celia L. Feetam, R.H. Leach, and M.J. Meynell. Lack of a clinically important interaction between warfarin and ascorbic acid. *Toxicology and Applied Pharmacology*, 31(3):544 – 547, 1975. URL: <http://www.sciencedirect.com/science/article/pii/0041008X75902781>, doi:[https://doi.org/10.1016/0041-008X\(75\)90278-1](https://doi.org/10.1016/0041-008X(75)90278-1).
- [54] Jennifer C. Felger, Li Li, Paul J. Marvar, Bobbi J. Woolwine, David G. Harrison, Charles L. Raison, and Andrew H. Miller. Tyrosine metabolism during interferon-alpha administration: Association with fatigue and csf dopamine concentrations. *Brain, Behavior, and Immunity*, 31:153 – 160, 2013. Inflammation and Mental Health. URL: <http://www.sciencedirect.com/science/article/pii/S0889159112004722>, doi:<https://doi.org/10.1016/j.bbi.2012.10.010>.
- [55] Richard A. Festa and Dennis J. Thiele. Copper at the front line of the host-pathogen battle. *PLOS Pathogens*, 8(9):1–4, 09 2012. URL: <https://doi.org/10.1371/journal.ppat.1002887>, doi:10.1371/journal.ppat.1002887.
- [56] Sharon E. Fox, Aibek Akmatbekov, Jack L. Harbert, Guang Li, J. Quincy Brown, and Richard S. Vander Heide. Pulmonary and cardiac pathology in covid-19: The first autopsy series from new orleans. apr 2020. URL: <https://doi.org/10.1101/2020.04.06.20050575>, doi:10.1101/2020.04.06.20050575.
- [57] M Friedman. Analysis, nutrition, and health benefits of tryptophan. *International Journal of Tryptophan Research : IJTR*, 11, 2018. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC6158605/>.
- [58] G Frumento, R Rotondo, M Tonetti, G Damonte, U Benatti, and GB Ferrara. Tryptophan-derived catabolites are responsible for inhibition of t and natural killer cell proliferation induced by indoleamine 2,3-dioxygenase. *The Journal of Experimental Medicine*, 196(4):459–68, 2002. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2196046/>.
- [59] Mohamed Zakaria Gad. Anti-aging effects of l-arginine. 2010. URL: <https://www.semanticscholar.org/paper/Anti-aging-effects-of-l-arginine-Gad/625b4dd6bd8089299b9846fd82e6eb28f2561c08>.
- [60] James Galloway, Sam Norton, Richard Barker, Andrew Brookes, Ivana Carey, Benjamin Clarke, Raeesa Jina, Carole Reid, Mark D.. Russell, Ruth Sneep, Leah Sugarman, Sarah Williams, Mark Yates, James Teo, Ajay M. Shah, and Fleur Cantle. A clinical risk score to identify patients with COVID-19 at high risk of critical care admission or death: An observational cohort study. *SSRN Electronic Journal*, 2020. URL: <https://doi.org/10.2139/ssrn.3590486>, doi:10.2139/ssrn.3590486.
- [61] Shikha Garg, Lindsay Kim, Michael Whitaker, Alissa O'Halloran, Charisse Cummings, Rachel Holstein, Mila Prill, Shua J. Chai, Pam D. Kirley, Nisha B. Alden, Breanna Kawasaki, Kimberly Yousey-Hindes, Linda Niccolai, Evan J. Anderson, Kyle P. Openo, Andrew Weigel, Maya L. Monroe, Patricia Ryan, Justin Henderson, Sue Kim, Kathy Como-Sabetti, Ruth Lynfield, Daniel Sosin, Salina Torres, Alison Muse, Nancy M. Bennett, Laurie Billing, Melissa Sutton, Nicole West, William Schaffner, H. Keipp Talbot, Clarissa Aquino, Andrea George, Alicia Budd, Lynnette Brammer, Gayle Langley, Aron J. Hall, and Alicia Fry. Hospitalization rates and characteristics of patients hospitalized with laboratory-confirmed coronavirus disease 2019 — COVID-NET, 14 states, march 1–30, 2020. *MMWR. Morbidity and Mortality Weekly Report*, 69(15):458–464, apr 2020. URL: <https://doi.org/10.15585/mmwr.mm6915e3>, doi:10.15585/mmwr.mm6915e3.
- [62] Giacomo Garibotto, Paolo Tessari, Daniela Verzola, and Laura Dertennois. The metabolic conversion of phenylalanine into tyrosine in the human kidney: Does it have nutritional implications in renal patients? *Journal of Renal Nutrition*,

- 12(1):8–16, jan 2002. URL: <https://doi.org/10.1053/jren.2002.29600>, doi:10.1053/jren.2002.29600.
- [63] Peter J. Garlick. The Nature of Human Hazards Associated with Excessive Intake of Amino Acids. *The Journal of Nutrition*, 134(6):1633S–1639S, 10 2004. URL: <https://doi.org/10.1093/jn/134.6.1633S>, arXiv:<https://academic.oup.com/jn/article-pdf/134/6/1633S/24007039/1633s.pdf>, doi:10.1093/jn/134.6.1633S.
- [64] A Gely-Pernot, V Coronas, T Harnois, L Prestoz, N Mandairon, A Didier, JM Berjeaud, A Monvoisin, N Bourmeyster, Frutos PG De, M Philippe, and O Benzakour. An endogenous vitamin k-dependent mechanism regulates cell proliferation in the brain subventricular stem cell niche. *Stem Cells (Dayton, Ohio)*, 30(4):719–31, 2012. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3601423/>.
- [65] FK Ghishan and PR Kiela. Vitamins and minerals in ibd. *Gastroenterology clinics of North America*, 46(4):797–808, 2017. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC6342481/>.
- [66] Leila Gobejishvili, Jack T. Saari, Ayotunde S.O. Adeagbo, Xiaodong Zhang, and Dale A. Schuschke. Dietary copper deficiency increases inducible nitric oxide synthase-mediated vascular dilation in rat aorta. *The Journal of Trace Elements in Experimental Medicine*, 15(2):85–95, 2002. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1002/jtra.10005>, arXiv:<https://onlinelibrary.wiley.com/doi/pdf/10.1002/jtra.10005>, doi:10.1002/jtra.10005.
- [67] Theodore L. Goodfriend and Seymour Kaufman. PHENYLALANINE METABOLISM AND FOLIC ACID ANTAGONISTS. *Journal of Clinical Investigation*, 40(9):1743–1750, sep 1961. URL: <https://doi.org/10.1172/jci104397>.
- [68] Arthur R. Van Gool, Durk Fekkes, Wim H.J. Kruit, Paul G.H. Mulder, Timo L.M. Ten Hagen, Marjolein Bannink, Michael Maes, and Alexander M.M. Eggermont. Serum amino acids, biopterin and neopterin during long-term immunotherapy with interferon-alpha in high-risk melanoma patients. *Psychiatry Research*, 119(1-2):125–132, jul 2003. URL: <https://doi.org/10.1016/s0165-1781%2803%2900113-6>, doi:10.1016/s0165-1781(03)00113-6.
- [69] Jan Oxholm Gordeladze. Introductory chapter: Vitamin k2. In *Vitamin K2 - Vital for Health and Wellbeing*. InTech, mar 2017. URL: <https://doi.org/10.5772/66384>, doi:10.5772/66384.
- [70] Johanna M. Gostner, Kathrin Becker, Katharina Kurz, and Dietmar Fuchs. Disturbed amino acid metabolism in HIV: Association with neuropsychiatric symptoms. *Frontiers in Psychiatry*, 6, jul 2015. URL: <https://doi.org/10.3389/fpsy.2015.00097>, doi:10.3389/fpsy.2015.00097.
- [71] Judith M. Graat, Evert G. Schouten, and Frans J. Kok. Effect of daily vitamin e and multivitamin-mineral supplementation on acute respiratory tract infections in elderly persons. *JAMA*, 288(6):715, aug 2002. URL: <https://doi.org/10.1001/jama.288.6.715>, doi:10.1001/jama.288.6.715.
- [72] Meritxell Espino Guarch, Mariona Font-Llitjós, Silvia Murillo-Cuesta, Ekaitz Errasti-Murugarren, Adelaida M Celaya, Georgia Giroto, Dragana Vuckovic, Massimo Mezzavilla, Clara Vilches, Susanna Bodoy, Ignasi Sahún, Laura González, Esther Prat, Antonio Zorzano, Mara Dierssen, Isabel Varela-Nieto, Paolo Gasparini, Manuel Palacín, and Virginia Nunes. Mutations in l-type amino acid transporter-2 support SLC7a8 as a novel gene involved in age-related hearing loss. *eLife*, 7, jan 2018. URL: <https://doi.org/10.7554/elife.31511>, doi:10.7554/elife.31511.
- [73] Albert Haas. The phagosome: Compartment with a license to kill. *Traffic*, 8(4):311–330, 2007. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1600-0854.2006.00531.x>, arXiv:<https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1600-0854.2006.00531.x>, doi:10.1111/j.1600-0854.2006.00531.x.
- [74] Fumiaki Hanzawa, Eri Sakuma, Saki Nomura, Tomono Uchida, Hiroaki Oda, and Saiko Ikeda. Excess α -tocopherol decreases extrahepatic phylloquinone in phylloquinone-fed rats but not menaquinone-4 in menaquinone-4-fed rats. *Molecular Nutrition & Food Research*, 58(8):1601–1609, apr 2014. URL: <https://doi.org/10.1002/mnfr.201300710>, doi:10.1002/mnfr.201300710.
- [75] Melissa L. Harris, Temesgen D. Fufa, Joseph W. Palmer, Sandeep S. Joshi, Denise M. Larson, Arturo Incao, Derek E. Gildea, Niraj S. Trivedi, Autumne N. Lee, Chi-Ping Day, Helen T. Michael, Thomas J. Hornyak, Glenn Merlino, NISC Comparative Sequencing Program, and William J. Pavan. A direct link between mitf, innate immunity, and hair graying. *PLOS Biology*, 16(5):1–27, 05 2018. URL: <https://doi.org/10.1371/journal.pbio.2003648>, doi:10.1371/journal.pbio.2003648.
- [76] I Hasanbasic, I Rajotte, and M Blostein. The role of γ -carboxylation in the anti-apoptotic function of gas6. *Journal of thrombosis and haemostasis : JTH*, 3:2790–7, 01 2006. URL: https://www.researchgate.net/publication/7409839_The_role_of_g-carboxylation_in_the_anti-apoptotic_function_of_Gas6, doi:10.1111/j.1538-7836.2005.01662.x.
- [77] S Haseli, N Khalili, M Bakhshayeshkaram, Taheri M Sanei, and Y Moharramzad. Lobar distribution of covid-19 pneumonia based on chest computed tomography findings; a retrospective study. *Archives of Academic Emergency Medicine*, 8(1), 2020. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC7212068/>.
- [78] Inaya Hayek, Fabian Fischer, Jan Schulze-Luehrmann, Katja Dettmer, Katharina Sobotta, Valentin Schatz, Lisa Kohl, Katharina Boden, Roland Lang, Peter J. Oefner, Stefan Wirtz, Jonathan Jantsch, and Anja Lhrmann. Limitation of TCA cycle intermediates represents an oxygen-independent nutritional antibacterial effector mechanism of macrophages. *Cell Reports*, 26(13):3502–3510.e6, mar 2019. URL: <https://doi.org/10.1016/j.celrep.2019.02.103>, doi:10.1016/j.celrep.2019.02.103.
- [79] Xia-Di He, Wei Gong, Jia-Nong Zhang, Ji Nie, Cui-Fang Yao, Fu-Shen Guo, Yan Lin, Xiao-Hui Wu, Feng Li, Jie Li, Wei-Cheng Sun, En-Duo Wang, Yan-Peng An, Hui-Ru Tang, Guo-Quan Yan, Peng-Yuan Yang, Yun Wei, Yun-Zi Mao, Peng-Cheng Lin, Jian-Yuan Zhao, Yanhui Xu, Wei Xu, and Shi-Min Zhao. Sensing and transmitting intracellular amino acid signals through reversible lysine aminoacylations. *Cell Metabolism*, 27(1):151–166.e6, jan 2018. URL: <https://doi.org/10.1016/j.cmet.2017.10.015>, doi:10.1016/j.cmet.2017.10.015.
- [80] Zhesheng He, Wencong Zhao, Wenchao Niu, Xuejiao Gao, Xingfa Gao, Yong Gong, and Xueyun Gao. Molecules inhibit

- the enzyme activity of 3-chymotrypsin-like cysteine protease of SARS-CoV-2 virus: the experimental and theory studies. may 2020. URL: <https://doi.org/10.1101/2020.05.28.120642>, doi:10.1101/2020.05.28.120642.
- [81] Collin D. Heer, Daniel J. Sanderson, Yousef M.O. Alhammad, Mark S. Schmidt, Samuel A.J. Trammell, Stanley Perlman, Michael S. Cohen, Anthony R. Fehr, and Charles Brenner. Coronavirus and PARP expression dysregulate the NAD metabolome: a potentially actionable component of innate immunity. apr 2020. URL: <https://doi.org/10.1101/2020.04.17.047480>, doi:10.1101/2020.04.17.047480.
- [82] Leonhard X. Heinz, JangEun Lee, Utkarsh Kapoor, Felix Kartnig, Vitaly Sedlyarov, Konstantinos Papakostas, Adrian César-Razquin, Patrick Essletzbichler, Ulrich Goldmann, Adrijana Stefanovic, Johannes W. Bigenzahn, Stefania Scorzoni, Mattia D. Pizzagalli, Ariel Bensimon, André C. Mller, F. James King, Jun Li, Enrico Girardi, M. Lamine Mbow, Charles E. Whitehurst, Manuele Rebsamen, and Giulio Superti-Furga. TASL is the SLC15a4-associated adaptor for IRF5 activation by TLR7–9. *Nature*, 581(7808):316–322, may 2020. URL: <https://doi.org/10.1038/s41586-020-2282-0>, doi:10.1038/s41586-020-2282-0.
- [83] LA Henderson, SW Canna, GS Schulert, S Volpi, PY Lee, KF Kernan, R Caricchio, S Mahmud, MM Hazen, O Halyabar, KJ Hoyt, J Han, AA Grom, M Gattorno, A Ravelli, Benedetti F De, EM Behrens, RQ Cron, and PA Nigrovic. On the alert for cytokine storm: Immunopathology in covid19. *Arthritis & Rheumatology (Hoboken, N.j.)*, 20200510. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC7262347/>.
- [84] Adeline Heurich, Heike Hofmann-Winkler, Stefanie Gierer, Thomas Liepold, Olaf Jahn, and Stefan Pöhlmann. Tmprss2 and adam17 cleave ace2 differentially and only proteolysis by tmprss2 augments entry driven by the severe acute respiratory syndrome coronavirus spike protein. *Journal of Virology*, 88(2):1293–1307, 2014. URL: <https://jvi.asm.org/content/88/2/1293>, arXiv:<https://jvi.asm.org/content/88/2/1293.full.pdf>, doi:10.1128/JVI.02202-13.
- [85] M Hinz, A Stein, T Cole, and P Ryan. Administration of supplemental l-tyrosine with phenelzine: a clinical literature review. *Clinical Pharmacology : Advances and Applications*, 6:107–10, 2014. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4112771/>.
- [86] Victoria Hodgkinson and Michael J. Petris. Copper homeostasis at the host-pathogen interface. *Journal of Biological Chemistry*, 287(17):13549–13555, 2012. URL: <http://www.jbc.org/content/287/17/13549.abstract>, arXiv:<http://www.jbc.org/content/287/17/13549.full.pdf+html>, doi:10.1074/jbc.R111.316406.
- [87] Jessica M. Hoffman, J. Veronika Kiklevich, Marika Austad, ViLinh Tran, Dean P. Jones, Angela Royal, Carolyn Henry, and Steven N. Austad. Tryptophan metabolism is differently regulated between large and small dogs. *GeroScience*, Nov 2019. URL: <https://doi.org/10.1007/s11357-019-00114-x>, doi:10.1007/s11357-019-00114-x.
- [88] Patrick M. Honore, David De Bels, Thierry Preseau, Sebastien Redant, and Herbert D. Spapen. Citrate: How to get started and what, when, and how to monitor? *Journal of Translational Internal Medicine*, 6(3):115–127, oct 2018. URL: <https://doi.org/10.2478/jtim-2018-0026>, doi:10.2478/jtim-2018-0026.
- [89] YC Hou, CL Lu, CM Zheng, RM Chen, YF Lin, WC Liu, TH Yen, R Chen, and KC Lu. Emerging role of vitamins d and k in modulating uremic vascular calcification: The aspect of passive calcification. *Nutrients*, 11(1), 2019. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC6356797/>.
- [90] Glen Huang, Alex J. Kovalic, and Christopher J. Graber. Prognostic value of leukocytosis and lymphopenia for coronavirus disease severity. *Emerging Infectious Diseases*, 26(8), aug 2020. URL: <https://doi.org/10.3201/eid2608.201160>, doi:10.3201/eid2608.201160.
- [91] X Huang, J Chen, W Cao, L Yang, Q Chen, J He, Q Yi, H Huang, E Zhang, and Z Cai. The many substrates and functions of nedd4-1. *Cell Death & Disease*, 10(12), 2019. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC6885513/>.
- [92] Mohammed Iddir, Alex Brito, Giulia Dingo, Sofia Fernandez Del Campo, Hanen Samouda, Michael La Frano, and Torsten Bohn. Strengthening the immune system and reducing inflammation and oxidative stress through diet and nutrition: Considerations during the COVID-19 crisis. *Nutrients*, 12(6):1562, may 2020. URL: <https://doi.org/10.3390/nu12061562>, doi:10.3390/nu12061562.
- [93] Theocharis Ispoglou, Kevin Deighton, Roderick FGJ King, Helen White, and Matthew Lees. Novel essential amino acid supplements enriched with l-leucine facilitate increased protein and energy intakes in older women: a randomised controlled trial. *Nutrition Journal*, 16(1), nov 2017. URL: <https://doi.org/10.1186/s12937-017-0298-6>, doi:10.1186/s12937-017-0298-6.
- [94] Hijiri ITO, Atsuko TAKAO, and Yoshihiko IDE. Relationship between the trace elements and abnormal electrocardiogram in bed-ridden patients. *The Japanese Journal of Rehabilitation Medicine*, 35(10):681–686, 1998. URL: https://www.jstage.jst.go.jp/article/jjrm1964/35/10/35_10_681/_article, doi:10.2490/jjrm1963.35.681.
- [95] D Ivanova, Z Zhelev, P Getsov, B Nikolova, I Aoki, T Higashi, and R Bakalova. Vitamin k: Redox-modulation, prevention of mitochondrial dysfunction and anticancer effect. *Redox Biology*, 16:352–8, 2018. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC5953218/>.
- [96] KL Ivey, MK Jensen, JM Hodgson, AH Eliassen, A Cassidy, and EB Rimm. Association of flavonoid-rich foods and flavonoids with risk of all-cause mortality. *The British journal of nutrition*, 117(10):1470–7, 2017. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC7233415/>.
- [97] A Jajodia, L Ebner, B Heidinger, A Chaturvedi, and H Prosch. Imaging in corona virus disease 2019 (covid-19)a scoping review. *European Journal of Radiology Open*, 7:100237, 2020. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC7211691/>.
- [98] Miguel Jiménez-Alcázar, Chandini Rangaswamy, Rachita Panda, Josephine Bitterling, Yashin J. Simsek, Andy T. Long, Rostyslav Bilyy, Veit Krenn, Christoph Renné, Thomas Renné, Stefan Kluge, Ulf Panzer, Ryushin Mizuta, Hans Georg Mannherz, Daisuke Kitamura, Martin Herrmann, Markus Napirei, and Tobias A. Fuchs. Host DNases prevent vascular occlusion by neutrophil extracellular traps. *Science*, 358(6367):1202–1206, nov 2017. URL: <https://doi.org/10.1126>

- science.aam8897, doi:10.1126/science.aam8897.
- [99] Li Jing, Zhang Yanyan, and Fan Junfeng. Acetic acid in aged vinegar affects molecular targets for thrombus disease management. *Food & Function*, 6(8):2845–2853, 2015. URL: <https://doi.org/10.1039/c5fo00327j>, doi:10.1039/c5fo00327j.
 - [100] Mary Ann Johnson. If high folic acid aggravates vitamin b12 deficiency what should be done about it? *Nutrition Reviews*, 65(10):451–458, jun 2008. URL: <https://doi.org/10.1111/j.1753-4887.2007.tb00270.x>, doi:10.1111/j.1753-4887.2007.tb00270.x.
 - [101] Eiji Kakazu, Noriatsu Kanno, Yoshiyuki Ueno, and Tooru Shimosegawa. Extracellular branched-chain amino acids, especially valine, regulate maturation and function of monocyte-derived dendritic cells. *The Journal of Immunology*, 179(10):7137–7146, 2007. URL: <https://www.jimmunol.org/content/179/10/7137>, arXiv:<https://www.jimmunol.org/content/179/10/7137.full.pdf>, doi:10.4049/jimmunol.179.10.7137.
 - [102] P Katulanda, HA Dissanayake, I Ranathunga, V Ratnasamy, PSA Wijewickrama, N Yogendranathan, KKK Gamage, Silva NL de, M Sumanatilleke, NP Somasundaram, and DR Matthews. Prevention and management of covid-19 among patients with diabetes: an appraisal of the literature. *Diabetologia*, pages 1–13, 20200514. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC7220850/>.
 - [103] T. Kawaguchi, T. Torimura, A. Takata, S. Satomi, and M. Sata. Valine, a branched-chain amino acid, reduced hcv viral load and led to eradication of hcv by interferon therapy in a decompensated cirrhotic patient. *Case Rep Gastroenterol*, 6(3), 2012. URL: <https://www.karger.com/DOI/10.1159/000343094>.
 - [104] Ying Ke, A. Grant Butt, Marianne Swart, Yong Feng Liu, and Fiona J. McDonald. COMMD1 downregulates the epithelial sodium channel through nedd4–2. *American Journal of Physiology-Renal Physiology*, 298(6):F1445–F1456, jun 2010. URL: <https://doi.org/10.1152/ajprenal.00257.2009>, doi:10.1152/ajprenal.00257.2009.
 - [105] W Khaliq, D Andreis, A Kleyman, M Grler, and M Singer. Reductions in tyrosine levels are associated with thyroid hormone and catecholamine disturbances in sepsis. *Intensive Care Medicine Experimental*, 3(Suppl 1), 2015. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4798095/>.
 - [106] Y Kim and OY Bang. Paradoxical procoagulant effect of early doses of warfarin: Possible role of non-vitamin k oral anticoagulant in patients with atrial fibrillation-related stroke. *Journal of Stroke*, 17(2):216–8, 2015. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4460341/>.
 - [107] LM Klevay. Cardiovascular disease from copper deficiency—a history. *The Journal of nutrition*, 130(2S Suppl):489S–492S, 2000. URL: <http://www.ncbi.nlm.nih.gov/pubmed/articles/10721936/>.
 - [108] Toshihiko Kobayashi, Shiho Shimabukuro-Demoto, Reiko Yoshida-Sugitani, Kaori Furuyama-Tanaka, Hitomi Karyu, Yuki Sugiura, Yukiko Shimizu, Toshiaki Hosaka, Motohito Goto, Norihiro Kato, Tadashi Okamura, Makoto Suematsu, Shigeyuki Yokoyama, and Noriko Toyama-Sorimachi. The histidine transporter slc15a4 coordinates mtor-dependent inflammatory responses and pathogenic antibody production. *Immunity*, 41(3):375 – 388, 2014. URL: <http://www.sciencedirect.com/science/article/pii/S1074761314003057>, doi:<https://doi.org/10.1016/j.immuni.2014.08.011>.
 - [109] Joel D. Kopple. Phenylalanine and tyrosine metabolism in chronic kidney failure. *The Journal of Nutrition*, 137(6):1586S–1590S, jun 2007. URL: <https://doi.org/10.1093/jn/137.6.1586s>, doi:10.1093/jn/137.6.1586s.
 - [110] Mirjana K. Kovac, Sanja Z. Lalic-Cosic, Jelena M. Dmitrovic, Valentina J. Djordjevic, and Dragica P. Radojkovic. Thrombin generation, d-dimer and protein s in uncomplicated pregnancy. *Clinical Chemistry and Laboratory Medicine (CCLM)*, 53(12), jan 2015. URL: <https://doi.org/10.1515/cclm-2014-1030>, doi:10.1515/cclm-2014-1030.
 - [111] M. Anna Kowalska, Guohua Zhao, Li Zhai, George David, Stephen Marcus, Sriram Krishnaswamy, and Mortimer Poncz. Modulation of protein c activation by histones, platelet factor 4, and heparinoids. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 34(1):120–126, jan 2014. URL: <https://doi.org/10.1161/atvbaha.113.302236>, doi:10.1161/atvbaha.113.302236.
 - [112] Sanjeev Kumar, William P. Malachowski, James B. DuHadaway, Judith M. LaLonde, Patrick J. Carroll, Daniel Jaller, Richard Metz, George C. Prendergast, and Alexander J. Muller. Indoleamine 2,3-dioxygenase is the anticancer target for a novel series of potent naphthoquinone-based inhibitors. *Journal of Medicinal Chemistry*, 51(6):1706–1718, mar 2008. URL: <https://doi.org/10.1021/jm7014155>, doi:10.1021/jm7014155.
 - [113] Z Kurugl, F Vardar, F zkinay, K Kavakh, and C zkinay. Lupus anticoagulant and protein s deficiency in otherwise healthy children with acute varicella infection. *Acta Paediatrica*, 89(10):1186–1189, 2000. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1651-2227.2000.tb00733.x>, arXiv:<https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1651-2227.2000.tb00733.x>, doi:10.1111/j.1651-2227.2000.tb00733.x.
 - [114] Adrian C. Kuyper and H. A. Mattill. Some aspects of citric acid metabolism. *Journal of Biological Chemistry*, 103(1):51–60, 1933. URL: <http://www.jbc.org/content/103/1/51.short>, arXiv:<http://www.jbc.org/content/103/1/51.full.pdf+html>.
 - [115] Jun Lan, Jiwan Ge, Jinfang Yu, Sisi Shan, Huan Zhou, Shilong Fan, Qi Zhang, Xuanling Shi, Qisheng Wang, Linqi Zhang, and Xinquan Wang. Structure of the SARS-CoV-2 spike receptor-binding domain bound to the ACE2 receptor. *Nature*, 581(7807):215–220, mar 2020. URL: <https://doi.org/10.1038/s41586-020-2180-5>, doi:10.1038/s41586-020-2180-5.
 - [116] M Larsen, C Bayard, H Lepetitcorps, J Cohen-Bittan, V Appay, J Boddaert, and D Sauce. Elevated neopterin levels predict early death in older hip-fracture patients. *EBioMedicine*, 26:157–64, 2017. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC5832560/>.
 - [117] GK Lee, HJ Park, M Macleod, P Chandler, DH Munn, and AL Mellor. Tryptophan deprivation sensitizes activated t cells to apoptosis prior to cell division. *Immunology*, 107(4):452–60, 2002. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1782830/>.

- [118] Ji-Hyun Lee, Seon-Ae Jeon, Byung-Gyu Kim, Michiko Takeda, Jae-Jin Cho, Dong-Ik Kim, Hiroshi Kawabe, and Je-Yoel Cho. Nedd4 deficiency in vascular smooth muscle promotes vascular calcification by stabilizing psmad1. *Journal of Bone and Mineral Research*, 32(5):927–938, 2017. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1002/jbmr.3073>, [arXiv:https://onlinelibrary.wiley.com/doi/pdf/10.1002/jbmr.3073](https://onlinelibrary.wiley.com/doi/pdf/10.1002/jbmr.3073), doi:10.1002/jbmr.3073.
- [119] Chong Lei, Binxiao Su, Hailong Dong, Andrea Bellavia, Raffaele Di Fenza, Bijan Safaei Fakhr, Stefano Gianni, Luigi Giuseppe Grassi, Robert Kacmarek, Caio Cesar Araujo Moraes, Riccardo Pinciroli, Emanuele Vassena, and Lorenzo Berra. Protocol of a randomized controlled trial testing inhaled nitric oxide in mechanically ventilated patients with severe acute respiratory syndrome in covid-19 (sars-cov-2). *medRxiv*, 2020. URL: <https://www.medrxiv.org/content/early/2020/05/24/2020.03.09.20033530>, [arXiv:https://www.medrxiv.org/content/early/2020/05/24/2020.03.09.20033530.full.pdf](https://www.medrxiv.org/content/early/2020/05/24/2020.03.09.20033530.full.pdf), doi:10.1101/2020.03.09.20033530.
- [120] Greg Lemke and Gregg J. Silverman. Blood clots and TAM receptor signalling in COVID-19 pathogenesis. *Nature Reviews Immunology*, 20(7):395–396, jun 2020. URL: <https://doi.org/10.1038/s41577-020-0354-x>, doi:10.1038/s41577-020-0354-x.
- [121] W Leslie and C Hankey. Aging, nutritional status and health. *Healthcare*, 3(3):648–58, 2015. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4939559/>.
- [122] Marcel Levi, Jecko Thachil, Toshiaki Iba, and Jerrold H Levy. Coagulation abnormalities and thrombosis in patients with COVID-19. *The Lancet Haematology*, may 2020. URL: [https://doi.org/10.1016/s2352-3026\(20\)30145-9](https://doi.org/10.1016/s2352-3026(20)30145-9), doi:10.1016/s2352-3026(20)30145-9.
- [123] M Leyton. Effects on mood of acute phenylalanine/tyrosine depletion in healthy women. *Neuropsychopharmacology*, 22(1):52–63, jan 2000. URL: [https://doi.org/10.1016/s0893-133x\(99\)00086-x](https://doi.org/10.1016/s0893-133x(99)00086-x), doi:10.1016/s0893-133x(99)00086-x.
- [124] Peng Li, Yu-Long Yin, Defa Li, Sung Woo Kim, and Guoyao Wu. Amino acids and immune function. *British Journal of Nutrition*, 98(2):237252, 2007. URL: <https://www.cambridge.org/core/journals/british-journal-of-nutrition/article/amino-acids-and-immune-function/B1A9C1587A8602613F6447BA8404D8E1/core-reader#>, doi:10.1017/S000711450769936X.
- [125] Q Li, L Wang, C Dong, Y Che, L Jiang, L Liu, H Zhao, Y Liao, Y Sheng, S Dong, and S Ma. The interaction of the sars coronavirus non-structural protein 10 with the cellular oxido-reductase system causes an extensive cytopathic effect. *Journal of Clinical Virology*, 34(2):133–9, 2005. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC7108382/>.
- [126] Andrew W. Lindsley, Justin T. Schwartz, and Marc E. Rothenberg. Eosinophil responses during COVID-19 infections and coronavirus vaccination. *Journal of Allergy and Clinical Immunology*, apr 2020. URL: <https://doi.org/10.1016/j.jaci.2020.04.021>, doi:10.1016/j.jaci.2020.04.021.
- [127] Brea Lipe and Deborah L. Ornstein. Deficiencies of natural anticoagulants, protein c, protein s, and antithrombin. *Circulation*, 124(14):e365–e368, 2011. URL: <https://www.ahajournals.org/doi/abs/10.1161/CIRCULATIONAHA.111.044412>, [arXiv:https://www.ahajournals.org/doi/pdf/10.1161/CIRCULATIONAHA.111.044412](https://www.ahajournals.org/doi/pdf/10.1161/CIRCULATIONAHA.111.044412), doi:10.1161/CIRCULATIONAHA.111.044412.
- [128] Jiheng Liu, Heng Li, Ming Luo, Jiyang Liu, Lingzhen Wu, Xianfeng Lin, Ruijuan Li, Zhihua Wang, Haiying Zhong, Wenli Zheng, Yan Zhou, Dixuan Jiang, Xin Tan, Zhiguo Zhou, Hongling Peng, and Guangsen Zhang. Lymphopenia acted as an adverse factor for severity in patients with COVID-19: a single-centered, retrospective study. apr 2020. URL: <https://doi.org/10.21203/rs.3.rs-22849/v1>, doi:10.21203/rs.3.rs-22849/v1.
- [129] Yang Liu, Bei Mao, Shuo Liang, Jia-wei Yang, Hai-wen Lu, Yan-hua Chai, Lan Wang, Li Zhang, Qiu-hong Li, Lan Zhao, Yan He, Xiao-long Gu, Xiao-bin Ji, Li Li, Zhi-jun Jie, Qiang Li, Xiang-yang Li, Hong-zhou Lu, Wen-hong Zhang, Yuan-lin Song, Jie-ming Qu, and Jin-fu Xu. Association between ages and clinical characteristics and outcomes of coronavirus disease 2019. *European Respiratory Journal*, 2020. URL: <https://erj.ersjournals.com/content/early/2020/04/16/13993003.01112-2020>, [arXiv:https://erj.ersjournals.com/content/early/2020/04/16/13993003.01112-2020.full.pdf](https://erj.ersjournals.com/content/early/2020/04/16/13993003.01112-2020.full.pdf), doi:10.1183/13993003.01112-2020.
- [130] Yuwei Liu, Xuebei Du, Jing Chen, Yalei Jin, Li Peng, Harry H.X. Wang, Mingqi Luo, Ling Chen, and Yan Zhao. Neutrophil-to-lymphocyte ratio as an independent risk factor for mortality in hospitalized patients with covid-19. *Journal of Infection*, 2020. URL: <http://www.sciencedirect.com/science/article/pii/S0163445320302085>, doi:https://doi.org/10.1016/j.jinf.2020.04.002.
- [131] Christopher A. Lopez and Eric P. Skaar. The impact of dietary transition metals on host-bacterial interactions. *Cell Host & Microbe*, 23(6):737–748, jun 2018. URL: <https://doi.org/10.1016/j.chom.2018.05.008>, doi:10.1016/j.chom.2018.05.008.
- [132] Inken Lorenzen, Julianne Lokau, Yvonne Korpys, Mirja Oldefest, Charlotte M. Flynn, Ulrike Knzel, Christoph Garbers, Matthew Freeman, Joachim Grtzing, and Stefan Dsterhft. Control of ADAM17 activity by regulation of its cellular localisation. *Scientific Reports*, 6(1), oct 2016. URL: <https://doi.org/10.1038/srep35067>, doi:10.1038/srep35067.
- [133] Sean M. Lynch and Leslie M. Klevay. Effects of a dietary copper deficiency on plasma coagulation factor activities in male and female mice. *The Journal of Nutritional Biochemistry*, 3(8):387 – 391, 1992. URL: <http://www.sciencedirect.com/science/article/pii/S0955286392900128>, doi:https://doi.org/10.1016/0955-2863(92)90012-8.
- [134] Jan Niklas Lke, Felix Neumaier, Serdar Alpdogan, Jrgen Hescheler, Toni Schneider, Walid Albanna, and Isha Akhtar-Schfer. Submicromolar copper (II) ions stimulate transretinal signaling in the isolated retina from wild type but not from cav2.3-deficient mice. *BMC Ophthalmology*, 20(1), may 2020. URL: <https://doi.org/10.1186/s12886-020-01451-8>, doi:10.1186/s12886-020-01451-8.
- [135] David J. Maggs, B. Keith Collins, James G. Thorne, and Mark P. Nasisse. Effects of l-lysine and l-arginine on in vitro replication of feline herpesvirus type-1. *American Journal of Veterinary Research*, 61(12):1474–1478, dec 2000. URL:

- <https://doi.org/10.2460/ajvr.2000.61.1474>, doi:10.2460/ajvr.2000.61.1474.
- [136] OA Malakhova and DE Zhang. Isg15 inhibits nedd4 ubiquitin e3 activity and enhances the innate antiviral response*. *The Journal of Biological Chemistry*, 283(14):8783–7, 2008. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2276364/>.
 - [137] Sathyavani Malyala, Yizhu Zhang, Jasiel O. Strubbe, and Jason N. Bazil. Calcium phosphate precipitation inhibits mitochondrial energy metabolism. *PLOS Computational Biology*, 15(1):1–19, 01 2019. URL: <https://doi.org/10.1371/journal.pcbi.1006719>, doi:10.1371/journal.pcbi.1006719.
 - [138] M.J. Marchywka. Canine heartworm treated with doxycycline, ivermectin and various supplements. Technical Report MJM-2019-001, not institutionalized , independent, 306 Charles Cox , Canton GA 30115, manuscript in preparation. May be recycled in appropriate media.
 - [139] M.J. Marchywka. Less trendy properties of common molecules. Technical Report MJM-2019-004, not institutionalized , independent, 306 Charles Cox , Canton GA 30115, manuscript in preparation. May be recycled in appropriate media.
 - [140] M.J. Marchywka. Notebook - experiences with supplemental vitamin k, copper, and herbs in dogs discussed in terms of inappropriate mineralization and disease. Technical Report MJM-2015-001, not institutionalized , independent, 306 Charles Cox , Canton GA 30115, manuscript in preparation. Not originally meant to publish, lab notebook.
 - [141] M.J. Marchywka. Old age as starvation. Technical Report MJM-2019-005, not institutionalized , independent, 306 Charles Cox , Canton GA 30115, manuscript in preparation. May be recycled in appropriate media.
 - [142] M.J. Marchywka. A series of informal canine cases, a methodology and hypotheses on therapeutic uses of copper and vitamins for dogs. Technical Report MJM-2017-001, not institutionalized , independent, 306 Charles Cox , Canton GA 30115, manuscript in preparation. May be recycled in appropriate media.
 - [143] Jan Martel, Yun-Fei Ko, John D. Young, and David M. Ojcius. Could nasal nitric oxide help to mitigate the severity of covid-19? *Microbes and Infection*, 22(4):168 – 171, 2020. URL: <http://www.sciencedirect.com/science/article/pii/S1286457920300800>, doi:<https://doi.org/10.1016/j.micinf.2020.05.002>.
 - [144] Michael A. Matthay and Lorraine B. Ware. Plasma protein c levels in patients with acute lung injury: Prognostic significance. *Critical Care Medicine*, 32(Supplement):S229–S232, may 2004. URL: <https://doi.org/10.1097/01.ccm.0000126121.56990.d3>, doi:10.1097/01.ccm.0000126121.56990.d3.
 - [145] Braedon McDonald, Rachelle P. Davis, Seok-Joo Kim, Mandy Tse, Charles T. Esmon, Elzbieta Kolaczowska, and Craig N. Jenne. Platelets and neutrophil extracellular traps collaborate to promote intravascular coagulation during sepsis in mice. *Blood*, 129(10):1357–1367, mar 2017. URL: <https://doi.org/10.1182/blood-2016-09-741298>, doi:10.1182/blood-2016-09-741298.
 - [146] Eileen McNeill, Elena Stylianou, Mark J. Crabtree, Rachel Harrington-Kandt, Anna-Lena Kolb, Marina Diotallevi, Ashley B. Hale, Paulo Bettencourt, Rachel Tanner, Matthew K. O'Shea, Magali Matsumiya, Helen Lockstone, Julius Mller, Helen A. Fletcher, David R. Greaves, Helen McShane, and Keith M. Channon. Regulation of mycobacterial infection by macrophage gh1 and tetrahydrobiopterin. *Nature Communications*, 9(1), dec 2018. URL: <https://doi.org/10.1038/s41467-018-07714-9>, doi:10.1038/s41467-018-07714-9.
 - [147] GaryG. Meadows, RokiaM. Abdallah, JeanR. Starkey, and CherylJ. Aslakson. Response of natural killer cells from dietary tyrosine-and phenylalanine-restricted mice to biological response modifiers. *Cancer Immunology Immunotherapy*, 26(1), feb 1988. URL: <https://doi.org/10.1007/bf00199850>, doi:10.1007/bf00199850.
 - [148] D B Milne and F H Nielsen. Effects of a diet low in copper on copper-status indicators in postmenopausal women. *The American Journal of Clinical Nutrition*, 63(3):358–364, mar 1996. URL: <https://doi.org/10.1093/ajcn/63.3.358>, doi:10.1093/ajcn/63.3.358.
 - [149] SK Mistry, Z Greenfeld, SM Morris, and C Baylis. The intestinalrenal arginine biosynthetic axis in the aging rat. *Mechanisms of ageing and development*, 123(8):1159–65, 2002. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2745251/>.
 - [150] Kenji Mizumoto, Ryosuke Omori, and Hiroshi Nishiura. Age specificity of cases and attack rate of novel coronavirus disease (covid-19). *medRxiv*, 2020. URL: <https://www.medrxiv.org/content/early/2020/03/13/2020.03.09.20033142>, arXiv:<https://www.medrxiv.org/content/early/2020/03/13/2020.03.09.20033142.full.pdf>, doi:10.1101/2020.03.09.20033142.
 - [151] Soenke Moehn, Paul B. Pencharz, and Ronald O. Ball. Lessons Learned Regarding Symptoms of Tryptophan Deficiency and Excess from Animal Requirement Studies. *The Journal of Nutrition*, 142(12):2231S–2235S, 10 2012. URL: <https://doi.org/10.3945/jn.112.159061>, arXiv:<https://academic.oup.com/jn/article-pdf/142/12/2231S/29585314/2231s.pdf>, doi:10.3945/jn.112.159061.
 - [152] C Mozzini and D Girelli. The role of neutrophil extracellular traps in covid-19: Only an hypothesis or a potential new field of research? *Thrombosis Research*, 20200427. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC7184981/>.
 - [153] MF Murray. Tryptophan depletion and hiv infection: a metabolic link to pathogenesis. *The Lancet. Infectious diseases*, 3(10):644–52, 2003. URL: <http://www.ncbi.nlm.nih.gov/pubmed/articles/14522263/>.
 - [154] Michael F Murray, Mary Langan, and Rob Roy MacGregor. Increased plasma tryptophan in hiv-infected patients treated with pharmacologic doses of nicotinamide. *Nutrition*, 17(7):654 – 656, 2001. URL: <http://www.sciencedirect.com/science/article/pii/S0899900701005688>, doi:[https://doi.org/10.1016/S0899-9007\(01\)00568-8](https://doi.org/10.1016/S0899-9007(01)00568-8).
 - [155] T. Nagatsu and H. Ichinose. Regulation of pteridine-requiring enzymes by the cofactor tetrahydrobiopterin. *Molecular Neurobiology*, 19(1):79–96, feb 1999. URL: <https://doi.org/10.1007/bf02741379>, doi:10.1007/bf02741379.
 - [156] H. Namkung, J. Gong, H. Yu, and C. F. M. de Lange. Effect of pharmacological intakes of zinc and copper on growth performance, circulating cytokines and gut microbiota of newly weaned piglets challenged with coliform lipopolysaccharides. *Canadian Journal of Animal Science*, 86(4):511–522, dec 2006. URL: <https://doi.org/10.4141/a05-075>,

- doi:10.4141/a05-075.
- [157] M. Negro, S. Perna, D. Spadaccini, L. Castelli, L. Calanni, M. Barbero, C. Cescon, M. Rondanelli, and Giuseppe D'Antona. Effects of 12 weeks of essential amino acids (eaa)-based multi-ingredient nutritional supplementation on muscle mass, muscle strength, muscle power and fatigue in healthy elderly subjects: A randomized controlled double-blind study. *The journal of nutrition, health & aging*, 23(5):414–424, May 2019. URL: <https://doi.org/10.1007/s12603-019-1163-4>, doi:10.1007/s12603-019-1163-4.
 - [158] P. Nguyễn, M. Munzer, J. Reynaud, O. Richard, P. Pouzol, and P. François. Varicella and thrombotic complications associated with transient protein c and protein s deficiencies in children. *European Journal of Pediatrics*, 153(9):646–649, Sep 1994. URL: <https://doi.org/10.1007/BF02190684>, doi:10.1007/BF02190684.
 - [159] P. G. NORTHUP, V. SUNDARAM, M. B. FALLON, K. R. REDDY, R. A. BALOGUN, A. J. SANYAL, Q. M. ANSTEE, M. R. HOFFMAN, Y. IKURA, S. H. CALDWELL, and THE COAGULATION IN LIVER DISEASE GROUP. Hypercoagulation and thrombophilia in liver disease. *Journal of Thrombosis and Haemostasis*, 6(1):2–9, 2008. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1538-7836.2007.02772.x>, arXiv:<https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1538-7836.2007.02772.x>, doi:10.1111/j.1538-7836.2007.02772.x.
 - [160] Blessing Ogunlade, Jesse J Guidry, Eric Lazartigues, and Catalin Filipeanu. Ace2 internalization and degradation is controlled by ubiquitin ligase nedd4. *The FASEB Journal*, 33(1-supplement):719.14–719.14, 2019. URL: https://www.fasebj.org/doi/abs/10.1096/fasebj.2019.33.1_supplement.719.14, arXiv:https://www.fasebj.org/doi/pdf/10.1096/fasebj.2019.33.1_supplement.719.14, doi:10.1096/fasebj.2019.33.1_supplement.719.14.
 - [161] Curtis T. Okamoto. Regulation of transporters and channels by membrane-trafficking complexes in epithelial cells. *Cold Spring Harbor Perspectives in Biology*, 9(11):a027839, feb 2017. URL: <https://doi.org/10.1101/cshperspect.a027839>, doi:10.1101/cshperspect.a027839.
 - [162] Hikaru Okamoto, Yasuaki Okamoto, Takeshi Hirokawa, and Andrei Timerbaev. Trace ion analysis of sea water by capillary electrophoresis: Determination of strontium and lithium pre-concentrated by transient isotachopheresis. *The Analyst*, 128:1439–42, 01 2004. URL: https://www.researchgate.net/publication/8908634_Trace_ion_analysis_of_sea_water_by_capillary_electrophoresis_Determination_of_strontium_and_lithium_pre-concentrated_by_transient_isotachopheresis, doi:10.1039/b310345p.
 - [163] Ricardo Brandt Oliveira, Ana Lourdes Candolo Martinelli, Luiz Ernesto Almeida Troncon, and Jorge Elias Jr. Small intestinal bacterial overgrowth (SIBO) and vitamin k-responsive coagulopathy: a previously unrecorded association. *BMJ Case Reports*, pages bcr-2017-223531, jun 2018. URL: <https://doi.org/10.1136/bcr-2017-223531>, doi:10.1136/bcr-2017-223531.
 - [164] Claudia M. Oomen, Marjan J. van Erk, Edith J. M. Feskens, Frans J. Kok, and Daan Kromhout. Arginine intake and risk of coronary heart disease mortality in elderly men. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 20(9):2134–2139, 2000. URL: <https://www.ahajournals.org/doi/abs/10.1161/01.ATV.20.9.2134>, arXiv:<https://www.ahajournals.org/doi/pdf/10.1161/01.ATV.20.9.2134>, doi:10.1161/01.ATV.20.9.2134.
 - [165] Isabelle Papet, Benot Ruot, Denis Breuille, Stephane Walrand, Marie-Chantal Farges, Marie-Paule Vasson, and Christiane Obled. Bacterial Infection Affects Protein Synthesis in Primary Lymphoid Tissues and Circulating Lymphocytes of Rats. *The Journal of Nutrition*, 132(7):2028–2032, 07 2002. URL: <https://doi.org/10.1093/jn/132.7.2028>, arXiv:<https://academic.oup.com/jn/article-pdf/132/7/2028/24013818/4w0702002028.pdf>, doi:10.1093/jn/132.7.2028.
 - [166] Sang Gyu Park, Paul Schimmel, and Sunghoon Kim. Aminoacyl trna synthetases and their connections to disease. *Proceedings of the National Academy of Sciences*, 105(32):11043–11049, 2008. URL: <https://www.pnas.org/content/105/32/11043>, arXiv:<https://www.pnas.org/content/105/32/11043.full.pdf>, doi:10.1073/pnas.0802862105.
 - [167] G. Parker and H. Brotchie. Mood effects of the amino acids tryptophan and tyrosine. *Acta Psychiatrica Scandinavica*, 124(6):417–426, 2011. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1600-0447.2011.01706.x>, arXiv:<https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1600-0447.2011.01706.x>, doi:10.1111/j.1600-0447.2011.01706.x.
 - [168] I Piscaer, den Ouweland JMW van, K Vermeersch, NL Reynaert, FME Franssen, S Keene, EFM Wouters, W Janssens, C Vermeer, and R Janssen. Low vitamin k status is associated with increased elastin degradation in chronic obstructive pulmonary disease. *Journal of Clinical Medicine*, 8(8), 2019. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC6724066/>.
 - [169] Giuseppe Pontrelli, Alessandra M Martino, Hyppolite K Tchidjou, Rita Citton, Nadia Mora, Lucilla Ravà, Alberto E Tozzi, Paolo Palma, Maurizio Muraca, Elisabetta Franco, Paolo Rossi, and Stefania Bernardi. HIV is associated with thrombophilia and high d-dimer in children and adolescents. *AIDS*, 24(8):1145–1151, may 2010. URL: <https://doi.org/10.1097/qad.0b013e328337b9a0>, doi:10.1097/qad.0b013e328337b9a0.
 - [170] Roger J. S. Preston, Sinh Tran, Jennifer A. Johnson, Fionnuala Ní Áinle, Shona Harmon, Barry White, Owen P. Smith, P. Vince Jenkins, Björn Dahlbäck, and James S. O'Donnell. Platelet factor 4 impairs the anticoagulant activity of activated protein c. *Journal of Biological Chemistry*, 284(9):5869–5875, jan 2009. URL: <https://doi.org/10.1074/jbc.M804703200>, doi:10.1074/jbc.M804703200.
 - [171] Joseph R. Prohaska and Lois J. Heller. Calcium reintroduction decreases viability of cardiac myocytes from copper-deficient rats. *The Journal of Nutrition*, 129(10):1842–1845, oct 1999. URL: <https://doi.org/10.1093/jn/129.10.1842>, doi:10.1093/jn/129.10.1842.
 - [172] Syamal Raha, Rahul Mallick, Sanjay Basak, and Asim K. Duttaroy. Is copper beneficial for covid-19 patients? *Medical Hypotheses*, 142:109814, 2020. URL: <http://www.sciencedirect.com/science/article/pii/S0306987720308136>, doi: <https://doi.org/10.1016/j.mehy.2020.109814>.

- [173] Jian-Guo Ren, Pankaj Seth, Huihui Ye, Kun Guo, Jun-ichi Hanai, Zaheed Husain, and Vikas Sukhatme. Citrate suppresses tumor growth in multiple models through inhibition of glycolysis, the tricarboxylic acid cycle and the igf-1r pathway. *Scientific Reports*, 7, 12 2017. URL: https://www.researchgate.net/publication/318166179_Citrate_Suppresses_Tumor_Growth_in_Multiple_Models_through_Inhibition_of_Glycolysis_the_Tricarboxylic_Acid_Cycle_and_the_IGF-1R_Pathway, doi:10.1038/s41598-017-04626-4.
- [174] DM Richard, MA Dawes, CW Mathias, A Acheson, N Hill-Kapturczak, and DM Dougherty. L-tryptophan: Basic metabolic functions, behavioral research and therapeutic indications. *International Journal of Tryptophan Research : IJTR*, 2:45–60, 2009. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2908021/>.
- [175] CV Rothlin, EA Carrera-Silva, L Bosurgi, and S Ghosh. Tam receptor signaling in immune homeostasis. *Annual review of immunology*, 33:355–91, 2015. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4491918/>.
- [176] Raúl A. Ruggiero, Juan Bruzzo, Paula Chiarella, Pedro di Gianni, Martín A. Isturiz, Susana Linskens, Norma Speziale, Roberto P. Meiss, Oscar D. Bustuabad, and Christiane D. Pasqualini. Tyrosine isomers mediate the classical phenomenon of concomitant tumor resistance. *Cancer Research*, 71(22):7113–7124, 2011. URL: <https://cancerres.aacrjournals.org/content/71/22/7113>, arXiv:<https://cancerres.aacrjournals.org/content/71/22/7113.full.pdf>, doi:10.1158/0008-5472.CAN-11-0581.
- [177] Robert M. Russell. Factors in Aging that Effect the Bioavailability of Nutrients. *The Journal of Nutrition*, 131(4):1359S–1361S, 04 2001. URL: <https://doi.org/10.1093/jn/131.4.1359S>, arXiv:<https://academic.oup.com/jn/article-pdf/131/4/1359S/30094645/4w04t1s1359.pdf>, doi:10.1093/jn/131.4.1359S.
- [178] C Saade, E Najem, K Asmar, R Salman, Achkar B El, and L Naffaa. Intracranial calcifications on ct: an updated review. *Journal of Radiology Case Reports*, 13(8):1–18, 2019. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC6738489/>.
- [179] J Saban-Ruiz and D Ly-Pen. Covid-19: A personalized cardiometabolic approach for reducing complications and costs. the role of aging beyond topics. *The Journal of Nutrition, Health & Aging*, pages 1–10, 20200512. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC7217344/>.
- [180] Toshiro Sato, Naoko Inaba, and Takatoshi Yamashita. Mk-7 and its effects on bone quality and strength. *Nutrients*, 12(4), 2020. URL: <https://www.mdpi.com/2072-6643/12/4/965>, doi:10.3390/nu12040965.
- [181] Adil Sattar, Jane E. Willman, and Raghu Kolluri. Possible warfarin resistance due to interaction with ascorbic acid: Case report and literature review. *American Journal of Health-System Pharmacy*, 70(9):782–786, may 2013. URL: <https://doi.org/10.2146/ajhp110704>, doi:10.2146/ajhp110704.
- [182] Naveed Sattar, Iain B. McInnes, and John J. V. McMurray. Obesity a risk factor for severe covid-19 infection: Multiple potential mechanisms. *Circulation*, 0(0). URL: <https://www.ahajournals.org/doi/abs/10.1161/CIRCULATIONAHA.120.047659>, arXiv:<https://www.ahajournals.org/doi/pdf/10.1161/CIRCULATIONAHA.120.047659>, doi:10.1161/CIRCULATIONAHA.120.047659.
- [183] SP Saxena, T Fan, M Li, ED Israels, and LG Israels. A novel role for vitamin k1 in a tyrosine phosphorylation cascade during chick embryogenesis. *Journal of Clinical Investigation*, 99(4):602–7, 1997. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC507841/>.
- [184] Tobias Schneider, Daniel Caviezel, C Korcan Ayata, Caroline Kiss, Jan Hendrik Niess, and Petr Hruz. The Copper/Zinc Ratio Correlates With Markers of Disease Activity in Patients With Inflammatory Bowel Disease. *Crohn's & Colitis 360*, 2(1), 01 2020. otaa001. URL: <https://doi.org/10.1093/crocol/otaa001>, arXiv:<https://academic.oup.com/crohnscolitis360/article-pdf/2/1/otaa001/33030054/otaa001.pdf>, doi:10.1093/crocol/otaa001.
- [185] Dewald Schoeman and Burtram C. Fielding. Coronavirus envelope protein: current knowledge. *Virology Journal*, 16(1), may 2019. URL: <https://doi.org/10.1186/s12985-019-1182-0>, doi:10.1186/s12985-019-1182-0.
- [186] Lea Schroeder and Amy E. Ikui. Tryptophan confers resistance to sds-associated cell membrane stress in saccharomyces cerevisiae. *PLOS ONE*, 14(3):1–16, 03 2019. URL: <https://doi.org/10.1371/journal.pone.0199484>, doi:10.1371/journal.pone.0199484.
- [187] E Schrom, M Huber, M Aneja, C Dohmen, D Emrich, J Geiger, G Hasenpusch, A Herrmann-Janson, V Kretzschmann, O Mykhailik, T Pasewald, P Oak, A Hilgendorff, D Wohlleber, HG Hoymann, D Schaudien, C Plank, C Rudolph, and R Kubisch-Dohmen. Translation of angiotensin-converting enzyme 2 upon liver- and lung-targeted delivery of optimized chemically modified mrna. *Molecular Therapy. Nucleic Acids*, 7:350–65, 2017. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC5423349/>.
- [188] Cecilia Schuck. Urinary excretion of citric acid. *The Journal of Nutrition*, 7(6):691–700, jun 1934. URL: <https://doi.org/10.1093/jn/7.6.691>, doi:10.1093/jn/7.6.691.
- [189] Rifka C Schulman and Jeffrey I Mechanick. Metabolic and nutrition support in the chronic critical illness syndrome. *Respiratory Care*, 57(6):958–978, 2012. URL: <http://rc.rcjournal.com/content/57/6/958>, arXiv:<http://rc.rcjournal.com/content/57/6/958.full.pdf>, doi:10.4187/respcare.01620.
- [190] Steven P. Schulman, Lewis C. Becker, David A. Kass, Hunter C. Champion, Michael L. Terrin, Sandra Forman, Kavita V. Ernst, Mark D. Kelemen, Susan N. Townsend, Anne Capriotti, Joshua M. Hare, and Gary Gerstenblith. L-arginine therapy in acute myocardial infarction. *JAMA*, 295(1):58, jan 2006. URL: <https://doi.org/10.1001/jama.295.1.58>, doi:10.1001/jama.295.1.58.
- [191] R Serge, J Vandromme, and M Charlotte. Are we equal in adversity? does covid-19 affect women and men differently? *Maturitas*, 20200515. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC7227588/>.
- [192] Kenneth D.R. Setchell, James E. Heubi, Sohela Shah, Joel E. Lavine, David Suskind, Mohammed Al’Edreesi, Carol Potter, David W. Russell, Nancy C. O’Connell, Brian Wolfe, Pinky Jha, Wujuan Zhang, Kevin E. Bove, Alex S. Knisely, Alan F. Hofmann, Philip Rosenthal, and Laura N. Bull. Genetic defects in bile acid conjugation cause fat-soluble vitamin deficiency. *Gastroenterology*, 144(5):945–955.e6, may 2013. URL: <https://doi.org/10.1053/j.gastro.2013.02.004>,

- doi:10.1053/j.gastro.2013.02.004.
- [193] Jian Shang, Yushun Wan, Chuming Luo, Gang Ye, Qibin Geng, Ashley Auerbach, and Fang Li. Cell entry mechanisms of sars-cov-2. *Proceedings of the National Academy of Sciences*, 117(21):11727–11734, 2020. URL: <https://www.pnas.org/content/117/21/11727>, arXiv:<https://www.pnas.org/content/117/21/11727.full.pdf>, doi:10.1073/pnas.2003138117.
 - [194] Martin J. Shearer and Paul Newman. Recent trends in the metabolism and cell biology of vitamin k with special reference to vitamin k cycling and menaquinone-4 biosynthesis. *Journal of Lipid Research*, 2014. URL: <http://www.jlr.org/content/early/2014/01/31/jlr.R045559.abstract>, arXiv:<http://www.jlr.org/content/early/2014/01/31/jlr.R045559.full.pdf+html>, doi:10.1194/jlr.R045559.
 - [195] Martin J. Shearer and Paul Newman. Recent trends in the metabolism and cell biology of vitamin k with special reference to vitamin k cycling and mk-4 biosynthesis. *Journal of Lipid Research*, 55(3):345–362, 2014. URL: <http://www.jlr.org/content/55/3/345.abstract>, arXiv:<http://www.jlr.org/content/55/3/345.full.pdf+html>, doi:10.1194/jlr.R045559.
 - [196] Bo Shen, Xiao Yi, Yaoting Sun, Xiaojie Bi, Juping Du, Chao Zhang, Sheng Quan, Fangfei Zhang, Rui Sun, Liujia Qian, Weigang Ge, Wei Liu, Shuang Liang, Hao Chen, Ying Zhang, Jun Li, Jiaqin Xu, Zebao He, Baofu Chen, Jing Wang, Haixi Yan, Yufen Zheng, Donglian Wang, Jiansheng Zhu, Ziqing Kong, Zhouyang Kang, Xiao Liang, Xuan Ding, Guan Ruan, Nan Xiang, Xue Cai, Huanhuan Gao, Lu Li, Sainan Li, Qi Xiao, Tian Lu, Yi Zhu, Huafen Liu, Haixiao Chen, and Tiannan Guo. Proteomic and metabolomic characterization of covid-19 patient sera. *Cell*, 2020. URL: <http://www.sciencedirect.com/science/article/pii/S0092867420306279>, doi:<https://doi.org/10.1016/j.cell.2020.05.032>.
 - [197] Katsumi Shibata and Satoko Toda. Effects of sex hormones on the metabolism of tryptophan to niacin and to serotonin in male rats. *Bioscience, Biotechnology, and Biochemistry*, 61(7):1200–1202, jan 1997. URL: <https://doi.org/10.1271/bbb.61.1200>, doi:10.1271/bbb.61.1200.
 - [198] Takehiko Shibata, David M. Habel, Ana L. Coelho, Steven L. Kunkel, Nicholas W. Lukacs, and Cory M. Hogaboam. Axl receptor blockade ameliorates pulmonary pathology resulting from primary viral infection and viral exacerbation of asthma. *The Journal of Immunology*, 192(8):3569–3581, 2014. URL: <https://www.jimmunol.org/content/192/8/3569>, arXiv:<https://www.jimmunol.org/content/192/8/3569.full.pdf>, doi:10.4049/jimmunol.1302766.
 - [199] DC Simes, CSB Viegas, N Arajo, and C Marreiros. Vitamin k as a diet supplement with impact in human health: Current evidence in age-related diseases. *Nutrients*, 12(1), 2020. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC7019739/>.
 - [200] AK Singal, C Kormos-Hallberg, C Lee, VM Sadagoparamanujam, JJ Grady, DH Freeman, and KE Anderson. Low-dose hydroxychloroquine is as effective as phlebotomy in treatment of patients with porphyria cutanea tarda. *of the American Gastroenterological Association*, 10(12):1402–9, 2012. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3501544/>.
 - [201] Rani H. Singh, Fran Rohr, Dianne Frazier, Amy Cunningham, Shideh Mofidi, Beth Ogata, Patricia L. Splett, Kathryn Moseley, Kathleen Huntington, Phyllis B. Acosta, Jerry Vockley, and Sandra C. Van Calcar. Recommendations for the nutrition management of phenylalanine hydroxylase deficiency. *Genetics in Medicine*, 16(2):121–131, jan 2014. URL: <https://doi.org/10.1038/gim.2013.179>, doi:10.1038/gim.2013.179.
 - [202] Ming Song, Miriam Vos, and Craig McClain. Copper-fructose interactions: A novel mechanism in the pathogenesis of NAFLD. *Nutrients*, 10(11):1815, nov 2018. URL: <https://doi.org/10.3390/nu10111815>, doi:10.3390/nu10111815.
 - [203] T Sugino, S Aoyagi, T Shirai, Y Kajimoto, and O Kajimoto. Effects of citric acid and l-carnitine on physical fatigue. *Journal of Clinical Biochemistry and Nutrition*, 41(3):224–30, 2007. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2243251/>.
 - [204] Kaiyuan Sun, Jenny Chen, and Cécile Viboud. Early epidemiological analysis of the coronavirus disease 2019 outbreak based on crowdsourced data: a population-level observational study. *The Lancet Digital Health*, 2(4):e201–e208, apr 2020. URL: [https://doi.org/10.1016/s2589-7500\(2029\)30026-1](https://doi.org/10.1016/s2589-7500(2029)30026-1), doi:10.1016/s2589-7500(20)30026-1.
 - [205] M Taura, E Kudo, R Kariya, H Goto, K Matsuda, S Hattori, K Vaeteewoottacharn, F McDonald, MA Suico, T Shuto, H Kai, and S Okada. Commd1/murrl reinforces hiv-1 latent infection through ib- stabilization. *Journal of Virology*, 89(5):2643–58, 2015. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4325709/>.
 - [206] Alicia A. Taylor, Joyce S. Tsuji, Michael R. Garry, Margaret E. McArdle, William L. Goodfellow, William J. Adams, and Charles A. Menzie. Critical review of exposure and effects: Implications for setting regulatory health criteria for ingested copper. *Environmental Management*, 65(1):131–159, dec 2019. URL: <https://doi.org/10.1007/s00267-019-01234-y>, doi:10.1007/s00267-019-01234-y.
 - [207] KF Trefz, AC Muntau, KM Kohlscheen, J Altevers, C Jacob, S Braun, W Greiner, A Jha, M Jain, I Alvarez, P Lane, C Schrder, and F Rutsch. Clinical burden of illness in patients with phenylketonuria (pku) and associated. *Orphanet Journal of Rare Diseases*, 14, 2019. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC6647060/>.
 - [208] Masashi Uchida, Yuka Sakaguchi, and Yohei Miyamoto. General toxicity of a vitamin k1 2,3-epoxide reductase (vkor) inhibitor, 3-acetyl-5-methyltetronic acid, in rats. *Fundamental Toxicological Sciences*, 4:285–293, 12 2017. URL: https://www.researchgate.net/publication/322059583_General_toxicity_of_a_vitamin_K1_23-epoxide_reductase_VKOR_inhibitor_3-acetyl-5-methyltetronic_acid_in_rats, doi:10.2131/fts.4.285.
 - [209] C. van Damme, E. Berlingin, S. Saussez, and O. Accaputo. Acute urticaria with pyrexia as the first manifestations of a covid-19 infection. *Journal of the European Academy of Dermatology and Venereology*, n/a(n/a). URL: <https://onlinelibrary.wiley.com/doi/abs/10.1111/jdv.16523>, arXiv:<https://onlinelibrary.wiley.com/doi/pdf/10.1111/jdv.16523>, doi:10.1111/jdv.16523.
 - [210] Annemieke T. van der Goot and Ellen A.A. Nollen. Tryptophan metabolism: entering the field of aging and age-related pathologies. *Trends in Molecular Medicine*, 19(6):336–344, jun 2013. URL: <https://doi.org/10.1016/j.molmed.2013>.

- 02.007, doi:10.1016/j.molmed.2013.02.007.
- [211] Annemieke T. van der Goot, Wentao Zhu, Rafael P. Vázquez-Manrique, Renée I. Seinstra, Katja Dettmer, Helen Michels, Francesca Farina, Jasper Krijnen, Ronald Melki, Rogier C. Buijsman, Mariana Ruiz Silva, Karen L. Thijssen, Ido P. Kema, Christian Neri, Peter J. Oefner, and Ellen A. A. Nollen. Delaying aging and the aging-associated decline in protein homeostasis by inhibition of tryptophan degradation. *Proceedings of the National Academy of Sciences*, 109(37):14912–14917, 2012. URL: <https://www.pnas.org/content/109/37/14912>, arXiv:<https://www.pnas.org/content/109/37/14912.full.pdf>, doi:10.1073/pnas.1203083109.
 - [212] Lucio Verdoni, Angelo Mazza, Annalisa Gervasoni, Laura Martelli, Maurizio Ruggeri, Matteo Ciuffreda, Ezio Bonanomi, and Lorenzo D’Antiga. An outbreak of severe kawasaki-like disease at the italian epicentre of the SARS-CoV-2 epidemic: an observational cohort study. *The Lancet*, may 2020. URL: [https://doi.org/10.1016/s0140-6736\(20\)31103-x](https://doi.org/10.1016/s0140-6736(20)31103-x), doi:10.1016/s0140-6736(20)31103-x.
 - [213] M Vinciguerra and E Greco. Sars-cov-2 and black population: Ace2 as shield or blade? *Infection, Genetics and Evolution*, 20200513. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC7219352/>.
 - [214] Yulin Wan, Sirui Yan, Ying Zhang, Shanshan An, Kaiqiao Yang, Huan Xu, Tao Gan, Weiguo Cao, and Qiuyun Liu. The pneumonia outbreak: High isoleucine and high valine plus glycine contents are features of the proteins of COVID-19 virus. mar 2020. URL: <https://doi.org/10.20944/preprints202002.0289.v4>, doi:10.20944/preprints202002.0289.v4.
 - [215] J Wang, Q Peng, Q Lin, C Childress, D Carey, and W Yang. Calcium activates nedd4 e3 ubiquitin ligases by releasing the c2 domain-mediated auto-inhibition*. *The Journal of Biological Chemistry*, 285(16):12279–88, 2010. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2852967/>.
 - [216] L Wang, W He, X Yu, D Hu, M Bao, H Liu, J Zhou, and H Jiang. Coronavirus disease 2019 in elderly patients: Characteristics and prognostic factors based on 4-week follow-up. *The Journal of Infection*, 80(6):639–45, 2020. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC7118526/>.
 - [217] S Wang, F Guo, K Liu, H Wang, S Rao, P Yang, and C Jiang. Endocytosis of the receptor-binding domain of sars-cov spike protein together with virus receptor ace2. *Virus Research*, 136(1):8–15, 2008. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC7114441/>.
 - [218] Weibin Wang, Xingyun Su, Yongfeng Ding, Weina Fan, Junwei Su, Zhendong Chen, Hong Zhao, Kaijin Xu, Qin Ni, Xiaowei Xu, Yunqing Qiu, and Lisong Teng. Thyroid function abnormalities in covid-19 patients. *medRxiv*, 2020. URL: <https://www.medrxiv.org/content/early/2020/06/16/2020.06.15.20130807>, arXiv:<https://www.medrxiv.org/content/early/2020/06/16/2020.06.15.20130807.full.pdf>, doi:10.1101/2020.06.15.20130807.
 - [219] Katherine Lisa Whitcroft and Thomas Hummel. Olfactory dysfunction in COVID-19. *JAMA*, 323(24):2512, jun 2020. URL: <https://doi.org/10.1001/jama.2020.8391>, doi:10.1001/jama.2020.8391.
 - [220] Wim A. Wuyts, Carlo Agostini, Katerina M. Antoniou, Demosthenes Bouros, Rachel C. Chambers, Vincent Cottin, Jim J. Egan, Bart N. Lambrecht, Rik Lories, Helen Parfrey, Antje Prasse, Carlos Robalo-Cordeiro, Eric Verbeken, Johny A. Verschakelen, Athol U. Wells, and Geert M. Verleden. The pathogenesis of pulmonary fibrosis: a moving target. *European Respiratory Journal*, 41(5):1207–1218, 2013. URL: <https://erj.ersjournals.com/content/41/5/1207>, arXiv:<https://erj.ersjournals.com/content/41/5/1207.full.pdf>, doi:10.1183/09031936.00073012.
 - [221] Chang Xia, Zachary Braunstein, Amelia C. Toomey, Jixin Zhong, and Xiaoquan Rao. S100 proteins as an important regulator of macrophage inflammation. *Frontiers in Immunology*, 8, jan 2018. URL: <https://doi.org/10.3389/fimmu.2017.01908>, doi:10.3389/fimmu.2017.01908.
 - [222] Yan xia Yu, Yi pei Li, Feng Gao, Qing song Hu, Yan Zhang, Dong Chen, and Guang hui Wang. Vitamin k2 suppresses rotenone-induced microglial activation in vitro. *Acta Pharmacologica Sinica*, 37(9):1178–1189, aug 2016. URL: <https://doi.org/10.1038/aps.2016.68>, doi:10.1038/aps.2016.68.
 - [223] Y Xiong, MF Fru, Y Yu, JP Montani, XF Ming, and Z Yang. Long term exposure to l-arginine accelerates endothelial cell senescence through arginase-ii and s6k1 signaling. *Aging (Albany NY)*, 6(5):369–79, 2014. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4069264/>.
 - [224] Zhe-rong Xu, Zhong-ju Tan, Qin Zhang, Qi-feng Gui, and Yun-mei Yang. Clinical effectiveness of protein and amino acid supplementation on building muscle mass in elderly people: A meta-analysis. *PLOS ONE*, 9(9):1–10, 09 2014. URL: <https://doi.org/10.1371/journal.pone.0109141>, doi:10.1371/journal.pone.0109141.
 - [225] J Xue, A Moyer, B Peng, J Wu, BN Hannafon, and WQ Ding. Chloroquine is a zinc ionophore. *PLoS ONE*, 9(10), 2014. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4182877/>.
 - [226] Ai-Ping Yang, Hui-Ming Li, Wen-Qiang Tao, Xue-Jing Yang, Min Wang, Wen-Juan Yang, and Jian-Ping Liu. Infection with SARS-CoV-2 causes abnormal laboratory results of multiple organs in patients. *Aging*, jun 2020. URL: <https://doi.org/10.18632/aging.103255>, doi:10.18632/aging.103255.
 - [227] Nan Yang, Julian A. Tanner, Bo-Jian Zheng, Rory M. Watt, Ming-Liang He, Lin-Yu Lu, Jie-Qing Jiang, Ka-To Shum, Yong-Ping Lin, Kin-Ling Wong, Marie C. M. Lin, Hsiang-Fu Kung, Hongzhe Sun, and Jian-Dong Huang. Bismuth complexes inhibit the SARS coronavirus. *Angewandte Chemie International Edition*, 46(34):6464–6468, aug 2007. URL: <https://doi.org/10.1002/anie.200701021>, doi:10.1002/anie.200701021.
 - [228] SJ Yang, CL Keen, L Lanoue, RB Rucker, and JY Uriu-Adams. Low nitric oxide: a key factor underlying copper deficiency teratogenicity. *Free radical biology & medicine*, 43(12):1639–48, 2007. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2289431/>.
 - [229] Wai-Ming Yau, William C. Wimley, Klaus Gawrisch, and Stephen H. White. The preference of tryptophan for membrane interfaces†. *Biochemistry*, 37(42):14713–14718, oct 1998. URL: <https://doi.org/10.1021/bi980809c>, doi:10.1021/bi980809c.
 - [230] Maimaiti Yisireyili, Yasuhiro Uchida, Koji Yamamoto, Takayuki Nakayama, Xian Wu Cheng, Tadashi Matsushita, Shigeo

- Nakamura, Toyoaki Murohara, and Kyosuke Takeshita. Angiotensin receptor blocker irbesartan reduces stress-induced intestinal inflammation via AT1a signaling and ACE2-dependent mechanism in mice. *Brain, Behavior, and Immunity*, 69:167–179, mar 2018. URL: <https://doi.org/10.1016/j.bbi.2017.11.010>, doi:10.1016/j.bbi.2017.11.010.
- [231] Pierre Yot, Monique Pinck, Anne-Lise Haenni, Henri M. Duranton, and François Chapeville. Valine-specific trna-like structure in turnip yellow mosaic virus rna. *Proceedings of the National Academy of Sciences*, 67(3):1345–1352, 1970. URL: <https://www.pnas.org/content/67/3/1345>, arXiv:<https://www.pnas.org/content/67/3/1345.full.pdf>, doi:10.1073/pnas.67.3.1345.
- [232] JM Zaborske, J Narasimhan, L Jiang, SA Wek, KA Dittmar, F Freimoser, T Pan, and RC Wek. Genome-wide analysis of trna charging and activation of the eif2 kinase gcn2p*. *The Journal of Biological Chemistry*, 284(37):25254–67, 2009. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2757228/>.
- [233] R Zangerle, K Kurz, G Neurauter, M Kitchen, M Sarcletti, and D Fuchs. Increased blood phenylalanine to tyrosine ratio in hiv-1 infection and correction following effective antiretroviral therapy. *Brain, behavior, and immunity*, 24(3):403–8, 2010. URL: <http://www.ncbi.nlm.nih.gov/pubmed/articles/19925861/>.
- [234] Lei Zhang and Yunhui Liu. Potential interventions for novel coronavirus in china: A systematic review. *Journal of Medical Virology*, 92(5):479–490, 2020. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1002/jmv.25707>, arXiv:<https://onlinelibrary.wiley.com/doi/pdf/10.1002/jmv.25707>, doi:10.1002/jmv.25707.
- [235] Wei Zhang, Youshu Yuan, Zhengqiao Yang, Jianxia Fu, Yun Zhang, Ming Ma, Weidong Wu, and Hourong Zhou. Leukopenia of asymptomatic COVID-19 infections under 18 years old in recovery stage. may 2020. URL: <https://doi.org/10.1101/2020.05.21.20074682>, doi:10.1101/2020.05.21.20074682.
- [236] Hangyu Zhou, Guangchuan Wang, Xiaoyu Wang, Zhiyong Song, and Ruikang Tang. Mineralized state of the avian influenza virus in the environment. *Angewandte Chemie International Edition*, 56(42):12908–12912, aug 2017. URL: <https://doi.org/10.1002/anie.201705769>, doi:10.1002/anie.201705769.
- [237] Zhuo Zhou, Lili Ren, Li Zhang, Jiaxin Zhong, Yan Xiao, Zhilong Jia, Li Guo, Jing Yang, Chun Wang, Shuai Jiang, Donghong Yang, Guoliang Zhang, Hongru Li, Fuhui Chen, Yu Xu, Mingwei Chen, Zhancheng Gao, Jian Yang, Jie Dong, Bo Liu, Xiannian Zhang, Weidong Wang, Kunlun He, Qi Jin, Mingkun Li, and Jianwei Wang. Heightened innate immune responses in the respiratory tract of COVID-19 patients. *Cell Host & Microbe*, may 2020. URL: <https://doi.org/10.1016/j.chom.2020.04.017>, doi:10.1016/j.chom.2020.04.017.
- [238] Carly G.K. Ziegler, Samuel J. Allon, Sarah K. Nyquist, Ian M. Mbano, Vincent N. Miao, Constantine N. Tzouanas, Yuming Cao, Ashraf S. Yousif, Julia Bals, Blake M. Hauser, Jared Feldman, Christoph Muus, Marc H. Wadsworth, Samuel W. Kazer, Travis K. Hughes, Benjamin Doran, G. James Gatter, Marko Vukovic, Faith Taliaferro, Benjamin E. Mead, Zhiru Guo, Jennifer P. Wang, Delphine Gras, Magali Plaisant, Meshal Ansari, Ilias Angelidis, Heiko Adler, Jennifer M.S. Sucre, Chase J. Taylor, Brian Lin, Avinash Waghay, Vanessa Mitsialis, Daniel F. Dwyer, Kathleen M. Buchheit, Joshua A. Boyce, Nora A. Barrett, Tanya M. Laidlaw, Shaina L. Carroll, Lucrezia Colonna, Victor Tkachev, Christopher W. Peterson, Alison Yu, Hengqi Betty Zheng, Hannah P. Gideon, Caylin G. Winchell, Philana Ling Lin, Colin D. Bingle, Scott B. Snapper, Jonathan A. Kropski, Fabian J. Theis, Herbert B. Schiller, Laure-Emmanuelle Zaragosi, Pascal Barbry, Alasdair Leslie, Hans-Peter Kiem, JoAnne L. Flynn, Sarah M. Fortune, Bonnie Berger, Robert W. Finberg, Leslie S. Kean, Manuel Garber, Aaron G. Schmidt, Daniel Lingwood, Alex K. Shalek, Jose Ordoñas-Montanes, Nicholas Banovich, Pascal Barbry, Alvis Brazma, Tushar Desai, Thu Elizabeth Duong, Oliver Eickelberg, Christine Falk, Michael Farzan, Ian Glass, Muzlifah Haniffa, Peter Horvath, Deborah Hung, Naftali Kaminski, Mark Krasnow, Jonathan A. Kropski, Malte Kuhnemund, Robert Lafyatis, Haeock Lee, Sylvie Leroy, Sten Linnarson, Joakim Lundberg, Kerstin Meyer, Alexander Misharin, Martijn Nawijn, Marko Z. Nikolic, Jose Ordoñas-Montanes, Dana Pe'er, Joseph Powell, Stephen Quake, Jay Rajagopal, Purushothama Rao Tata, Emma L. Rawlins, Aviv Regev, Paul A. Reyfman, Mauricio Rojas, Orit Rosen, Kourosh Saeb-Parsy, Christos Samakovlis, Herbert Schiller, Joachim L. Schultze, Max A. Seibold, Alex K. Shalek, Douglas Shepherd, Jason Spence, Avrum Spira, Xin Sun, Sarah Teichmann, Fabian Theis, Alexander Tsankov, Maarten van den Berge, Michael von Papen, Jeffrey Whitsett, Ramnik Xavier, Yan Xu, Laure-Emmanuelle Zaragosi, and Kun Zhang. SARS-CoV-2 receptor ACE2 is an interferon-stimulated gene in human airway epithelial cells and is detected in specific cell subsets across tissues. *Cell*, apr 2020. URL: <https://doi.org/10.1016/j.cell.2020.04.035>, doi:10.1016/j.cell.2020.04.035.
- [239] A Zinkeviciene, D Kainov, I Girkontaite, E Lastauskiene, V Kvedariene, Y Fu, S Anders, and V Velagapudi. Activation of tryptophan and phenylalanine catabolism in the remission phase of allergic contact dermatitis: A pilot study. *International archives of allergy and immunology*, 170(4):262–268, 2016. URL: <http://www.ncbi.nlm.nih.gov/pubmed/articles/27771694/>.
- [240] AZ Zucoloto and CN Jenne. Platelet-neutrophil interplay: Insights into neutrophil extracellular trap (net)-driven coagulation in infection. *Frontiers in Cardiovascular Medicine*, 6, 2019. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC6595231/>.
- [241] Yu Zuo, Melanie Zuo, Srilakshmi Yalavarthi, Kelsey Gockman, Jacqueline A. Madison, Hui Shi, Jason S. Knight, and Yogendra Kanthi. Neutrophil extracellular traps and thrombosis in covid-19. *medRxiv*, 2020. URL: <https://www.medrxiv.org/content/early/2020/05/13/2020.04.30.20086736>, arXiv:<https://www.medrxiv.org/content/early/2020/05/13/2020.04.30.20086736.full.pdf>, doi:10.1101/2020.04.30.20086736.

Acknowledgments

1. Pubmed eutils facilities and the basic research it provides.
2. Free software including Linux, R, LaTeX etc.
3. Thanks everyone who contributed incidental support.

Appendix A: Statement of conflicts

No specific funding was used in this effort as the original goal was to optimize feeding for dogs with various conditions. There are no relationships with others that could create a conflict of interest. I would like to develop these ideas further and have obvious bias towards making them appear successful. I have no interest in the makers of any of the products named in this work.

Appendix B: About the author

This work was performed at a dog rescue run by Barbara Cade and housed in rural Georgia. The author of this report ,Mike Marchywka, has a background in electrical engineering (BSEE 1984, MSEE 1985) and has done extensive research using free online literature sources. I hope to find additional people interested in critically examining the results and verify that they can be reproduced effectively to treat other dogs.

Appendix C: Symbols, abbreviations and colloquialisms

TERM	definition and meaning
AFIB	Atrial FIBrillation
BH4	Tetrahydrobiopterin
COPD	Chronic obstructive pulmonary disease
CMV	Cytomegalovirus
DIC	Disseminated Intravascular Coagulation
EET	Eosinophil Extracellular Trap
ERGIC	Endoplasmic Reticulum - Golgi intermediate compartment
IBD	Inflammatory Bowel Disease
IDO	Indoleamine Dioxygenase
NET	Neutrophil Extracellular Trap
NAFLD	Non Alcoholic Fatty Liver Disease
NOAEL	No Observed Adverse Event Level
PAH	Phenylalanine Hydroxylase
RDA	Recommended Daily Allowance
RDI	Recommended Daily Intake
SVZ	Subventricular zone
VKDP	Vitamin K dependent proteins
VKOR	Vitamin K oxidoreductase

Appendix D: General caveats and disclaimer

This document was created in the hope it will be interesting to someone including me by providing information about some topic that may include personal experience or a literature review or description of a speculative theory or idea. There is no assurance that the content of this work will be useful for any particular purpose.

All statements in this document were true to the best of my knowledge at the time they were made and every attempt is made to assure they are not misleading or confusing. However, information provided by others and observations that can be manipulated by unknown causes may be misleading. Any use of this information should be preceded by validation including replication where feasible. Errors may enter into the final work at every step from conception and research to final editing.

Documents labelled "NOTES" or "not public" contain substantial informal or speculative content that may be terse and poorly edited or even sarcastic or profane. Documents labelled as "public" have generally been edited to be more coherent but probably have not been reviewed or proof read.

Generally non-public documents are labelled as such to avoid confusion and embarrassment and should be read with that understanding. ‘

Appendix E: Citing this as a tech report or white paper

Note: This is mostly manually entered and not assured to be error free.
This is tech report MJM-2020-002-0.10 .

Version	Date	Comments
0.01	2020-05-11	Create from empty.tex template
0.10	2020-07-01	NOTES version for discussion
-	July 1, 2020	version 0.10 MJM-2020-002-0.10
1.0	20xx-xx-xx	First revision for distribution

Released versions,

Version	Date	
0.10	2020-07-01	https://media-exp1.licdn.com/dms/document/C4E1FAQGHDj69cUckbw/feedshare-document-pdf-analyzed/0?e=15
0.10	2020-07-01	https://www.linkedin.com/posts/marchywka_notes-on-agin

Suggested Bibtex:

```
@TECHREPORT{mmarchywka-MJM-2020-002-0.10 ,
AUTHOR = {M.J. Marchywka},
TITLE = { On the age distribution of SARS-Cov-2 Patients},
NUMBER = {MJM-2020-002-0.10 },
VERSION = { 0.10 July 1, 2020 PUBLIC NOTES },
INSTITUTION = { not institutionalized , independent},
ADDRESS = {306 Charles Cox , Canton GA 30115},
NOTE = {Version 0.10 , may change significantly if less than 1.00 },
DATE = {July 1, 2020},
DAY = {1},
MONTH = {7},
YEAR = {2020},
AUTHOR1EMAIL = {marchywka@hotmail.com},
AUTHOR1ID = {orcid.org/0000-0001-9237-455X},
PAGES = { 33 },
CONTACT = {marchywka@hotmail.com},
FILENAME = {covid19}
}
```

Supporting files. Note that some dates,sizes, and md5's will change as this is rebuilt.

This really needs to include the data analysis code but right now it is auto generated picking up things from prior build in many cases

```
3888 Jul 1 09:05 ./comment.cut 27da0107de2fb32d4ca436e40684f4e1
22473 Jul 1 09:05 ./covid19.aux cd8686fe1e3b383bca8319991357542b
112984 Jul 1 09:05 ./covid19.bbl d803a12b13b4d311cb518e18718ee81f
415456 Jul 1 09:05 ./covid19.bib 76324d0b9566fa165482ac29651c0e4a
2467 Jul 1 09:05 covid19.blg 2980a948f03636af3bfd934aebada34b
324 Jul 1 09:06 ./covid19.bundle_checksums 23a697b22eb0e64a5ec63d4816188ff8
3 Jul 1 09:05 ./covid19.last_page 4fbafd6948b6529caa2b78e476359875
69109 Jul 1 09:05 covid19.log 0e6276a0bfb0038642c5eea42c71cb89
1817 Jul 1 09:05 ./covid19.out 65505d95c2ff929f09c1ecdca1f6a3f3
118109 Jul 1 09:05 ./covid19.tex c9b25026c254802ddf2ee52fbf204e1a
2329 Jul 1 09:05 ./covid19.toc 80089ceea1f2764c87850a2d98dd19dc
12488 Jun 11 16:26 /home/documents/latex/bib/mjm_tr_prep.bib a5b25ac287b05d33ebf88d4dada5e375
7331 Jan 24 2019 /home/documents/latex/pkg/fltpage.sty 73b3a2493ca297ef0d59d6c1b921684b
7434 Oct 21 1999 /home/documents/latex/pkg/lgrind.sty ea74beead1aa2b711ec2669ba60562c3
7162 Nov 13 2015 /home/documents/latex/pkg/mol2chemfig.sty f5a8b1719cee30a4df0739275ac7f5f8a
```


2858 Jun 18 07:15 /home/documents/latex/share/includes/bibtex.txt 9fb628fb3f5269ec5fe11e0ede188418
1050 Jun 30 18:59 /home/documents/latex/share/includes/disclaimer-informal.tex 82
adbad09c56f90ace278c22aaa14b08
424 Sep 2 2019 /home/documents/latex/share/includes/disclaimer-status.tex f374e3907a4aac2ddee51e7269823223
1286 Nov 14 2019 /home/documents/latex/share/includes/mycommands.tex 18011c7f850bd7ab15625df8328e3cf8
2901 Jun 17 15:04 /home/documents/latex/share/includes/myskeletonpackages.tex
fcfcd2e3c8d69d533932edaaa47f53a1
331065 Jul 1 06:24 non_pmc_covid19.bib 04cbd619516722e9c2adaa70db7de746
83988 Jun 30 08:34 pmc_covid19.bib 56dbe74591edaf84f265ec9b5206aa9a
63 Jul 1 06:55 ./releases.tex 161c46fc24fc73868762881b91917ae5
31050 Jul 21 2011 /usr/share/texlive/texmf-dist/bibtex/bst/urlbst/plainurl.bst
ffdaefb09013f5fd4b31e485c13933c1

454799 Jul 1 09:05 covid19.pdf 58d1d0a4fd1e8f0cbdea7b4377e8cc2a