Release Notes 2021-12-22: This is an extract from a longer (thesis length) work (explaining results feeding amino acids etc to older dogs) that is taking forever to organize and I wanted to get something out today. It motivates complexities with interpreting amino acid levels as a basis for picking an intervention but is discussed as a more general problem perceiving paradigms as paradoxes. The equation probably appears somewhere in other literature, I found a quote from the kidney lit expressing the sentiment, but I did not bother to look. It could certainly use some cleanup but seems to make the immediate points. Several peripheral points are not well documented. This is bad scholarship but can be fixed if they become important. To say the least, the paradox/paradigm point is very simplistic and intended for a variety of audiences but not sure if it is well suited for any one in particular.

This is a draft which may not have been fully proofread and certainly not peer reviewed. Read the disclaimers and take them seriously. The reader is assumed familiar with the related literature and controversial issues. For information and thought only not intended for any particular purpose. Caveat Emptor

The Paradox Paradigm: When Could Amino Acid Supplements Be Beneficial?

Mike Marchywka*
306 Charles Cox , Canton GA 30115
(Dated: December 22, 2021)

Amino acids are important components of proteins but also consumed as signalling molecules such as neurotransmitters or oxidized as fuel. They may also be modified non-enzymatically creating amino acid analogs that may act as the parent molecule to some degree. Blood concentrations of specific amino acids or proteins may be obvious clues about possible benefits of supplementation but that need not be the case. This paradox is elaborated by examining various correlations and contrasting them to interventions but then rationalizing with simple general equations and feedback mechanisms. While details are lacking and the model is greatly simplified, overall conservation laws, things like global nitrogen balance, are considered as sanity checks on the concentration-sufficiency assumption. Illustrations with the hypertension literature and recent amino acid profiles in covid-19 patients are considered along with some support from the dementia and kidney disease literature. Regulation patterns may be altruistic, shuttling scare amino acids from muscles to the brain, as has been observed with biotin for example. In this case, blood levels may be fortuitously normal but the 'direction" of flow is an unsustainable compensation perhaps for low glucose due to some other reason. Assays often include "damaged" components in the case of proteins and these may be elevated due to conservation measures but supply may still benefit from supplements. In brief, the paradox is due to mistaking a concentration for a velocity or flux.

1. INTRODUCTION

There is a lot of literature on blood concentrations of amino acids and proteins in health and disease. There is also literature on overall nitrogen balance attempting to add up input and outputs and determine if it is being accumulated or depleted. However, it is not clear how correlations between blood levels relate to disease modifying interventions or cures. Some of the discrepancies between correlations and "obvious" interventions are discussed after the introduction of a general rate equation as a focus for discussion. Hopefully this can point to more useful interventions or at least more informative failures.

2. ANALYSIS

Most molecules in the blood such as amino acids can be described as coming from some source "a" and going to some destination "b" while the latter may include modification without exit from the blood stream. Consider a general rate equation as follows,

^{*}Electronic address: marchywka@hotmail.com; to cite or credit this work, see bibtex in Appendix E

$$\frac{d[x_i]}{dt} = \sum_{i} a_{i,j} - [x_i] \sum_{k} b_{i,k} \tag{1}$$

where $[x_i]$ is the concentration of some molecule such as an amino acid, $a_{i,j}$ is a production rate from source j for x_i and $b_{i,k}$ is a consumption rate constant for some sink k. In the present cases, sources can include the GI tract or muscle and sinks can be organs such as muscle and brain, or the urine. These include components from metabolic transformations such as conversions from one amino acid to another or conversion into neurotransmitters. The source and sink values may be adaptive and vary with other parameters. Usually the thing that matters most is the consumption rate(s), $[x_i]b_{i,k}$, which can be optimized independently via "b" but total consumption and $[x_i]$ is limited by production rate.

In steady state the blood level or concentration is,

$$[x_i] = \frac{\sum_j a_{i,j}}{\sum_k b_{i,k}} \tag{2}$$

which may be important if this level modifies the properties of the blood or the consumers are not adaptive. Otherwise, the consumption rates matter and these may be compromised in times of nutrient shortages or stress requiring "repair" while maintaining some normal concentrations. The time constant related to the circulation time for molecules of x_i is

$$\tau_i = \frac{1}{\sum_k b_{i,k}} \tag{3}$$

and

$$[x_i] = \tau_i \sum_j a_{i,j} \tag{4}$$

suggesting that the same blood level can be obtained with varying production rates and lifetimes (production and consumption velocity vectors or "directions" could be defined as $\overrightarrow{\mathbf{a_i}}$ and $[x_i]\overrightarrow{\mathbf{b_i}}$).

As τ increases, , more may be destined for spontaneous modification and may still function somewhat as the parent molecule especially with proteins.

In the case of multiple deficiencies, amino acid consumption may increase as a compensatory measure raising blood levels but not indicating excess. Effects such as this lead to changes in blood levels that do not relate to sufficiency in obvious ways. In this case, the expectation that deficiency leads to low blood levels, an assumed correlation, may discourage useful interventions rather than encourage useless ones as is often the case. Ideally, the other deficiencies would be fixed but if amio acids have been mobilized for any length of time replacing them is still likely to be beneficial. The discrepancy between correlations and interventions is explored in some limited literature from hypertension, dementia, kidney disease, and covid-19.

2.1. Hypertension

Hypertension literature is an excellent example of this issue. While not exhaustive by any means, this selection appears reasonably representative. Studies of correlations are interpreted to suggest amino acids are not a factor. A 2019 review works on amino acids and hypertension was inconclusive [22] but the literature reviewed mostly explored plasma amino acids and blood pressure. High levels of blood BCAA's were found to be predictive of developing hypertension [7]. BCAA's sound even more suspicious as correlations between known risk factors and blood levels of BCAA's have been observed [8]. Protein or amino acid intake studies however have shown a protective effect of higher consumption of some amino acids inferred from diet [12] or mixed results [32] or in favor of trendy plant based diets [31]. A BCAA intake study suggested higher consumption was beneficial [11] and even beneficial in cirrhosis patients with portal hypertension [16]. Lysine supplementation was shown to improve hypertension in a small group of adults in Ghana [33]. Tryptophan normalized high blood pressure within 2 hours in a group of 14 patients [4]. Showing a consistent benefit from just tryptophan may require good patient selection or a combination of nutrients but none of them need demonstrate low blood concentrations. Tyrosine administration to spontaneously hypertensive rat reduced blood pressure but the effect was blocked by other large neutral amino acids [29] and one small trial on mildly hypertensive people failed to show a benefit [27].

The apparent inconsistency between correlation and intake trials can be resolved with hypothesized causality and the source and sink equations above. For example, muscle uptake can be reduced in starvation even at a fixed plasma level [34] which may allow for more amino acids to be used for gluconeogenesis. (Interestingly, in older people there is some indication muscles increase amino acid export after exercise [3] which would be an altruistic move anticipating brain requirements. As precedent, biotin altruism among SMVT expressors has been observed.) In this case, intervention trials would make sense even lacking associations between low plasma levels and BP.

2.2. Dementia

Dementia appears to be another area where amino acid profiles don't follow glib expectations. One case of amino acid profiling in elderly dementia patients demonstrates increases in serine, arginine, and isoleucine [26] Similarly, increased amino acids in CSF for dementia patients (methionine and alanine) [19] have been observed. In cases where the brain's glucose metabolism is impaired, amino acid oxidation may make up the difference [10] further contributing to loss and probably mobilization from various reserves increasing concentrations in fluids. Interestingly, as a possible confounding or alternative effect, insulin and glucose were shown to increase Trp and Tyr in the brain while decreasing serum levels suggestive of transport into the CNS [30] although the situation is complicated by factors including competing amino acids binding to albumin [2]. While serum tryptophan was shown to correlate with cognitive impairment in a small group of older women [23], it was elevated in cancer patients [14] who probably do not have excess protein supply. If this is compensatory or maladaptive is unknown.

Limited intervention trials however have shown promise. As early as the 1970's, brain neurotransmitters including acetylcholine, were found to be responsive to dietary precursor inputs to varying degrees [5]. Neurodegeneration was corrected in one mouse model with essential amino acids and the authors point out there is epidemiological association with senile dementia [24] suggesting the result is applicable to real disease. An intervention trial with "Amino LP7" on adults 55 and older with other criteria demonstrated benefits in various aspects of functioning [28] but that likely can be improved due to considerations discussed here and in forthcoming work.

Lithium was recently found to suppress tryptophan catabolism [9] although a role for tryptophan depletion in mental disorders did not appear to be considered. Many drugs such as MAO inhibitors ultimately conserve products derived from amino acids and therefore could conserve the amino acids. SSRI's, MAOI's, anti-histamines, etc could all be expected to change fluxes and adjust control mechanisms.

2.3. Kidney Disease

Kidney disease is often considered in terms of modification to protein intake but recommendations vary. In this context, serum albumin illustrates problems with protein assays. Serum albumin as with all proteins has many modified forms both physiological and pathological [15] [25] [21] [1] so trying to measure a "total" amount may be uninformative for nutritional interventions. Longer circulation lifetime which may occur with conservation attempts likely allow for more damage as is thought to occur with fibrinogen [18]. The multiple effects on source and sink have been noted for kidney patients [6],

Despite its sensitivity as a screening and prognostic tool, serum albumin provides limited information about the complex nature of the underlying nutritional problem in the setting of AKI and CKD. The albumin concentration is the net result of its synthesis, breakdown, the volume of distribution, and exchange between intra- and extra-vascular spaces, as well as losses [75]. Besides, it is a negative acute phase reactant, i.e., during acute illness its synthesis is reduced, resulting in low serum levels. Albumin level values should not be interpreted alone, and the appropriate nutritional assessment should also include a thorough physical exam and clinical judgment [76].

While high protein diets with kidney disease are controversial, some consideration of "protein quality" exists [13] and its even possible that specific amino acid deficiencies contribute to kidney disease. In the specific case of kidney, protein and in particular tyrosine are important issues although glutamine supplementation is thought to be harmful [6] which itself has been a surprising or "paradoxical" result even as I had not recommended that [17] but suggested it could be synthesized from the TCA and "waste" nitrogen sources.

It has been said [20],

A deficiency of even a single AA leads to a negative nitrogen balance. In this state, more protein is degraded than is synthesized, so more nitrogen is excreted than is ingested (38). The latter is particularly relevant for CKD patients under low protein diet (LPD), who may be at risk for developing protein-energy wasting (PEW) if not adequately monitored.

In theory large proteins could be catabolized to gain a few necessary amino acids leaving the rest as waste products. In such a case, essential amino acids even in slight surplus, would be beneficial over wasting large amounts of nitrogen. Urinary protein "wasting' could reflect the only means to eliminate crosslinked/modified proteins or simply eliminating excess scavenged proteins (the "haystack" as the "needle' is found).

2.4. covid-19

Various derangements were observed in covid-19 patients [17] but with no clear interpretation. Given the descriptions of hypermetabolism, GI infections, and immune activation, its most reasonable to assume essential amino acids were depleted and elevated blood levels are clues about some compensatory actions not surplus. Interestingly, the much discussed ACE2 is associated with amino acid transport which is largely ignored. My work is explicitly concerned with tyrosine isomers which have significant existing literature but have not been studied in covid-19 or in relation to coagulation status.

2.5. Discussion

These paradoxes, successful interventions that don't seem consistent with blood levels, may be a pattern or actually a paradigm covered by the general rate equation although other issues may cloud correlation studies. Headlines about vitamin D associations with various health outcomes are much more numerous than successful intervention trials. In this case, vitamin D status could correlate with fat soluble nutrient absorption and suggest issues with vitamin K or lipophilic amino acids. With neutrophils, the left shift is known to indicate velocity but those clues may not exist for other quantities. A1c is an integrated damage measure but that may also depend on RBC half life (another dynamical component usually negligible). With amino acids, there is a lot of unexplored territory but once the correlation studies are considered less discouraging when a hypothesis on cause and effect exists, interventions such as those I proposed earlier make a lot more sense. In the case of older people with impaired digestion and intake or protein wasting in the urine, amino acid supplements may make sense especially with any empirical indications of benefit absent randomized controlled clinical trials.

With blood sugar, high blood sugar causes damage as measured with "A1c" but also often represents a failure to consume the sugar. Some damage may be avoided by reducing supply or wasting it, but ultimately there is likely a problem with consuming it that should be fixed.

In reality, many patients have issues with diet or GI tract making protein deficiency a reasonable suspect. In other cases, with protein wasting via the kidneys, even more amino acids may be needed even if protein wasting is thought to be related to excess protein. Tryptophan in particular may be a concern due to its scarcity, physical properties, and immune activated destruction.

While it remains to be determined if results hold up in definitive well controlled clinical trials, limited Intervention trials can be successful contrary to what may be expected based on correlation studies.

3. CONCLUSIONS

In many cases, where negative nitrogen or amino acid balance can be reasonably assumed, the existence of normal or high blood levels should not be given particular significance. Amino acid supplementation may make a lot of sense then in situations where it is remains controversial such as with hypertension, kidney disease, dementia, or in relation to covid-19. Not only do naive correlations prompt futile intervention trials but they may also discourage effective treatment investigations too. Direct intervention and empirical successes need to be seriously considered even when less direct measures conflict. Ultimately well controlled randomized clinical intervention trials with clinical endpoints will have to settle these issues rather than just correlations or associations.

4. SUPPLEMENTAL INFORMATION

4.1. Computer Code

5. BIBLIOGRAPHY

[1] Daria A. Belinskaia, Polina A. Voronina, Vladimir I. Shmurak, Mikhail A. Vovk, Anastasia A. Batalova, Richard O. Jenkins, and Nikolay V. Goncharov. The universal soldier: Enzymatic and non-enzymatic antioxidant functions of serum albumin. *Antioxidants*, 9(10), 2020. URL: https://www.mdpi.com/2076-3921/9/10/966, doi:10.3390/antiox9100966.

- [2] P M Daniel, E R Love, S R Moorhouse, and O E Pratt. The effect of insulin upon the influx of tryptophan into the brain of the rabbit. The Journal of Physiology, 312(1):551-562, 1981. URL: https://onlinelibrary.wiley.com/doi/abs/10.1113/jphysiol.1981.sp013643, arXiv:https://onlinelibrary.wiley.com/doi/pdf/10.1113/jphysiol.1981.sp013643.
- [3] Micah J. Drummond, Christopher S. Fry, Erin L. Glynn, Kyle L. Timmerman, Jared M. Dickinson, Dillon K. Walker, David M. Gundermann, Elena Volpi, and Blake B. Rasmussen. Skeletal muscle amino acid transporter expression is increased in young and older adults following resistance exercise. *Journal of Applied Physiology*, 111(1):135–142, 2011. PMID: 21527663. URL: https://doi.org/10.1152/japplphysiol.01408.2010, arXiv:https://doi.org/10.1152/japplphysiol.01408.2010.
- [4] H. Feltkamp, K. A. Meurer, and E. Godehardt. Tryptophan-induced lowering of blood pressure and changes of serotonin uptake by platelets in patients with essential hypertension. *Klinische Wochenschrift*, 62, 12 1984. URL: http://dx.doi.org/10.1007/bf01782468, doi:10.1007/bf01782468.
- [5] John D. Fernstrom. Effects of the diet on brain neurotransmitters. Metabolism, 26(2):207-223, 1977. URL: https://www.sciencedirect.com/science/article/pii/0026049577900579, doi:https://doi.org/10.1016/0026-0495(77)90057-9.
- [6] Enrico Fiaccadori, Alice Sabatino, Rocco Barazzoni, Juan Jesus Carrero, Adamasco Cupisti, Elisabeth De Waele, Joop Jonckheer, Pierre Singer, and Cristina Cuerda. Espen guideline on clinical nutrition in hospitalized patients with acute or chronic kidney disease. Clinical Nutrition, 40, 4 2021. URL: http://dx.doi.org/10.1016/j.clnu.2021.01.028, doi: 10.1016/j.clnu.2021.01.028.
- [7] Jose L. Flores-Guerrero, Dion Groothof, Margery A. Connelly, James D. Otvos, Stephan J.L. Bakker, and Robin P.F. Dullaart. Concentration of branched-chain amino acids is a strong risk marker for incident hypertension. *Hypertension*, 74(6):1428-1435, 2019. URL: https://www.ahajournals.org/doi/abs/10.1161/HYPERTENSIONAHA.119.13735, arXiv:https://www.ahajournals.org/doi/pdf/10.1161/HYPERTENSIONAHA.119.13735, doi:10.1161/HYPERTENSIONAHA.119.13735.
- [8] Keiko Fukushima, Sei Harada, Ayano Takeuchi, Ayako Kurihara, Miho Iida, Kota Fukai, Kazuyo Kuwabara, Suzuka Kato, Minako Matsumoto, Aya Hirata, Miki Akiyama, Masaru Tomita, Akiyoshi Hirayama, Asako Sato, Chizuru Suzuki, Masahiro Sugimoto, Tomoyoshi Soga, Daisuke Sugiyama, Tomonori Okamura, and Toru Takebayashi. Association between dyslipidemia and plasma levels of branched-chain amino acids in the japanese population without diabetes mellitus. *Journal of Clinical Lipidology*, 13, 11 2019. URL: http://dx.doi.org/10.1016/j.jacl.2019.09.002, doi:10.1016/j.jacl.2019.09.002.
- [9] R Gottert, P Fidzinski, L Kraus, UC Schneider, M Holtkamp, M Endres, K Gertz, and G Kronenberg. Lithium inhibits tryptophan catabolism via the inflammation-induced kynurenine pathway in human microglia. *Glia*, 2021. URL: http://www.ncbi.nlm.nih.gov/pubmed/articles/34862988/.
- [10] Jeddidiah W. D. Griffin and Patrick C. Bradshaw. Amino acid catabolism in alzheimers disease brain: Friend or foe? Oxidative Medicine and Cellular Longevity, page 5472792, 02 2017. URL: https://doi.org/10.1155/2017/5472792, doi:10.1155/2017/5472792.
- [11] Amy Jennings, Alex MacGregor, Tess Pallister, Tim Spector, and Aedn Cassidy. Associations between branched chain amino acid intake and biomarkers of adiposity and cardiometabolic health independent of genetic factors: A twin study. International Journal of Cardiology, 223:992-998, 2016. URL: https://www.sciencedirect.com/science/article/pii/S0167527316320277, doi:https://doi.org/10.1016/j.ijcard.2016.08.307.
- [12] Amy Jennings, Alex MacGregor, Ailsa Welch, Phil Chowienczyk, Tim Spector, and Aedn Cassidy. Amino Acid Intakes Are Inversely Associated with Arterial Stiffness and Central Blood Pressure in Women. The Journal of Nutrition, 145(9):2130– 2138, 07 2015. URL: https://doi.org/10.3945/jn.115.214700, arXiv:https://academic.oup.com/jn/article-pdf/ 145/9/2130/30060530/jn214700.pdf, doi:10.3945/jn.115.214700.
- [13] Gang-Jee Ko, Connie M. Rhee, Kamyar Kalantar-Zadeh, and Shivam Joshi. The effects of high-protein diets on kidney health and longevity. *Journal of the American Society of Nephrology*, 31(8):1667-1679, 2020. URL: https://jasn.asnjournals.org/content/31/8/1667.full.pdf, doi:10.1681/ASN.2020010028.
- [14] Alessandro Laviano, Antonia Cascino, Maurizio Muscaritoli, Francesca Fanfarillo, and Filippo Rossi Fanelli. Tumor-Induced Changes In Host Metabolism: A Possible Role For Free Tryptophan As A Marker Of Neoplastic Disease, pages 363–366. Springer US, Boston, MA, 2003. URL: https://doi.org/10.1007/978-1-4615-0135-0_41, doi: 10.1007/978-1-4615-0135-0_41.
- [15] Magorzata Maciek-Jurczyk, Agnieszka Szkudlarek, Mariola Chudzik, Jadwiga Poycka, and Anna Sukowska. Alteration of human serum albumin binding properties induced by modifications: A review. Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy, 188:675–683, 2018. URL: https://www.sciencedirect.com/science/article/pii/

- S1386142517303906, doi:https://doi.org/10.1016/j.saa.2017.05.023.
- [16] Giulio Marchesini, Giampaolo Bianchi, Manuela Merli, Piero Amodio, Carmine Panella, Carmela Loguercio, Fillipo Rossi Fanelli, and Roberto Abbiati. Nutritional supplementation with branched-chain amino acids in advanced cirrhosis: a double-blind, randomized trial. *Gastroenterology*, 124, 6 2003. URL: http://dx.doi.org/10.1016/s0016-5085(03) 00323-8, doi:10.1016/s0016-5085(03)00323-8.
- [17] Mike J Marchywka. Update to covid-19 age distribution notes. Technical Report MJM-2021-007, not institutionalized, independent, 306 Charles Cox, Canton GA 30115, 8 2021. Version .1, may change signficantly if less than 1.00. URL: https://www.researchgate.net/publication/353946686_Draft_table_comparing_expectations_to_recent_results_with_covid-19.
- [18] M.J. Marchywka. Considering alternative fibrinogen fates in diseased states. Technical Report MJM-2021-006, not institutionalized, independent, 306 Charles Cox, Canton GA 30115, 07 2021. Version 0.50, may change signficantly if less than 1.00. URL: https://www.researchgate.net/publication/353314699_Considering_Alternative_Fibrinogen_Fates_in_Diseased_States.
- [19] Y Mochizuki, M Oishi, M Hara, and T Takasu. Amino acid concentration in dementia of the alzheimer type and multi-infarct dementia. Annals of Clinical & Laboratory Science, 26, 05 1996. URL: http://www.annclinlabsci.org/content/26/3/275.full.pdf.
- [20] Pablo Molina, Eva Gavela, Beln Vizcano, Emma Huarte, and Juan Jess Carrero. Optimizing diet to slow ckd progression. Frontiers in Medicine, 06 2021. URL: https://www.ncbi.nlm.nih.gov/labs/pmc/articles/PMC8267004/, doi:10.3389/fmed.2021.654250.
- [21] Masaki Otagiri and Victor Tuan Giam Chuang. Pharmaceutically important pre- and posttranslational modifications on human serum albumin. *Biological and Pharmaceutical Bulletin*, 32(4):527–534, 2009. URL: https://www.jstage.jst.go.jp/article/bpb/32/4/32_4_527/_article/-char/ja/, doi:10.1248/bpb.32.527.
- [22] Eleonora Poggiogalle, Mario Fontana, Anna Maria Giusti, Alessandro Pinto, Gino Iannucci, Andrea Lenzi, and Lorenzo Maria Donini. Amino acids and hypertension in adults. *Nutrients*, 06 2019. URL: https://pubmed.ncbi.nlm.nih.gov/31252583/, doi:10.3390/nu11071459.
- [23] L. A. Ramos-Chvez, G. Roldn-Roldn, B. Garca-Jurez, D. Gonzlez-Esquivel, G. Prez de la Cruz, B. Pineda, D. Ramrez-Ortega, I. Garca Muoz, B. Jimnez Herrera, C. Ros, S. Gmez-Manzo, J. Marcial-Quino, L. Snchez Chapul, P. Carrillo Mora, and V. Prez de la Cruz. Low serum tryptophan levels as an indicator of global cognitive performance in nondemented women over 50 years of age. Oxidative Medicine and Cellular Longevity, 2018, 11 2018. URL: http://dx.doi.org/10.1155/2018/8604718, doi:10.1155/2018/8604718.
- [24] Hideaki Sato, Yuhei Takado, Sakiko Toyoda, Masako Tsukamoto-Yasui, Keiichiro Minatohara, Hiroyuki Takuwa, Takuya Urushihata, Manami Takahashi, Masafumi Shimojo, Maiko Ono, Jun Maeda, Asumi Orihara, Naruhiko Sahara, Ichio Aoki, Sachise Karakawa, Muneki Isokawa, Noriko Kawasaki, Mika Kawasaki, Satoko Ueno, Mayuka Kanda, Mai Nishimura, Katsuya Suzuki, Akira Mitsui, Kenji Nagao, Akihiko Kitamura, and Makoto Higuchi. Neurodegenerative processes accelerated by protein malnutrition and decelerated by essential amino acids in a tauopathy mouse model. *Science Advances*, 7, 10 2021. URL: http://dx.doi.org/10.1126/sciadv.abd5046, doi:10.1126/sciadv.abd5046.
- [25] Andre M.N. Silva and Robert C. Hider. Influence of non-enzymatic post-translation modifications on the ability of human serum albumin to bind iron: Implications for non-transferrin-bound iron speciation. *Biochimica et Biophysica Acta (BBA)* Proteins and Proteomics, 1794(10):1449-1458, 2009. URL: https://www.sciencedirect.com/science/article/pii/S1570963909001319, doi:https://doi.org/10.1016/j.bbapap.2009.06.003.
- [26] Edyta Socha, Piotr Koliski, Marcin Koba, Katarzyna Mdra-Gackowska, Kornelia Kdziora-Kornatowska, Marcin Gackowski, and Emilia Daghir-Wojtkowiak. Amino acid levels as potential biomarker of elderly patients with dementia. *Brain Sciences*, 10(12), 2020. URL: https://www.mdpi.com/2076-3425/10/12/914, doi:10.3390/brainsci10120914.
- [27] M J Sole, C R Benedict, M G Myers, F H Leenen, and G H Anderson. Chronic dietary tyrosine supplements do not affect mild essential hypertension. *Hypertension*, 7(4):593-596, 1985. URL: https://www.ahajournals.org/doi/abs/10.1161/01.HYP.7.4.593, arXiv:https://www.ahajournals.org/doi/pdf/10.1161/01.HYP.7.4.593, doi:10.1161/01.HYP.7.4.593
- [28] Hiroyuki Suzuki, Daichi Yamashiro, Susumu Ogawa, Momoko Kobayashi, Daisuke Cho, Ai Iizuka, Masako Tsukamoto-Yasui, Michihiro Takada, Muneki Isokawa, Kenji Nagao, and Yoshinori Fujiwara. Intake of seven essential amino acids improves cognitive function and psychological and social function in middle-aged and older adults: A double-blind, randomized, placebo-controlled trial. Frontiers in Nutrition, 7:240, 2020. URL: https://www.frontiersin.org/article/10.3389/fnut.2020.586166, doi:10.3389/fnut.2020.586166.
- [29] A. F. Sved, J. D. Fernstrom, and R. J. Wurtman. Tyrosine administration reduces blood pressure and enhances brain norepinephrine release in spontaneously hypertensive rats. *Proceedings of the National Academy of Sciences*, 76, 07 1979. URL: http://dx.doi.org/10.1073/pnas.76.7.3511, doi:10.1073/pnas.76.7.3511.
- [30] A. Tagliamonte, M.G. DeMontis, M. Olianas, P.L. Onali, and G.L. Gessa. Possible role of insulin in the transport of tyrosine and tryptophan from blood to brain. *Pharmacological Research Communications*, 7(6):493–499, 1975. URL: https://www.sciencedirect.com/science/article/pii/0031698975900107, doi:https://doi.org/10.1016/0031-6989(75)90010-7.
- [31] Farshad Teymoori, Golaleh Asghari, Parvin Mirmiran, and Fereidoun Azizi. Dietary amino acids and incidence of hypertension: A principle component analysis approach. *Scientific Reports*, 7, 12 2017. URL: https://www.nature.com/articles/s41598-017-17047-0, doi:10.1038/s41598-017-17047-0.
- [32] Katherine R. Tuttle, Joan E. Milton, Diane P. Packard, Lynn A. Shuler, and Robert A. Short. Dietary amino acids and blood pressure: A cohort study of patients with cardiovascular disease. *American Journal of Kidney Diseases*, 59, 6 2012. URL: http://dx.doi.org/10.1053/j.ajkd.2011.12.026, doi:10.1053/j.ajkd.2011.12.026.

- [33] Frederick Vuvor, Husein Mohammed, Thomas Ndanu, and Obed Harrison. Effect of lysine supplementation on hypertensive men and women in selected peri-urban community in ghana. *BMC Nutrition*, 3, 07 2017. URL: http://dx.doi.org/10.1186/s40795-017-0187-6, doi:10.1186/s40795-017-0187-6.
- [34] B W Warner, R P 3rd Hummel, P O Hasselgren, J H James, and J E Fischer. Inhibited amino acid uptake in skeletal muscle during starvation. *JPEN. Journal of parenteral and enteral nutrition*, pages 344–8, Jul-Aug 1989. URL: https://pubmed.ncbi.nlm.nih.gov/2778939/, doi:10.1177/0148607189013004344.

Acknowledgments

- 1. Pubmed eutils facilities and the basic research it provides.
- 2. Free software including Linux, R, LaTex etc.
- 3. Thanks everyone who contributed incidental support.

Appendix A: Statement of Conflicts

No specific funding was used in this effort and there are no relationships with others that could create a conflict of interest. I would like to develop these ideas further and have obvious bias towards making them appear successful. Barbara Cade, the dog owner, has worked in the pet food industry but this does not likely create a conflict. We have no interest in the makers of any of the products named in this work.

Appendix B: About the Authors and Facility

This work was performed at a dog rescue run by Barbara Cade and housed in rural Georgia. The author of this report ,Mike Marchywka, has a background in electrical engineering and has done extensive research using free online literature sources. I hope to find additional people interested in critically examining the results and verify that they can be reproduced effectively to treat other dogs.

Appendix C: Symbols, Abbreviations and Colloquialisms

TERM definition and meaning		
	AA	amino acid
	AKI	acute kidney injury
	BCAA	branched chain amino acid
	CKD	chronic kidney disease
	CNS	central nervous system
	CSF	cerebrospinal fluid
	GI	gastrointestinal tract
	MAO	monoamine oxidase
	RBC	red blood cell
	SMVT	sodium dependent multivitamin transporter

tri-carboxylic acid cycle

uniform resource locator

TCA

URL

Appendix D: General caveats and disclaimer

This document was created in the hope it will be interesting to someone including me by providing information about some topic that may include personal experience or a literature review or description of a speculative theory or idea. There is no assurance that the content of this work will be useful for any paricular purpose.

All statements in this document were true to the best of my knowledge at the time they were made and every attempt is made to assure they are not misleading or confusing. However, information provided by others and observations that can be manipulated by unknown causes ("gaslighting") may be misleading. Any use of this information should be preceded by validation including replication where feasible. Errors may enter into the final work at every step from conception and research to final editing.

Documents labelled "NOTES" or "not public" contain substantial informal or speculative content that may be terse and poorly edited or even sarcastic or profane. Documents labelled as "public" have generally been edited to be more coherent but probably have not been reviewed or proof read.

Generally non-public documents are labelled as such to avoid confusion and embarassment and should be read with that understanding.

Appendix E: Citing this as a tech report or white paper

Note: This is mostly manually entered and not assured to be error free. This is tech report MJM-2021-015.

Version	Date	Comments
0.01	2021-12-20	Create from empty.tex template
0.50	2021-12-22	Release for comment
-	December 22, 2021	version 0.50 MJM-2021-015
1.0	20xx-xx-xx	First revision for distribution

Released versions, build script needs to include empty releases.tex

```
VersionDateURL0.502021-12-22https://www.researchgate.net/publication/357250784_The_Paradox_Paradigm_When_Could_Amino_Acid_Supplem0.502021-12-22https://www.academia.edu/s/3b3273469a0.502021-12-22https://www.linkedin.com/posts/marchywka_amino-acid-blood-levels-versus-production-activity-68794158
```

```
@techreport{marchywka-MJM-2021-015-0.50,
filename = "aapp",
run-date ="December 22, 2021",
title ="The Paradox Paradigm: When Could Amino Acid Supplements Be Beneficial?",
author ="Mike J Marchywka",
type ="techreport",
name = "marchywka-MJM-2021-015-0.50",
number = "MJM-2021-015",
version = "0.50",
institution ="not institutionalized, independent"
address =" 306 Charles Cox, Canton GA 30115",
date ="December 22, 2021",
startdate = "2021-12-20",
day = "22",
month = "12"
year ="2021"
author1email = "marchywka@hotmail.com",
contact ="marchywka@hotmail.com"
author1id = "orcid.org/0000-0001-9237-455X",
pages =" 10"
```

Supporting files. Note that some dates, sizes, and md5's will change as this is rebuilt.

This really needs to include the data analysis code but right now it is auto generated picking up things from prior build in many cases

```
6496 Dec 22 08:16 aapp.aux d2ddde6c1ad386460e1bf3331c40a642
18158 Dec 22 08:16 aapp.bbl 9c3b9e94117da15581085dd5e1ccebf1
803 Dec 22 08:16 aapp.bib a4513cf6c43b9c230937e5dc809cde83
1522 Dec 22 08:16 aapp.blg c27630feb6e6848df57856d7656b897c
0 Dec 22 08:16 aapp.bundle_checksums d41d8cd98f00b204e9800998ecf8427e
29411 Dec 22 08:16 aapp.fls f21999091c9481928486c7994fc17b1b
3 Dec 22 08:16 aapp.last_page 31d30eea8d0968d6458e0ad0027c9f80
57726 Dec 22 08:16 aapp.log 17f8dbd0fe4b299eb74ab579936141bc
1055 Dec 22 08:16 aapp.out ddaf88e5d1756c05434d22c5bf742bf6
243728 Dec 22 08:16 aapp.pdf cdf72e90cfbfe6809435bd9978d809d5
31935 Dec 22 08:14 aapp.tex bb3efd47fdec01758f8b2e4f6f29c167
3953 Dec 22 08:16 comment.cut 49a6350413f114d877e997ae0d93535f
1101 Aug 15 2019 /etc/texmf/web2c/texmf.cnf af7716885e081ab43982cab7b4672c1a
23946 Dec 20 15:57 /home/documents/latex/bib/mjm_tr.bib 5d23d75d60273fbd1c30bd661aaa168b
21058 Dec 5 10:59 /home/documents/latex/bib/releases.bib 9b39a44e8bf390c1a65babfe00e5884c
7331 Jan 24 2019 /home/documents/latex/pkg/fltpage.sty 73b3a2493ca297ef0d59d6c1b921684b
7434 Oct 21 1999 /home/documents/latex/pkg/lgrind.sty ea74beead1aa2b711ec2669ba60562c3
7162 Nov 13 2015 /home/documents/latex/pkg/mol2chemfig.sty f5a8b1719cee30a4df0739275ac75f8a
1069 Oct 15 19:43 /home/documents/latex/share/includes/disclaimer-gaslight.tex 94142
    bbe063984d082bff3b400abe0fb
```

- 425 Oct 11 2020 /home/documents/latex/share/includes/disclaimer-status.tex b276f09e06a3a9114f927e4199f379f7 1403 Dec 4 20:31 /home/documents/latex/share/includes/mjmaddbib.tex 8e515fcae10a0e8a83e7c7cb0a2b633a 2921 Dec 19 09:24 /home/documents/latex/share/includes/mycommands.tex 0acd774532a2c513df384d1a80f02dd2 2901 Jun 17 2020 /home/documents/latex/share/includes/myskeletonpackages.tex fcfcd2e3c8d69d533932edaaa47f53a1 1538 Aug 14 07:26 /home/documents/latex/share/includes/recent_template.tex 49763d2c29f74e4b54fa53b25c2cc439 940 Jul 24 2019 /home/marchywka/.texmf-var/fonts/tfm/public/amsfonts/cmextra/cmex7.tfm f9e66c0105a30e64e3a0f5c4f79efb8d 852 Jul 24 2019 /home/marchywka/.texmf-var/fonts/tfm/public/amsfonts/symbols/msam10.tfm b4a46d2c220ee4ffaaf87c608f8593cd 860 Jul 24 2019 /home/marchywka/.texmf-var/fonts/tfm/public/amsfonts/symbols/msam5.tfm c4142ffef6136ff95621fbe99efb7cec 864 Jul 24 2019 /home/marchywka/.texmf-var/fonts/tfm/public/amsfonts/symbols/msam7.tfm 2998 d813a00ebf21070684f214a50f7e 844 Jul 24 2019 /home/marchywka/.texmf-var/fonts/tfm/public/amsfonts/symbols/msbm10.tfm f7721eee07bdc9e743e6c5f3f7e3d06d 876 Jul 24 2019 /home/marchywka/.texmf-var/fonts/tfm/public/amsfonts/symbols/msbm5.tfm 9 e3df3efef7afc4b0381e88a6402f777 876 Jul 24 2019 /home/marchywka/.texmf-var/fonts/tfm/public/amsfonts/symbols/msbm7.tfm 374365713297 d597717720c5786882e5 1260 Jul 24 2019 /home/marchywka/.texmf-var/fonts/tfm/public/cm/cmbx12.tfm 41596a2c763cf972bbdd853b378ec55a 1264 Jul 24 2019 /home/marchywka/.texmf-var/fonts/tfm/public/cm/cmbx9.tfm c3f8c3f0292777e1e9153581c59f8506 1448 Jul 24 2019 /home/marchywka/.texmf-var/fonts/tfm/public/cm/cmmi6.tfm be0f1d444547257aeb3f042af14f3e47 1456 Jul 24 2019 /home/marchywka/.texmf-var/fonts/tfm/public/cm/cmmi8.tfm e7bb485e28fc530112b40f5c89496200 1224 Jul 24 2019 /home/marchywka/.texmf-var/fonts/tfm/public/cm/cmr12.tfm 48d5728dc6473917c0e45f34e6a0e9cd 1236 Jul 24 2019 /home/marchywka/.texmf-var/fonts/tfm/public/cm/cmr6.tfm 63e3c1344d1e22a058a5cb87731337e0 1228 Jul 24 2019 /home/marchywka/.texmf-var/fonts/tfm/public/cm/cmr8.tfm 29a15bf51bfb16348a5cabb3215cf3fd 1228 Jul 24 2019 /home/marchywka/.texmf-var/fonts/tfm/public/cm/cmr9.tfm b0280c40050dc3527dafc7c425060d31 1052 Jul 24 2019 /home/marchywka/.texmf-var/fonts/tfm/public/cm/cmsy6.tfm 00c03700e0e2f29cde6c0b50a5c56df5 1056 Jul 24 2019 /home/marchywka/.texmf-var/fonts/tfm/public/cm/cmsy8.tfm 7ef80e56d3b9e223d3bce5a9065b95ad 273271 Dec 22 05:18 ../oldvit/oldvit.bib 9868023eea3bfe619f5dabad07270847 415 Dec 21 06:51 pmc_aapp.bib 8a4f32ea634eee6ed82e5af426ba8c29 429 Dec 22 08:14 releases.tex 62d3e038c1290b0c44fc490fd496c0db
- ffdaefb09013f5fd4b31e485c13933c1
 15 Jul 23 2019 /var/lib/texmf/fonts/map/pdftex/updmap/pdftex.map -> pdftex_dl14.map 341
 e900b5c25cd7029252688d79936e4

31050 Jul 21 2011 /usr/share/texlive/texmf-dist/bibtex/bst/urlbst/plainurl.bst

1431330 Jul 23 2019 /var/lib/texmf/web2c/pdftex/pdflatex.fmt c2a1d33979bfd7080ba601d397e5e158 243728 Dec 22 08:16 aapp.pdf cdf72e90cfbfe6809435bd9978d809d5