


# Modeling Intervention Strategies for TB Control in the United States

Jessica Ginepro, Emma Hartman, Ryo Kimura, Matthew McDermott, Colin Pawlowski, & Dylan Shepardson  
Mathematical Modeling Group, Mount Holyoke College, South Hadley, MA, USA

## Introduction

Epidemiological models offer insight into the structure of disease outbreaks and the merits of various interventions. The most common epidemiological models are compartmental differential equation models, such as the SIR system, illustrated in figures 1 and 2.




**Figure:** This flowchart depicts the standard SIR epidemiological model. It is accompanied by the system of differential equations 2.

$$\begin{aligned}\frac{dS}{dt} &= -\beta S(t)I(t) \\ \frac{dI}{dt} &= \beta S(t)I(t) - \gamma I(t) \\ \frac{dR}{dt} &= \gamma I(t) \\ N &= S(t) + I(t) + R(t)\end{aligned}$$

**Figure:** The system of differential equations governing the SIR model.

## The Basic Hill Model

In order to model tuberculosis (TB) in the United States (US), Hill, Becerra, and Castro designed a complex compartmental model called the hill model.



**Figure:** A flow chart representing the compartments of the Hill Model.


Populations:

- US Born Individuals (USB)
  - Foreign Born Individuals (FB)
- Individuals also leave the model due to natural death.

## Analyzing US TB Reduction Strategies


1. Implemented in R, with various numerical DE solvers.
2. Tracks US Health Care System (HCS) cost.
3. Tracks statistics about various health states.

## Basic Hill Behaviour



**Figure:** A plot showing the source population of yearly US TB incidence.

**Figure:** New cases per year of various types of TB in the US.




**Figure:** A graph of the projected incidence levels per million in the US-born, Foreign-born, and Total US population, given that LTBI rates in Foreign-born arrivals are reduced by 0%, 50%, 75%, and 100%.

## Economic Modeling


We extended our basic implementation of the Hill model to incorporate economic data of treatment costs for Active and Latent Tuberculosis. In our model, we estimate the average health care costs to be \$14,014.90 and \$403.45 to treat a single case of Active or Latent TB respectively. Given these costs and the

## An Agent Based Implementation

In addition to the extended Hill model (implemented in R), we also created a stochastic agent-based counterpart in both NetLogo and C++. Stochastic models add notions of uncertainty into the model structure, while agent-based models simulate global behavior of the system by describing local behavior of individual agents in the population.



**Figure:** Incidence per million for R and NetLogo models (12 runs)



**Figure:** Incidence per million for R and C++ models (2100 runs)

## Stochastic Models as a Measure of Variability