Modeling Intervention Strategies for United States TB Control

Jessica Ginepro, Emma Hartman, Ryo Kimura, Matthew McDermott, Colin Pawlowski, & Dylan Shepardson Mathematical Modeling Group, Mount Holyoke College, South Hadley, MA, USA





Introduction

- ► Epidemiological models describe disease outbreaks.
- Compartmental Differential Equation models are common and rigorous epidemiological models.
- ► The Hill Model is a complex compartmental model of tuberculosis (TB) in the United States (US).

The Basic Hill Model




Figure: A flow chart representing the compartments of the Hill Model.

Populations:

- US Born Individuals (USB)
- Foreign Born Individuals (FB)

Individuals also leave the model due to natural death.





Figure: The source population of US TB incindecence



Analyzing US TB Reduction Strategies

- Implemented in R, with various numerical DE solvers.
- ► Tracks US Health Care System (HCS) cost.
- Tracks statistics about various health states.

Intervention Analysis




Figure: Incidence/million in USB, FB, and total populations, given 0%, 50%, 75%, or 100% treatment of incoming LTBI.




Figure: Incidence/million in USB, FB, and total populations, given 0%, 100%, or 300% LTBI treatment increase.

Base HCS

Active TB:

\$14,014.90

\$403.45

Losts:

TBI:

Economic Modeling

- ► Tracks treatment costs for various disease states
- Estimates implementation cost of intervention




Figure: Cumulative implementation costs, US HCS savings, and net US costs of LTBI arrival cure rates. Cost/case cured was \$600, \$800, and \$1000 for 50%, 75%, and 100% cured.





Figure: Incidence/million for R and NetLogo models (12 runs, $\Delta t = 0.1$, popConst = 100)






Figure: Incidence/million for R and C++ models (2100 runs, $\Delta t = 0.01$, popConst = 1)

Stochastic Models as a Measure of Variability


An Agent Based Implementation


Agent based models capture disease dynamics on the

individual level, and reflect stochasticity and

granularity lost in compartmental models. Agent

The stochastic model provides data on the variability of the results of the deterministic model.








Figure: Distribution of FB Incidence (C++) with fitted Normal curve

Future Extensions

- Including contact structure
- Multi-drug resistant TB
- ► HIV