

Correction

Loss of mouse *Stmn2* function causes motor neuropathy

Iruñe Guerra San Juan, Leslie A. Nash, Kevin S. Smith, Marcel F. Leyton-Jaimes, Menglu Qian, Joseph R. Klim, Francesco Limone, Alexander B. Dorr, Alexander Couto, Greta Pintacuda, Brian J. Joseph, D. Eric Whisenant, Caroline Noble, Veronika Melnik, Deirdre Potter, Amie Holmes, Aaron Burberry, Matthijs Verhage, and Kevin Eggan*

*Correspondence: kevin.eggan@bmrn.com
<https://doi.org/10.1016/j.neuron.2022.11.003>

(Neuron 110, 1671–1688; May 18, 2022)

It was brought to our attention that there is a clerical mistake in one of the panels of Figure 1 (Figure 1E) in our recent publication. The representative blots for the CNS *Stmn2* protein levels in the F₂ generation of *Stmn2* mutants included an incorrect duplicated blot for the GAPDH loading control in the brain samples. We have corrected this error and replaced the figure online with one that includes the correct blot for the GAPDH loading control for those samples. We would like to profoundly apologize to the readers for the oversight and honest mistake.

E Stmn2 Spinal Cord and Brain protein levels




Figure 1E. CRISPR/Cas9 editing of *Stmn2* (corrected)

E Stmn2 Spinal Cord and Brain protein levels




Figure 1E. CRISPR/Cas9 editing of *Stmn2* (original)