

Submission for BCX-GRAPH

OTIS (internal use)

Michael Middlezong

August 8, 2024

Example (Canada 2006/4, 0♣). Consider a round-robin tournament with $2k + 1$ teams, where each team plays each other team exactly once. We say that three teams X , Y and Z , form a *cyclic triplet* if X beats Y , Y beats Z and Z beats X . There are no ties. Find the minimum and maximum possible number of cyclic triplets.

06CAN4

Walkthrough. The minimum bound is not that interesting.

- (a) Give an example of a tournament with no cyclic triplet. This finds the minimum.

It's the maximum that we'll be most interested in. For a team v , let $\text{outdeg } v$ denote the number of teams beaten by v . (This notation is the standard graph-theoretic one.) In order to count it, it will actually be parametrize our target in terms of degrees.

- (b) Rephrase the “maximum” problem in terms of the number of *non-cyclic* triplets.
- (c) By double-counting, find an expression for the number of non-cyclic triplets in terms of the outdegrees of the vertices. (Possible hint: every non-cyclic triplet can be labeled vwx with $v \rightarrow w$, $v \rightarrow x$.)

Thus we are reduced to an algebraic calculation.

- (d) Use Jensen's inequality to show there are at least $(2k + 1)\binom{k}{2}$ non-cyclic triplets. (If you've never seen Jensen's inequality before, you can read about it in Section 2.5.1 of the OTIS Excerpts.)
- (e) Give an example where equality holds; thus the maximum is $\binom{2k+1}{3} - (2k + 1)\binom{k}{2}$.

Example (BAMO 2005/4, 0♣). Let G be a connected graph on n vertices. Cynthia wishes to select some subset S of the edges so that every vertex is an endpoint of an odd number of edges in S .

- (a) Prove that Cynthia can never fulfill her wish if n is odd.
- (b) Prove that Cynthia can always fulfill her wish if n is even.

05BAMO4

Walkthrough. The first part (a) is not so hard (parity arguments) so this walkthrough focuses on (b).

- (a) Why can't we just use induction by removing vertices?
- (b) Figure out under what circumstances you can remove an edge, though.
- (c) Reduce the case to where G is a tree.
- (d) When G is a tree, certain moves are now "forced"; what are they? (Possible hint: look at leaves.)
- (e) By "repeating" (d), describe an algorithm such that, for any tree G , produces a set S with the desired property. Prove that it exists.
- (f) Optionally, show that S is unique (when G is a tree). There is a good chance this follows from your solution to (d) and (e) and doesn't require any additional insights.
- (g) Where did we use the hypothesis that G is connected?

Practice problems

Instructions: Solve [30♣]. If you have time, solve [40♣].

Induction is the coordinate bashing of graph theory.

A MOP 2012 grader

Problem 1 (Chartrand-Zhang 2.14, 2♣)

Let G be a finite graph such that every edge of G has exactly one endpoint of even degree. Show that G has an even number of edges.

CZHG214

We use a double counting approach. Let an edge sum be the sum of the degrees of the two endpoints of an edge. Consider the total edge sum over all edges in the graph. Since each individual edge sum must be odd, it suffices to show that this total is even.

It's not hard to see that this total should equal

$$\sum_v (\deg v)^2.$$

Finally, this quantity is even because it has the same parity as $\sum_v \deg v$, so we are done.

Required Problem 2 (2♣)

Prove:

- (a) Let G be a simple undirected graph. Show that

$$\sum_v \deg v = 2\#E.$$

- (b) Let G be a directed graph. We say the *indegree* of a vertex v (denoted $\text{indeg } v$) is the number of incoming edges, while the *outdegree* (denoted $\text{outdeg } v$) is the number of outgoing edges. Show that

$$\sum_v \text{indeg } v = \sum_v \text{outdeg } v = \#E.$$

- (c) Continuing the notation of (b), suppose further that G is a tournament, i.e. every pair of vertices has exactly one edge between them oriented in one of the two directions. Prove that

$$\sum_v (\text{indeg } v)^2 = \sum_v (\text{outdeg } v)^2.$$

GRTH5

- (a) Each edge contributes 2 to the total degree, one for each vertex it touches. (Formally we could use induction.)
- (b) Every directed edge contributes 1 to the total indegree and 1 to the total outdegree.

(c) In a graph with n vertices, we have

$$\begin{aligned}
 \sum_v (\text{indeg } v)^2 - \sum_v (\text{outdeg } v)^2 &= \sum_v (\text{indeg } v)^2 - (\text{outdeg } v)^2 \\
 &= \sum_v (\text{indeg } v + \text{outdeg } v)(\text{indeg } v - \text{outdeg } v) \\
 &= (n-1) \sum_v \text{indeg } v - \text{outdeg } v \\
 &= 0,
 \end{aligned}$$

with the last step using the result of part (b). This concludes the proof.

Problem 3 (Chartrand-Zhang 2.36, 2♣)

Find the smallest positive integer k for which there exists a simple graph on $3k$ vertices, for which exactly k vertices have degree 2, exactly k vertices have degree 6, and exactly k vertices have degree 7.

CZHG236

The answer is $k = 4$. In order for a vertex to have degree 7, we must have $k \geq 3$. We also know that $k = 3$ is impossible, because the total degree must be even.

A construction for $k = 4$ is to first split the vertices into three groups of 4. Then, each vertex in group 1 is connected with all four vertices in group 2 and the other three vertices in group 1. Next, each vertex in group 2 is connected with all four vertices in group 1 and two vertices in group 3, in such a way that each vertex in group 3 is hit twice. In this construction, the vertices in group 1 have degree 7, the vertices in group 2 have degree 6, and the vertices in group 3 have degree 2.

Problem 4 (2♣)

Let G be a graph on $n \geq 2$ vertices. Prove that either G or its complement is a connected graph. (The *complement* of a graph, denoted \overline{G} , is the one which has the same vertices as G but has an edge between two vertices if and only if the corresponding edge G is *not* present.)

GRTH4

Assume G is not a connected graph. Then, it has at least two connected components.

We aim to show that there is a path between any two vertices in \overline{G} . Let a and b be two vertices, and let A and B be their respective connected components in G .

If A and B are distinct connected components, then notice that a and b must be directly connected by an edge in \overline{G} .

Otherwise, let C be a connected component distinct from A and B , and let $c \in C$. Our desired path is $a \rightarrow c \rightarrow b$.

Problem 5 (CGMO 2012/6, 3♣)

There are n cities, 2 airline companies in a country. Between any two cities, there is exactly one 2-way flight connecting them which is operated by one of the two companies. A female mathematician plans a travel route, so that it starts and ends at the same city, passes through at least two other cities, and each city in the route is visited once. She finds out that wherever she starts and whatever route she chooses, she must take flights of both companies. Find the maximum value of n .

12CGM06

The answer is $n = 4$. The problem statement's condition is equivalent to saying that there are no monochromatic cycles in some red-blue edge labeling of K_n . It is not hard to construct such an edge coloring for K_4 that works by manually checking.

It is well known that when $n \geq 6$, any red-blue edge coloring of K_n will contain a monochromatic triangle (proof is by Pigeonhole). In order to not create a monochromatic triangle when $n = 5$, each vertex must have exactly 2 red edges and 2 blue edges emanating from it. (Otherwise, in an attempt to not create monochromatic triangles, the endpoints of any 3 vertices of the same color will form a monochromatic triangle.) This information is enough to make a constructive argument for why $n = 5$ is impossible, using casework at any point where a non-obvious decision must be made.

Problem 6 (USAMO 1981/2, 3♣)

Every pair of communities in a county are linked directly by one mode of transportation: bus, train, or airplane. All three methods of transportation are used in the county with no community being serviced by all three modes and no three communities being linked pairwise by the same mode. Determine the largest number of communities in this county.

81AM02

Problem 7 (3♣)

One day, some students living in the University of Nebraska-Lincoln are friends with each other. The flu affects each student for one day; then the student is immune the next day. On any day n , a healthy student cares for all their sick friends, and falls ill on day $n + 1$ if they were not immune on day n .

- (a) Prove that eventually the epidemic will die out.
- (b) Suppose that immediately before the epidemic some students got vaccinated and thus had an additional immunity on day 1. Give an example to show the epidemic may continue indefinitely (for some choice of initial students and knowledge relation).

SICKNESS

Problem 8 (5♣)

Let $n \geq 3$ be an integer. Prove that if the edges of K_n are colored with n colors, with each color used at least once, then some triangle has its three edges of different colors.

H2024647

Problem 9 (Chartrand-Zhang 1.17, 3♣)

Let G be a connected graph on finitely many vertices. Suppose that the longest path in G has length ℓ . Prove that any two paths of length ℓ share a vertex.

CZHGT117

Suppose for contradiction that two paths of length ℓ do not share a vertex. Let $a_1, a_2, \dots, a_{\ell+1}$ and $b_1, b_2, \dots, b_{\ell+1}$ denote the vertex sequences of two paths of length ℓ which do not share a vertex.

Since G is connected, there must exist vertices a_i and b_j such that there exists a path from a_i to b_j which is disjoint from the two original paths. (Proof sketch: consider a

path from a_1 to b_1 and "remove" the two original paths from it.) We claim that either the path from $a_1 \rightarrow a_i \rightarrow b_j \rightarrow b_1$ or the path from $a_{l+1} \rightarrow a_i \rightarrow b_j \rightarrow b_{l+1}$ must have length greater than l .

Let m be the length of the path from a_i to b_j which is disjoint from the two original paths. Then, the length of the path from $a_1 \rightarrow a_i \rightarrow b_j \rightarrow b_1$ is

$$i + m + j,$$

and the length of the path from $a_{l+1} \rightarrow a_i \rightarrow b_j \rightarrow b_{l+1}$ is

$$(l - i) + m + (l - j).$$

These two lengths add up to $2l + 2m > 2l$, so clearly, one of them exceeds l .

Problem 10 (Chartrand-Zhang 3.16, 3♣)

Up to isomorphism, how many graphs on 9 vertices have every vertex of degree 6? (Two graphs are isomorphic if one can relabel the vertices of one to get the other.)

CZHGT316

Problem 11 (Shortlist 2001 C3, 3♣)

Let G be a finite simple graph with no K_5 , and for which any two triangles share at least one common vertex. Prove that one can delete two vertices from G and obtain a graph with no triangles.

01SLC3

Required Problem 12 (PUMaC Finals 2019 A1, 5♣)

A finite graph G is drawn on a blackboard. The following operation is permitted: pick any cycle C of G , draw a new vertex v , connect it to all vertices of C , and finally erase all the edges of C . Prove that this operation can only be done a finite number of times.

19PUMACFA1

Problem 13 (Tuymaada 2016/8, 5♣)

Let G be a finite connected graph. Prove that one can color the vertices blue and green, and the edges black and white, such that the following three conditions are fulfilled:

- Black edges join vertices of different colors.
- White edges have at least one blue endpoint.
- Any two vertices are connected by a black path.

16TMD8

Problem 14 (Shortlist 2017 C2, 3♣)

Let n be a positive integer. A *chameleon* is a sequence of $3n$ letters with exactly n occurrences of each of the letters a, b, c . Define a *swap* to be the transposition of two adjacent letters in the chameleon. Prove that for any X there exists a chameleon Y such that X cannot be changed to Y using fewer than $3n^2/2$ swaps.

17SLC2

Problem 15 (Balkan 2013/4, 9♣)

Let G be a graph with the property that for any chordless cycle C and any vertex v not in C , at most one vertex of C is adjacent to v . Prove that G is three-colorable.

(A *chordless cycle* is a cycle where no two vertices are adjacent unless they are consecutive in the cycle.)

13BALK4