

Spin-averated $\Lambda_c^{**+} \rightarrow \Lambda_c^+ \pi^+ \pi^-$ distribution

The notebook computes the **spin-averated** angular functions for the three body decays of the excited Λ_c^{**+} states under different hypotheses of its quantum numbers.

The spin factors are evaluated using conventions of the three-body decays introduced in the Dalitz plot decomposition paper.

The implementation in Julia comes from the [ThreeBodyDecay.jl](#) package. A dispatch on SymPy object is used to provide result in a nice symbolic form. The dispatch is defined in [SymbolicThreeBodyDecay.jl](#) package.

The spin sums are evaluated symbolically. The resulting decay intensity distribution is a bilinear form on isobars lineshapes.

$$I = |R^1|^2 T_{1,1}(\theta, \zeta) + |R^2|^2 T_{2,2}(\theta, \zeta) + 2\text{Re}(R^{1*} R^2) T_{1,2}(\theta, \zeta)$$

where I is a differential decay rate (density of the events on the Dalitz plot), R^i is the parametrization of the lineshape of an intermediate resonance i , and θ, ζ are kinematic angles that are unambiguously expressed using the Mandelstam invariants.

- The LS couplings are used for parametrizing the decay vertices. The minimal orbital angular momentum is chosen is several value are possible.
- The matrix T gives only the angular factors, the break-up momentum for $p^L q^l$ is attributed to the lineshape function R , with L being the orbital angular momentum in the decay of Λ_c^{**+} , and l for the decay of an intermediate resonance.
- We incorporate **six** subchannel resonances: f_0 , and ρ in $\pi\pi$, as well as $\Sigma_c(2455)$ and $\Sigma_c(2520)$

Selection deleted for both $\Lambda_c \pi$ channels

- Value of l is unambiguously in all channels, $l = 0$ for $f_0 \rightarrow \pi\pi$, and $l = 1$ for all other resonances.
- The value of L in the decay of Λ_c^{**+} is indicated in the lineshape function:

$$R_{L,S}^{\text{resonance}}$$

How to related $R^{\Sigma(2455)++}$ and $R^{\Sigma(2455)0}$?

The isospin Clebsch in the Λ_c^{*++} decay is the same

$$\langle \Sigma(2245)^{++}; \pi^- | \Lambda_c^{*++} \rangle = \langle 1, 1; 1, -1 | 0, 0 \rangle = \frac{1}{\sqrt{3}},$$
$$\langle \Sigma(2245)^0; \pi^+ | \Lambda_c^{*++} \rangle = \langle 1, -1; 1, 1 | 0, 0 \rangle = \frac{1}{\sqrt{3}},$$

There is a difference treatment of the Σ_c decays.

For $\Sigma_c^{++}(12)$ decaying to $\Lambda_c^+(1)\pi^+(2)$ in P-wave, the helicity coupling is related to the LS coupling with Clebsches which are the same

$$H_{\nu; \nu, 0}^{\Sigma_c^{++} \rightarrow \Lambda_c^+ \pi^+} = \langle 1, 0; 1/2, \nu | 1/2, \nu \rangle \langle 1/2, \nu; 0, 0 | 1/2, \nu \rangle H_{P\text{-wave}}^{\Sigma_c^{++} \rightarrow \Lambda_c^+ \pi^+},$$
$$H_{\nu; 0, \nu}^{\Sigma_c^0 \rightarrow \pi^- \Lambda_c^+} = \langle 1, 0; 1/2, \nu | 1/2, \nu \rangle \langle 1/2, \nu; 0, 0 | 1/2, \nu \rangle H_{P\text{-wave}}^{\Sigma_c^0 \rightarrow \pi^- \Lambda_c^+}$$

The symmetry of the LS coupling under permutation is the simplest,

$$H_{L\text{-wave}}^{\Sigma_c \rightarrow \Lambda_c \pi} = (-1)^L H_{L\text{-wave}}^{\Sigma_c \rightarrow \pi \Lambda_c}$$

The relation gives the way to relate the lineshape functions of Σ_c^0 and Σ_c^{++} :

$$R^{\Sigma_c^0} = -R^{\Sigma_c^{++}}.$$

```
1 begin
2   cd(mktempdir())
3   import Pkg
4   Pkg.activate(".")
5   Pkg.add([
6     Pkg.PackageSpec(url="https://github.com/mmikhasenko/ThreeBodyDecay.jl"),
7     Pkg.PackageSpec(url="https://github.com/mmikhasenko/SymbolicThreeBodyDecays.jl"),
8     Pkg.PackageSpec("Plots"), Pkg.PackageSpec("Parameters")
9   ])
10  #
11  using Parameters
12  using LinearAlgebra
13  using ThreeBodyDecay
14  using SymbolicThreeBodyDecays
15  using SymbolicThreeBodyDecays.SymPy
16  #
17  using Plots
18  import Plots.PlotMeasures: mm
19 end
```

```

1 theme(:wong2, frame=:box, grid=false, minorticks=true,
2       guidefontvalign=:top, guidefonthalign=:right,
3       xlim=(:auto, :auto), ylim=(:auto, :auto),
4       lw=1.2, lab="", colorbar=false,
5       bottom_margin=3mm, left_margin=3mm)

```

Integration with Symbolic Computation

```

1 begin
2     @syms m1 m2 m3 m0
3     @syms σ1 σ2 σ3
4 end;

```

```
const ms = (m1 =  $m_1$ , m2 =  $m_2$ , m3 =  $m_3$ , m0 =  $m_0$ )
```

```
1 const ms = ThreeBodyMasses(m1, m2, m3; m0)
```

```
const σs = (σ1 =  $\sigma_1$ , σ2 =  $\sigma_2$ , σ3 =  $m_0^2 + m_1^2 + m_2^2 + m_3^2 - \sigma_1 - \sigma_2$ )
```

```
1 const σs = Invariants(ms; σ1, σ2)
```

spinparity (generic function with 1 method)

```

1 function spinparity(p)
2     pt = (p[2]..., p[1])
3     jpv = str2jp.(pt)
4     getproperty.(jpv, :j) .|> x2 |> ThreeBodyDecay.SpinTuple,
5     getproperty.(jpv, :p) |> ThreeBodyDecay.ParityTuple
6 end

```

Selection deleted

Setup

```

1 begin
2   Base.@kwdef struct Setup{T}
3     two_js
4     decay_chains::Vector{T}
5   end
6   function Setup(chains::Vector{Tuple{String, String, Int64}}, ifstate)
7     js, Ps = ifstate |> spinparity
8     tbs = ThreeBodySystem(ms, js)
9     #
10    dcv = map(chains) do chain
11      nameR, jpR, k = chain
12      Rjp = str2jp(jpR)
13      #
14      d0 = DecayChainLS(k, identity; two_s=Rjp.j |> x2, parity=Rjp.p, Ps, tbs)
15      two_L, two_S = d0.HRk.two_ls
16      L = div(two_L, 2)
17      symR = sympy.Symbol(nameR * "-{$(L), $(two_S)/2}", real=true)
18      d = DecayChainLS(k, σ->symR; two_s=Rjp.j |> x2, parity=Rjp.p, Ps, tbs)
19    end
20    return Setup(tbs.two_js, dcv)
21  end
22 end

```

unpolarized_intensity (generic function with 1 method)

```

1 function unpolarized_intensity(setup; refζs=(1, 1, 1, 1))
2   #
3   @unpack two_js, decay_chains = setup
4   full_amplitude = sum(itr(two_js)) do two_λs
5     A = sum(decay_chains) do dc
6       amplitude(dc, σ |> StickySymTuple, two_λs .|> Sym; refζs)
7     end
8     abs2(A)
9   end
10  full_amplitude
11 end

```

intensity_matrix (generic function with 1 method)

```

1 function intensity_matrix(setup)
2   I_symm = unpolarized_intensity(setup; refζs=(1, 1, 1, 1));
3   lineshapes = map(d->d.Xlineshape(nothing), setup.decay_chains)
4   H = sympy.hessian(I_symm, lineshapes)
5   H_fully_simp = map(eachindex(Symmetric(H))) do i
6     simplify(H[i][2] ? Sym(0) :
7       simplify(H[i].doit()))
8   end |> Symmetric;
9   H_normalized = H_fully_simp.
10     xreplace(sign_replacements |> Dict).
11     xreplace(combine_replacements |> Dict)#.
12     # xreplace(unsafe_uncombine_replacement[2:3] |> Dict);
13   return H_normalized, lineshapes
14 end

```

Nice printing

latexalign (generic function with 1 method)

```
1 latexalign(H::Matrix) =
2     """
3     ``math
4     \\small
5     \\begin{align}
6     """ *
7     prod(map(eachindex(Symmetric(H))) do i
8         i[1]!=i[2] ? "" :
9         "T_{$(i[1]),$(i[2])} &= " * sympy.latex(H[i[1],i[2]]) * "\\,,\\\\"
10    end) *
11    prod(map(eachindex(Symmetric(H))) do i
12        i[1]<i[2] ?
13        "T_{$(i[1]),$(i[2])} &= " * sympy.latex(H[i[1],i[2]]) * "\\,,\\\\" : ""
14    end) *
15    prod(map(eachindex(Symmetric(H))) do i
16        i[1]>i[2] ? "T_{$(i[1]),$(i[2])} &= T_{$(i[2]),$(i[1])}" * "\\,,\\\\" : ""
17    end) *
18    """
19    \\end{align}
20    ``
21    """
```

Selection deleted

latexintensity (generic function with 2 methods)

```
1 latexintensity(V::Vector, Isubscript="") =
2 """
3 ```math
4 \\\small
5 \\\begin{equation}
6 {\\Large I_{$(Isubscript)}} =
7 \\\begin{pmatrix}
8 """ *
9     sympy.latex(V[1]) *
10     prod(map(V[2:end]) do v
11         " \\\ " * sympy.latex(v)
12     end) *
13 """
14 \\\end{pmatrix}^{\\\Large\\dagger}
15 \\\,,\\,, {\\Huge [T]} \\\,,\\,,
16 \\\begin{pmatrix}
17 """ *
18     sympy.latex(V[1]) *
19     prod(map(V[2:end]) do v
20         " \\\ " * sympy.latex(v)
21     end) *
22 """
23 \\\end{pmatrix}
24 \\\end{equation}
25 ```
26 """
```

printalign (generic function with 1 method)

```
1 printalign(H::Matrix) = Markdown.parse(latexalign(H))
```

Application

.....

final_state = ("1/2+", "0-", "0-")

```
1 final_state = ("1/2+", "0-", "0-")
```

reaction = "3/2-" ⇒ ("1/2+", "0-", "0-")

```
1 reaction = "3/2-" => final_state
```

Selection deleted

```
chains =
  [ ("R^{\rho(770)0}", "1-", 1), ("R^{f_0}", "0+", 1), ("R^{\Sigma(2455)0}", "1/2+", 2), ("
1 chains = [
2   ("R^{\rho(770)0}", "1-", 1), #pi+pi-
3   ("R^{f_0}", "0+", 1), #pi+pi-
4   #
5   ("R^{\Sigma(2455)0}", "1/2+", 2), #pi-Lc+
6   ("R^{\Sigma(2520)0}", "3/2+", 2), #pi-Lc+
7   #
8   ("R^{\Sigma(2455)++}", "1/2+", 3), #Lc+pi+
9   ("R^{\Sigma(2520)++}", "3/2+", 3), #Lc+pi+
10 ]
```

1/2- decay

1P states: 1/2-, 3/2-

```
s1h- =
  Setup((two_h1 = 1, two_h2 = 0, two_h3 = 0, two_h0 = 1), [DecayChain{var"#15#17"{Sym}, Rec
1 s1h- = Setup(chains, "1/2-" => final_state)
1 @time T1h- = intensity_matrix(s1h-);
92.660335 seconds (148.57 k allocations: 4.713 MiB)
```

Selection deleted

$$I_{1/2-} = \begin{pmatrix} R_{0,1/2}^{\rho(770)0} \\ R_{1,1/2}^{f_0} \\ R_{0,1/2}^{\Sigma(2455)0} \\ R_{2,3/2}^{\Sigma(2520)0} \\ R_{0,1/2}^{\Sigma(2455)++} \\ R_{2,3/2}^{\Sigma(2520)++} \end{pmatrix}^{\dagger} [T] \begin{pmatrix} R_{0,1/2}^{\rho(770)0} \\ R_{1,1/2}^{f_0} \\ R_{0,1/2}^{\Sigma(2455)0} \\ R_{2,3/2}^{\Sigma(2520)0} \\ R_{0,1/2}^{\Sigma(2455)++} \\ R_{2,3/2}^{\Sigma(2520)++} \end{pmatrix}$$

```
1 latexintensity(T1h-[2], "1/2-") |> Markdown.parse
```

Selection deleted

$$\begin{aligned}
T_{1,1} &= 2, \\
T_{2,2} &= 2, \\
T_{3,3} &= 2, \\
T_{4,4} &= 4 - 3 \sin^2(\theta_{31}), \\
T_{5,5} &= 2, \\
T_{6,6} &= 4 - 3 \sin^2(\theta_{12}), \\
T_{1,2} &= 2 \cos(\theta_{23}), \\
T_{1,3} &= 2 \cos\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} + \theta_{23} - \frac{\theta_{31}}{2}\right), \\
T_{2,3} &= 2 \cos\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\theta_{31}}{2}\right), \\
T_{1,4} &= -\frac{\sqrt{2}\left(\cos\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} + \theta_{23} - \frac{\theta_{31}}{2}\right) + 3 \cos\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} + \theta_{23} + \frac{3\theta_{31}}{2}\right)\right)}{2}, \\
T_{2,4} &= -\frac{\sqrt{2}\left(\cos\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\theta_{31}}{2}\right) + 3 \cos\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} + \frac{3\theta_{31}}{2}\right)\right)}{2}, \\
T_{3,4} &= \sqrt{2} \cdot (3 \sin^2(\theta_{31}) - 2), \\
T_{1,5} &= -2 \sin\left(\frac{\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} - \theta_{23}\right), \\
T_{2,5} &= -2 \sin\left(\frac{\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2}\right), \\
T_{3,5} &= -2 \sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} - \frac{\theta_{31}}{2}\right), \\
T_{4,5} &= \frac{\sqrt{2}\left(\sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} - \frac{\theta_{31}}{2}\right) + 3 \sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} + \frac{3\theta_{31}}{2}\right)\right)}{2}, \\
T_{1,6} &= \frac{\sqrt{2}\left(-3 \sin\left(-\frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\zeta_{2(3)}^1}{2} + \frac{3\theta_{12}}{2} + \theta_{23}\right) + \sin\left(\frac{\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} - \theta_{23}\right)\right)}{2}, \\
T_{2,6} &= \frac{\sqrt{2} \cdot \left(3 \sin\left(\frac{\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\zeta_{2(3)}^1}{2} - \frac{3\theta_{12}}{2}\right) + \sin\left(\frac{\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2}\right)\right)}{2}, \\
T_{3,6} &= -\frac{\sqrt{2}\left(-3 \sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(3)}^1}{2} - \frac{3\theta_{12}}{2} - \frac{\theta_{31}}{2}\right) - \sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} - \frac{\theta_{31}}{2}\right)\right)}{2}, \\
T_{4,6} &= \frac{3 \sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(3)}^1}{2} - \frac{3\theta_{12}}{2} - \frac{\theta_{31}}{2}\right)}{4} - \frac{9 \sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(3)}^1}{2} - \frac{3\theta_{12}}{2} + \frac{3\theta_{31}}{2}\right)}{4} - \frac{\sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} - \theta_{23}\right)}{4}, \\
T_{5,6} &= \sqrt{2} \cdot (3 \sin^2(\theta_{12}) - 2), \\
T_{2,1} &= T_{1,2}, \\
T_{3,1} &= T_{1,3}, \\
T_{4,1} &= T_{1,4}, \\
T_{5,1} &= T_{1,5}, \\
T_{6,1} &= T_{1,6}, \\
T_{3,2} &= T_{2,3}, \\
T_{4,2} &= T_{2,4}, \\
T_{5,2} &= T_{2,5}, \\
T_{6,2} &= T_{2,6}, \\
T_{3,3} &= T_{3,3}, \\
T_{4,3} &= T_{3,4}, \\
T_{5,3} &= T_{3,5}, \\
T_{6,3} &= T_{3,6}, \\
T_{4,4} &= T_{4,4}, \\
T_{5,4} &= T_{4,5}, \\
T_{6,4} &= T_{4,6}, \\
T_{5,5} &= T_{5,5}, \\
T_{6,5} &= T_{5,6}, \\
T_{6,6} &= T_{6,6}.
\end{aligned}$$

```

-2,4 -4,2,
T5,2 = T2,5,
T6,2 = T2,6,
T4,3 = T3,4,
T5,3 = T3,5,
T6,3 = T3,6,
T5,4 = T4,5,
T6,4 = T4,6,
T6,5 = T5,6,

```

```
1 printalign(T1h-[1])
```

```
s3h- =
```

```
Setup((two_h1 = 1, two_h2 = 0, two_h3 = 0, two_h0 = 3), [DecayChain{var"#15#17"{Sym}, Rec
```

```
1 s3h- = Setup(chains, "3/2-" => final_state)
```

```
1 @time T3h- = intensity_matrix(s3h-);
```

```
331.829005 seconds (511.30 k allocations: 16.139 MiB)
```

$$I_{3/2-} = \begin{pmatrix} R_{0,3/2}^{\rho(770)0} \\ R_{1,1/2}^{f_0} \\ R_{2,1/2}^{\Sigma(2455)0} \\ R_{0,3/2}^{\Sigma(2520)0} \\ R_{2,1/2}^{\Sigma(2455)++} \\ R_{0,3/2}^{\Sigma(2520)++} \end{pmatrix}^{\dagger} [T] \begin{pmatrix} R_{0,3/2}^{\rho(770)0} \\ R_{1,1/2}^{f_0} \\ R_{2,1/2}^{\Sigma(2455)0} \\ R_{0,3/2}^{\Sigma(2520)0} \\ R_{2,1/2}^{\Sigma(2455)++} \\ R_{0,3/2}^{\Sigma(2520)++} \end{pmatrix}$$

```
1 latexintensity(T3h-[2], "3/2-") |> Markdown.parse
```

Selection deleted

$$T_{1,1} = 2,$$

$$T_{2,2} = 2,$$

$$T_{3,3} = 2,$$

$$T_{4,4} = 2,$$

$$T_{5,5} = 2,$$

$$T_{6,6} = 2,$$

$$T_{1,2} = 2 \cos(\theta_{23}),$$

$$T_{1,3} = \frac{\cos\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} + \theta_{23} - \frac{\theta_{31}}{2}\right)}{2} + \frac{3 \cos\left(\frac{3\zeta_{1(2)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \theta_{23} + \frac{\theta_{31}}{2}\right)}{2},$$

$$T_{2,3} = \frac{\cos\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\theta_{31}}{2}\right)}{2} + \frac{3 \cos\left(\frac{3\zeta_{1(2)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\theta_{31}}{2}\right)}{2},$$

$$T_{1,4} = \frac{\cos\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} + \theta_{23} - \frac{\theta_{31}}{2}\right)}{2} + \frac{3 \cos\left(\frac{3\zeta_{1(2)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \theta_{23} - \frac{3\theta_{31}}{2}\right)}{2},$$

$$T_{2,4} = \frac{3 \cos\left(-\frac{3\zeta_{1(2)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} + \frac{3\theta_{31}}{2}\right)}{2} + \frac{\cos\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\theta_{31}}{2}\right)}{2},$$

$$T_{3,4} = 2 - 3 \sin^2(\theta_{31}),$$

$$T_{1,5} = -\frac{3 \sin\left(-\frac{3\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} + \theta_{23}\right)}{2} - \frac{\sin\left(\frac{\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} - \theta_{23}\right)}{2},$$

$$T_{2,5} = -\frac{\sin\left(\frac{\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2}\right)}{2} + \frac{3 \sin\left(\frac{3\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2}\right)}{2},$$

$$T_{3,5} = -\frac{\sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} - \frac{\theta_{31}}{2}\right)}{2} + \frac{3 \sin\left(\frac{3\zeta_{1(2)}^0}{2} + \frac{3\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2} + \frac{\theta_{31}}{2}\right)}{2},$$

$$T_{4,5} = -\frac{3 \sin\left(-\frac{3\zeta_{1(2)}^0}{2} - \frac{3\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} + \frac{3\theta_{31}}{2}\right)}{2} - \frac{\sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} - \frac{\theta_{31}}{2}\right)}{2},$$

$$T_{1,6} = -\frac{\sin\left(\frac{\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} - \theta_{23}\right)}{2} + \frac{3 \sin\left(\frac{3\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\zeta_{2(3)}^1}{2} + \frac{3\theta_{12}}{2} - \theta_{23}\right)}{2},$$

$$T_{2,6} = -\frac{\sin\left(\frac{\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2}\right)}{2} + \frac{3 \sin\left(\frac{3\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\zeta_{2(3)}^1}{2} + \frac{3\theta_{12}}{2}\right)}{2},$$

$$T_{3,6} = -\frac{\sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} - \frac{\theta_{31}}{2}\right)}{2} + \frac{3 \sin\left(\frac{3\zeta_{1(2)}^0}{2} + \frac{3\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(3)}^1}{2} + \frac{3\theta_{12}}{2} + \frac{\theta_{31}}{2}\right)}{2},$$

$$T_{4,6} = -\frac{\sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} - \frac{\theta_{31}}{2}\right)}{2} + \frac{3 \sin\left(\frac{3\zeta_{1(2)}^0}{2} + \frac{3\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(3)}^1}{2} + \frac{3\theta_{12}}{2} - \frac{3\theta_{31}}{2}\right)}{2},$$

$$T_{5,6} = 2 - 3 \sin^2(\theta_{12}),$$

$$T_{2,1} = T_{1,2},$$

$$T_{3,1} = T_{1,3},$$

$$T_{4,1} = T_{1,4},$$

$$T_{5,1} = T_{1,5},$$

$$T_{6,1} = T_{1,6},$$

$$\pi = \pi$$

$$\begin{aligned} I_{3,2} &= I_{2,3}, \\ T_{4,2} &= T_{2,4}, \\ T_{5,2} &= T_{2,5}, \\ T_{6,2} &= T_{2,6}, \\ T_{4,3} &= T_{3,4}, \\ T_{5,3} &= T_{3,5}, \\ T_{6,3} &= T_{3,6}, \\ T_{5,4} &= T_{4,5}, \\ T_{6,4} &= T_{4,6}, \\ T_{6,5} &= T_{5,6}, \end{aligned}$$

```
1 printalign(T3h[1])
```

1D states: 3/2+ 5/2+

s3h* =

```
Setup((two_h1 = 1, two_h2 = 0, two_h3 = 0, two_h0 = 3), [DecayChain{var"#15#17"{Sym}, Rec
```

```
1 s3h* = Setup(chains, "3/2+" => final_state)
```

```
1 @time T3h* = intensity_matrix(s3h*);
```

```
374.349996 seconds (511.30 k allocations: 16.139 MiB, 0.02% gc time)
```

$$I_{3/2+} = \begin{pmatrix} R_{1,1/2}^{\rho(770)0} \\ R_{2,1/2}^{f_0} \\ R_{1,1/2}^{\Sigma(2455)0} \\ R_{1,3/2}^{\Sigma(2520)0} \\ R_{1,1/2}^{\Sigma(2455)++} \\ R_{1,3/2}^{\Sigma(2520)++} \end{pmatrix}^{\dagger} [T] \begin{pmatrix} R_{1,1/2}^{\rho(770)0} \\ R_{2,1/2}^{f_0} \\ R_{1,1/2}^{\Sigma(2455)0} \\ R_{1,3/2}^{\Sigma(2520)0} \\ R_{1,1/2}^{\Sigma(2455)++} \\ R_{1,3/2}^{\Sigma(2520)++} \end{pmatrix}$$

```
1 latexintensity(T3h*[2], "3/2+") |> Markdown.parse
```


$$T_{1,1} = 2,$$

$$T_{2,2} = 2,$$

$$T_{3,3} = 2,$$

$$T_{4,4} = \frac{12 \sin^2(\theta_{31})}{5} + \frac{2}{5},$$

$$T_{5,5} = 2,$$

$$T_{6,6} = \frac{12 \sin^2(\theta_{12})}{5} + \frac{2}{5},$$

$$T_{1,2} = 2 \cos(\theta_{23}),$$

$$T_{1,3} = \frac{\cos\left(\frac{\zeta_{1(2)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} - \theta_{23} + \frac{\theta_{31}}{2}\right)}{2} + \frac{3 \cos\left(\frac{3\zeta_{1(2)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} + \theta_{23} - \frac{\theta_{31}}{2}\right)}{2},$$

$$T_{2,3} = \frac{\cos\left(\frac{\zeta_{1(2)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\theta_{31}}{2}\right)}{2} + \frac{3 \cos\left(\frac{3\zeta_{1(2)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\theta_{31}}{2}\right)}{2},$$

$$T_{1,4} = \frac{\sqrt{5} \cdot \left(12 \cos\left(-\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} + \theta_{23} + \frac{3\theta_{31}}{2}\right) - 8 \cos\left(\frac{\zeta_{1(2)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} - \theta_{23} + \frac{\theta_{31}}{2}\right) + 12 \cos\left(\frac{3\zeta_{1(2)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\theta_{31}}{2}\right) - 40\right)}{40}$$

$$T_{2,4} = \frac{\sqrt{5} \cdot \left(3 \cos\left(-\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} + \frac{3\theta_{31}}{2}\right) - 2 \cos\left(\frac{\zeta_{1(2)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\theta_{31}}{2}\right) + 3 \cos\left(\frac{3\zeta_{1(2)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\theta_{31}}{2}\right)\right)}{10},$$

$$T_{3,4} = \frac{\sqrt{5} \cdot (3 \cos(2\theta_{31}) + 1)}{10},$$

$$T_{1,5} = \frac{\sin\left(\frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2} + \theta_{23}\right)}{2} - \frac{3 \sin\left(\frac{3\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} - \theta_{23}\right)}{2},$$

$$T_{2,5} = \frac{\sin\left(\frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2}\right)}{2} - \frac{3 \sin\left(\frac{3\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2}\right)}{2},$$

$$T_{3,5} = \frac{\sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2} + \frac{\theta_{31}}{2}\right)}{2} - \frac{3 \sin\left(\frac{3\zeta_{1(2)}^0}{2} + \frac{3\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} - \frac{\theta_{31}}{2}\right)}{2},$$

$$T_{4,5} = \frac{\sqrt{5} \left(-3 \sin\left(-\frac{\zeta_{1(2)}^0}{2} - \frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} + \frac{3\theta_{31}}{2}\right) - 2 \sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2} + \frac{\theta_{31}}{2}\right) - 3 \sin\left(-\frac{3\zeta_{1(2)}^0}{2} - \frac{3\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} - \frac{\theta_{31}}{2}\right) + 10\right)}{10}$$

$$T_{1,6} = \frac{\sqrt{5} \left(-8 \sin\left(\frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2} + \theta_{23}\right) + 12 \sin\left(\frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\zeta_{2(3)}^1}{2} + \frac{3\theta_{12}}{2} + \theta_{23}\right) - 12 \sin\left(-\frac{3\zeta_{3(1)}^0}{2} - \frac{3\zeta_{2(1)}^1}{2} + \frac{\zeta_{2(3)}^1}{2} - \frac{3\theta_{12}}{2} - \theta_{23}\right) + 40\right)}{40}$$

$$T_{2,6} = \frac{\sqrt{5} \left(-2 \sin\left(\frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2}\right) + 3 \sin\left(\frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\zeta_{2(3)}^1}{2} + \frac{3\theta_{12}}{2}\right) - 3 \sin\left(\frac{3\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2}\right) + 10\right)}{10}$$

$$T_{3,6} = \frac{\sqrt{5} \left(-2 \sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2} + \frac{\theta_{31}}{2}\right) + 3 \sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(3)}^1}{2} + \frac{3\theta_{12}}{2} + \frac{\theta_{31}}{2}\right) - 3 \sin\left(\frac{3\zeta_{1(2)}^0}{2} + \frac{3\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2} - \frac{\theta_{31}}{2}\right) + 10\right)}{10}$$

$$T_{4,6} = \frac{3 \sin\left(-\frac{\zeta_{1(2)}^0}{2} - \frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} + \frac{3\theta_{31}}{2}\right)}{5} + \frac{2 \sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2} + \frac{\theta_{31}}{2}\right)}{5} + \frac{9 \sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2} - \frac{\theta_{31}}{2}\right)}{5}$$

$$T_{5,6} = \frac{\sqrt{5} \cdot (3 \cos(2\theta_{12}) + 1)}{10},$$

$$T_{2,1} = T_{1,2},$$

$$T_{3,1} = T_{1,3}.$$

```
1 printalign(T3h+[1])
```

$$T_{5,1} = T_{1,5},$$

s5h+ =

```
Setup((two_h1 = 1, two_h2 = 0, two_h3 = 0, two_h0 = 5), [DecayChain{var"#15#17"{Sym}, Rec
```

```
1 s5h+ = Setup(chains, "5/2+" => final_state)
```

$$T_{6,2} = T_{2,6},$$

```
1 @time T5h+ = intensity_matrix(s5h+);
```

735.694538 seconds (1.08 M allocations: 34.103 MiB)

$$T_{6,4} = T_{4,6},$$

$$I_{5/2+} = \begin{pmatrix} R_{1,3/2}^{\rho(770)0} \\ R_{2,1/2}^{f_0} \\ R_{3,1/2}^{\Sigma(2455)0} \\ R_{1,3/2}^{\Sigma(2520)0} \\ R_{3,1/2}^{\Sigma(2455)++} \\ R_{1,3/2}^{\Sigma(2520)++} \end{pmatrix}^{\dagger} [T] \begin{pmatrix} R_{1,3/2}^{\rho(770)0} \\ R_{2,1/2}^{f_0} \\ R_{3,1/2}^{\Sigma(2455)0} \\ R_{1,3/2}^{\Sigma(2520)0} \\ R_{3,1/2}^{\Sigma(2455)++} \\ R_{1,3/2}^{\Sigma(2520)++} \end{pmatrix}$$

```
1 latexintensity(T5h+[2], "5/2+") |> Markdown.parse
```


$$\begin{aligned}
T_{1,1} &= \frac{12}{5} - \frac{3 \sin^2(\theta_{23})}{5}, \\
T_{2,2} &= 2, \\
T_{3,3} &= 2, \\
T_{4,4} &= \frac{3 \cos(2\theta_{31})}{10} + \frac{21}{10}, \\
T_{5,5} &= 2, \\
T_{6,6} &= \frac{3 \cos(2\theta_{12})}{10} + \frac{21}{10}, \\
T_{1,2} &= \frac{2\sqrt{30} \cos(\theta_{23})}{5}, \\
T_{1,3} &= \frac{\sqrt{30} \left(\cos\left(\frac{\zeta_{1(2)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} - \theta_{23} + \frac{\theta_{31}}{2}\right) + \cos\left(\frac{\zeta_{1(2)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \theta_{23} + \frac{\theta_{31}}{2}\right) + \cos\left(\frac{3\zeta_{1(2)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} + \theta_{23} - \frac{\theta_{31}}{2}\right) \right)}{20}, \\
T_{2,3} &= \frac{\cos\left(\frac{\zeta_{1(2)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\theta_{31}}{2}\right)}{2} + \frac{\cos\left(\frac{3\zeta_{1(2)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\theta_{31}}{2}\right)}{4} + \frac{5 \cos\left(\frac{5\zeta_{1(2)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\theta_{31}}{2}\right)}{4}, \\
T_{1,4} &= \frac{3 \cos\left(-\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} + \theta_{23} + \frac{3\theta_{31}}{2}\right)}{20} + \frac{3 \cos\left(\frac{\zeta_{1(2)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} - \theta_{23} + \frac{\theta_{31}}{2}\right)}{20} + \frac{3 \cos\left(\frac{\zeta_{1(2)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \theta_{23} - \frac{\theta_{31}}{2}\right)}{20}, \\
T_{2,4} &= \frac{\sqrt{30} \cdot \left(5 \cos\left(-\frac{5\zeta_{1(2)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} + \frac{3\theta_{31}}{2}\right) + \cos\left(-\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} + \frac{3\theta_{31}}{2}\right) + \cos\left(\frac{\zeta_{1(2)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\theta_{31}}{2}\right) + \cos\left(\frac{\zeta_{1(2)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} - \theta_{23} + \frac{\theta_{31}}{2}\right) \right)}{20}, \\
T_{3,4} &= \frac{\sqrt{30} \cdot (6 \cos(2\theta_{31}) + 2)}{20}, \\
T_{1,5} &= \sqrt{30} \left(\frac{\sin\left(\frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2}\right) \cos(\theta_{23})}{10} - \frac{\sin\left(-\frac{5\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} + \theta_{23}\right)}{4} - \frac{\sin\left(-\frac{3\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} + \theta_{23}\right)}{4} \right), \\
T_{2,5} &= \frac{\sin\left(\frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2}\right)}{2} - \frac{\sin\left(\frac{3\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2}\right)}{4} + \frac{5 \sin\left(\frac{5\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2}\right)}{4}, \\
T_{3,5} &= \frac{\sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2} + \frac{\theta_{31}}{2}\right)}{2} - \frac{\sin\left(\frac{3\zeta_{1(2)}^0}{2} + \frac{3\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} - \frac{\theta_{31}}{2}\right)}{4} + \frac{5 \sin\left(\frac{5\zeta_{1(2)}^0}{2} + \frac{5\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2} + \frac{\theta_{31}}{2}\right)}{4}, \\
T_{4,5} &= \frac{\sqrt{30} \left(-5 \sin\left(-\frac{5\zeta_{1(2)}^0}{2} - \frac{5\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} + \frac{3\theta_{31}}{2}\right) - \sin\left(-\frac{\zeta_{1(2)}^0}{2} - \frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} + \frac{3\theta_{31}}{2}\right) + \sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2} + \frac{\theta_{31}}{2}\right) \right)}{20}, \\
T_{1,6} &= -\frac{3 \sin\left(-\frac{\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} + \theta_{23}\right)}{20} + \frac{3 \sin\left(\frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2} + \theta_{23}\right)}{20} + \frac{3 \sin\left(\frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2} + \theta_{23}\right)}{20}, \\
T_{2,6} &= \frac{\sqrt{30} \left(\sin\left(\frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2}\right) + \sin\left(\frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(1)}^1}{2} - \frac{\zeta_{2(3)}^1}{2} + \frac{3\theta_{12}}{2}\right) - \sin\left(\frac{3\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(1)}^1}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2}\right) \right)}{20}, \\
T_{3,6} &= \frac{\sqrt{30} \left(\sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2} + \frac{\theta_{31}}{2}\right) + \sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(3)}^1}{2} + \frac{3\theta_{12}}{2} + \frac{\theta_{31}}{2}\right) - \sin\left(\frac{3\zeta_{1(2)}^0}{2} + \frac{3\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2} + \frac{\theta_{31}}{2}\right) \right)}{20}, \\
T_{4,6} &= -\frac{3 \sin\left(-\frac{\zeta_{1(2)}^0}{2} - \frac{\zeta_{3(1)}^0}{2} + \frac{\zeta_{2(3)}^1}{2} + \frac{\theta_{12}}{2} + \frac{3\theta_{31}}{2}\right)}{20} + \frac{3 \sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2} + \frac{\theta_{31}}{2}\right)}{20} + \frac{3 \sin\left(\frac{\zeta_{1(2)}^0}{2} + \frac{\zeta_{3(1)}^0}{2} - \frac{\zeta_{2(3)}^1}{2} - \frac{\theta_{12}}{2} + \frac{\theta_{31}}{2}\right)}{20}, \\
T_{5,6} &= \frac{\sqrt{30} \cdot (6 \cos(2\theta_{31}) + 2)}{20},
\end{aligned}$$

```
1 printalign(T5h+[1])
```

$T_{2,1} = T_{1,2},$

Save to file

$T_{6,1} = T_{1,6},$

```
1 Pkg.add("JSON")
```

```
Resolving package versions...
Updating `C:\Users\Mikhasenko\AppData\Local\Temp\jl_9dSubJ\Project.toml`
[682c06a0] + JSON v0.21.4
No Changes to `C:\Users\Mikhasenko\AppData\Local\Temp\jl_9dSubJ\Manifest.toml`
```

$T_{5,4} = T_{4,5},$

```
1 using JSON
```

$T_{6,5} = T_{5,6},$

wrap2dict (generic function with 1 method)

```
1 wrap2dict(TV) = Dict(
2     "intensity" => latexintensity(T1h-[2]),
3     "matrix" => latexalign(T1h-[1]))
```

writejson (generic function with 1 method)

```
1 function writejson(path, obj)
2     open(path, "w") do io
3         JSON.print(io, obj, 4)
4     end
5 end
```

computation_result =

Dict("3/2+" => Dict("intensity" => "``math\n\\small\n\\begin{equation}\n{\\Large I_{}"

```
1 computation_result = Dict("1/2-"=>wrap2dict(T1h-),
2     "3/2-"=>wrap2dict(T3h-),
3     "3/2+"=>wrap2dict(T3h+),
4     "5/2+"=>wrap2dict(T5h+))
```

```
1 writejson(joinpath(@__DIR__, "LcXX2Lcpipi.json"), computation_result)
```

Simplifications

sign_replacements =

$$[\zeta_{2(1)}^0, \zeta_{3(2)}^0, \zeta_{1(3)}^0, \zeta_{3(1)}^1, \zeta_{1(2)}^1, \zeta_{3(2)}^1, \zeta_{1(2)}^2, \zeta_{1(3)}^2]$$

$$\Rightarrow -\zeta_{1(2)}^0 \Rightarrow -\zeta_{2(3)}^0 \Rightarrow -\zeta_{3(1)}^0 \Rightarrow -\zeta_{1(3)}^1 \Rightarrow -\zeta_{2(1)}^1 \Rightarrow -\zeta_{2(3)}^1 \Rightarrow -\zeta_{2(1)}^2 \Rightarrow -\zeta_{3(1)}^2$$

```

1 sign_replacements = let
2   all_rotations = [(i,j,k) for (i,j,k)
3     in Iterators.product(1:3,1:3,0:3) if i!=j]
4   negative_rotations = filter(all_rotations) do (i,j,k)
5     _wr = wr(i,j,k)
6     !ispositive(_wr)
7   end
8   map(negative_rotations) do (i,j,k)
9     symζ(wr(i,j,k)) => -symζ(wr(j,i,k))
10  end
11 end

```

uncombine_replacement = [$\zeta_{2(3)}^1$, $\zeta_{3(1)}^2$, $\zeta_{1(2)}^3$]

$$\Rightarrow \zeta_{1(3)}^1 + \zeta_{2(1)}^1 \Rightarrow \zeta_{2(1)}^2 + \zeta_{3(2)}^2 \Rightarrow \zeta_{1(3)}^3 + \zeta_{3(2)}^3$$

```

1 uncombine_replacement = [
2   symζ(wr(2,3,1)) => symζ(wr(2,1,1))+symζ(wr(1,3,1)),
3   symζ(wr(3,1,2)) => symζ(wr(3,2,2))+symζ(wr(2,1,2)),
4   symζ(wr(1,2,3)) => symζ(wr(1,3,3))+symζ(wr(3,2,3))]

```

combine_replacements = [$\zeta_{1(3)}^1$, $\zeta_{2(1)}^2$, $\zeta_{3(2)}^3$]

$$\Rightarrow -\zeta_{2(1)}^1 + \zeta_{2(3)}^1 \Rightarrow \zeta_{3(1)}^2 - \zeta_{3(2)}^2 \Rightarrow \zeta_{1(2)}^3 - \zeta_{1(3)}^3$$

```

1 combine_replacements = [
2   symζ(wr(1,3,1)) => symζ(wr(2,3,1)) - symζ(wr(2,1,1)),
3   symζ(wr(2,1,2)) => symζ(wr(3,1,2)) - symζ(wr(3,2,2)),
4   symζ(wr(3,2,3)) => symζ(wr(1,2,3)) - symζ(wr(1,3,3))]

```

unsafe_uncombine_replacement =

[$\zeta_{2(3)}^0$, $\zeta_{3(1)}^0$, $\zeta_{1(2)}^0$]

$$\Rightarrow -\zeta_{1(2)}^0 - \zeta_{3(1)}^0 + 2\pi \Rightarrow -\zeta_{1(2)}^0 - \zeta_{2(3)}^0 + 2\pi \Rightarrow -\zeta_{2(3)}^0 - \zeta_{3(1)}^0 + 2\pi$$

```

1 unsafe_uncombine_replacement = [
2   symζ(wr(2,3,0)) => 2PI-symζ(wr(3,1,0))-symζ(wr(1,2,0)),
3   symζ(wr(3,1,0)) => 2PI-symζ(wr(1,2,0))-symζ(wr(2,3,0)),
4   symζ(wr(1,2,0)) => 2PI-symζ(wr(2,3,0))-symζ(wr(3,1,0))
5 ]

```

