

# Gametic Selection, Sex Ratio Bias, and Transitions Between Sex Determination Systems

Michael F Scott\*<sup>1</sup> and Matthew M Osmond\*<sup>2</sup>, and Sarah P Otto<sup>2</sup>

\* These authors contributed equally to this work

<sup>1</sup> Department of Botany, University of British Columbia, #3529 - 6270 University Boulevard, Vancouver, BC, Canada V6T 1Z4

<sup>2</sup> Department of Zoology, University of British Columbia, #4200 - 6270 University Boulevard, Vancouver, BC, Canada V6T 1Z4

email: mfcscott@biodiversity.ubc.ca, mmosmond@zoology.ubc.ca

Contributions:

## Abstract

Sex determination systems are remarkably dynamic; many studied taxa display transitions of sex-determining genes between chromosomes or the evolution of new sex-determining systems. Here, we utilize population genetic models to study the spread of novel sex-determining systems when there is haploid gametic selection, e.g., pollen or sperm competition. Haploid selected loci experience a form of sex-specific selection (because gametic competition occurs predominantly among haploids produced by males) and can cause sex ratios at birth to become biased (because sex ratios are determined by the fertilization success of X- versus Y-bearing pollen/sperm). We find that the evolution of sex determination systems where mothers determine sex at birth (e.g., environmental sex determination where sex is determined at birth) is influenced by classic Fisherian sex ratio selection. (Maybe not true???) However, notably, we find that the spread of new genetic sex determination systems is not affected by sex ratio biases that are caused by gametic selection because sex ratios become biased after parental provisioning has occurred (even if pollen/sperm competition occurs within the mother). In addition, we find that linkage of an ancestral sex chromosome to a locus under haploid selection can favour transitions between male and female heterogamety (e.g., XY to ZW), which is not the case for any forms of diploid sex specific selection (e.g., sexually antagonistic selection). During these transitions, new sex-determining alleles spread despite breaking up favourable associations that build up between ancestral sex-determining loci and selected loci, reducing population mean fitness. Furthermore, a period of selection among haploids can favour the stable maintenance of polymorphic sex determination systems. Thus, our models offer several new insights to be explored as information about sex determination in non-model taxa accumulates.

## Introduction

Animals and angiosperms exhibit extremely diverse sex determination systems, as reviewed by Bull (1983), Charlesworth and Mank (2010), Beukeboom and Perrin (2014), Bachtrog et al. (2014). Among species with genetic sex determination of diploid sexes, some taxa have heterogametic males (XY) and homogametic females (XX), including mammals and most dioecious plants (Ming et al. 2011); whereas other taxa have homogametic males (ZZ) and heterogametic females (ZW), including Lepidoptera and birds. Within several taxa, the chromosome that harbours the master sex-determining region changes. For example, transitions of the master sex-determining gene between chromosomes or the evolution of new master sex-determining genes have occurred in Salmonids (Li et al. 2011, Yano et al. 2012), Diptera (Vicoso and Bachtrog 2015), and *Oryzias* (Myosho et al. 2012). In addition, many gonochoric/dioecious clades with genetic sex determination exhibit transitions between male (XY) and female (ZW) heterogamety, including lizards (Ezaz et al. 2009), eight of 26 teleost fish families (Mank et al. 2006), true fruit flies (Tephritids, Vicoso and Bachtrog 2015), amphibians (Hillis and Green 1990), the angiosperm genus *Silene* (Slancarova et al. 2013), Coleoptera and Hemiptera (Beukeboom and Perrin 2014, plate 2). Indeed, in some cases, both male and female heterogametic sex determination systems can be found in the same species, as exhibited by some cichlid species (Ser et al. 2010) and *Rana rugosa* (Ogata et al. 2007). In addition, multiple transitions have occurred between genetic and environmental sex determination systems, e.g., in reptiles and fishes (Conover and Heins 1987, Mank et al. 2006, Pokorná and Kratochvíl 2009, Ezaz et al. 2009, Pen et al. 2010, Holleley et al. 2015).

Predominant theories in which new sex determination systems are favoured by selection involve fitness differences between sexes (e.g., sexually antagonistic selection) or sex ratio selection. van Doorn and Kirkpatrick (2007; 2010) show that new sex determination loci can be favoured if they arise in close linkage with a locus that experiences sexual antagonism. For ex-

ample, linkage allows favourable associations to build up between a male-beneficial allele and a neo-Y chromosome. Such associations can favour a new master sex-determining gene on a new chromosome (van Doorn and Kirkpatrick 2007) and can also favour a transition between male and female heterogamety (e.g., a ZW to XY transition, van Doorn and Kirkpatrick 2010). However, any sexually-antagonistic loci that are linked to the ancestral sex-determination locus will develop similar, favourable associations and select against the spread of a new sex-determination system.

It has been suggested that sex ratio selection could be a particularly important force driving transitions between sex-determining systems (Beukeboom and Perrin 2014, Chapter 7). For example, flexible sex determination systems may be favoured in order to exploit local environmental conditions that are optimal for males or females, which creates locally biased sex ratios (Charnov and Bull 1977, Werren and Taylor 1984, Pen et al. 2010). In addition, feminizing mutations may invade when female biased sex ratios are favoured due to selection among demes (Wilson and Colwell 1981, Vuilleumier et al. 2007). In other situations, sex ratio selection may favour transitions in order to restore equal sex ratios. For example, Kozielska et al. (2010) consider systems in which the ancestral sex chromosomes experience meiotic drive (e.g., where driving X or Y chromosomes are inherited disproportionately often), which causes sex ratios to become biased (Hamilton 1967). They find that new, unlinked sex-determining loci (masculinizing or feminizing mutations, i.e., neo-Y or neo-W loci) can then spread, restoring an even sex ratio.

Here, we use mathematical models to find the conditions under which new sex determination systems are favoured by selection where we include a period of selection among haploid gametes/gametophytes. Selection among haploid genotypes is thought to occur primarily among pollen/sperm, which can compete whenever there are more pollen/sperm than required for fertilization (Mulcahy et al. 1996, Joseph and Kirkpatrick 2004). Haploid selection

may be particularly common in plants, in which 60-70% of all genes are expressed in the male gametophyte and these genes exhibit stronger signatures of selection than random genes (Borg et al. 2009, Arunkumar et al. 2013, Gossmann et al. 2014). In addition, artificial selection pressures applied to male gametophytes cause the frequency of resistant alleles to increase (e.g., Hormaza and Herrero 1996, Ravikumar et al. 2003, Hedhly et al. 2004, Clarke et al. 2004). A smaller (but non-negligible) proportion of genes are thought to be expressed and selected in animal sperm, although precise estimates are uncertain (Zheng et al. 2001, Joseph and Kirkpatrick 2004, Vibranovski et al. 2010).

There are various ways in which a period of haploid selection could influence transitions between sex determination systems. Firstly, if we assume that haploid selection at any particular locus predominantly occurs in one sex (e.g., pollen/sperm competition), then such loci experience a form of sex-specific selection. In this respect, we might expect that haploid selection might affect transitions between sex determination systems in a similar manner to sex-specific diploid selection (as explored by van Doorn and Kirkpatrick 2007; 2010). That is, new masculinizing mutations (neo-Y chromosomes) could be favoured via linkage associations with alleles that are beneficial in pollen/sperm. However, sex ratios can also become biased if there is linkage between the sex-determining region and a locus that harbours genetic variation in haploid fitness. For example, differences in fitness between X- and Y-bearing pollen tubes can cause the sex ratio among seeds to become biased when there is pollen competition (Lloyd 1974, Conn and Blum 1981, Stehlik and Barrett 2005; 2006, Field et al. 2012; 2013). It is not immediately clear how the spread of new sex determination systems would be influenced by the combination of sex ratio biases and favourable associations between haploid selected loci and sex-determining regions.

Surprisingly, our models show that haploid selection influences the evolution of new sex determination systems in a way that is distinct from both

diploid sex-specific selection and sex ratio selection. We find that new genetic sex determination systems are not affected by any sex ratio biases caused by associations between sex-determining regions and haploid selected loci. In addition, we find that associations that build up between an ancestral sex-determining locus and a haploid-selected locus can favour transitions between male and female heterogamety (e.g., a neo-Z/neo-W locus spreads in an ancestrally XY system), despite the fact that these ancestral associations were built up by selection. This does not occur in models that do not include haploid selection.

NOTE RE: DRIVE. I expect drive (that occurs specifically in one sex, e.g., during spermatogenesis) to behave almost exactly like haploid selection. That is, I think that a XY-linked driver that is maintained by selection (e.g., because it causes sterility when homozygous, which is common in known drive systems) will only favour invasion of a more tightly linked neo-Y (worsening sex ratio biases) and could favour invasion of a neo-W. This may run counter to generic expectations from new sex chromosome systems evolving to balance the sex ratio. So, do you think it would significantly enhance the paper to model drive explicitly or just discuss it as being similar???

FOR RESULTS?

FROM PREVIOUS PAPER: The maintenance of polymorphism at loci that experience sex specific selection in both haploid and diploid phases was considered by Immler et al. Immler et al. (2012), demonstrating that polymorphisms can be maintained by sexually antagonistic selection or overdominance as well as by conflicting selection pressures in haploids and diploids (haploid-diploid conflict or ploidy antagonistic selection) or a combination of these selective regimes.

## Model

We consider a three locus model. Locus **X** is the ancestral sex-determining region (SDR), with alleles  $X$  and  $Y$  (or  $Z$  and  $W$ ). Locus **A** is a region under selection, with alleles  $A$  and  $a$ . And locus **M** is a novel sex-determining region, with alleles  $M$  and  $m$ . With genotype  $MM$  at locus **M**, the sex of a zygote is determined by the genotype at locus **X** ( $XX$  become females and  $XY$  males, or  $ZW$  become females and  $ZZ$  males). With at least one  $m$  allele at locus **M**, a zygote develops as a female with probability  $k$  and as a male with probability  $1 - k$ . With  $k = 0$ , the novel sex determiner is a masculinizer (e.g., a neo-Y chromosome) and with  $k = 1$  the novel sex determiner is a feminizer (e.g., a neo-W chromosome). With intermediate  $k$ , locus **M** is interpreted as an offspring controlled environmental sex-determining region. Finally, we also analyze a model of maternally-controlled environmental sex-determination, where mothers with at least one  $m$  allele produce daughters with probability  $k$ .

The life-cycle begins with competition between haploid gametes/gametophytes from each sex, where selection depends on the sex of the diploid they came from and their allele at the **A** locus. Gametes/gametophytes from males then pair randomly with gametes/gametophytes from females. The resulting zygotes develop as males or females, depending on their genotypes at the **X** and **M** loci (or the **M** genotype of their mother in the case of maternal control). Diploids compete with others of the same sex, where selection depends on the sex of the individual and its genotype at the **A** locus. This is followed by meiosis with recombination and sex-specific meiotic drive. Recombination occurs between loci **X** and **A** with probability  $r$ , between loci **A** and **M** with probability  $R$ , and between loci **X** and **M** with probability  $\chi$ . Any order of the loci can be modelled with appropriate choices of  $r$ ,  $R$ , and  $\chi$ . We model meiotic drive at the viability locus only; individuals of sex  $i$  who are heterozygous at the **A** locus produce haploids bearing allele  $A$  with probability  $\alpha^i$ . We track the frequency of haploid genotypes produced by each sex from

one generation to the next (recursion equations in Sup. Mat.).

With allele  $M$  fixed at the **M** locus, sex is determined by locus **X** and an equilibrium is reached at locus **A**. We then examine the ability of a rare, novel sex-determiner,  $m$ , to increase in frequency from this equilibrium. Numerical simulations are used to examine when an invading mutation goes to fixation.

## Results

### Resident equilibrium and stability

With allele  $M$  fixed, we follow the dynamics of the frequency of  $A$  in gametes/gametophytes from the homogametic sex (e.g., in eggs from an XX female,  $p_X^f$ ), the frequencies of  $A$  in the two types of gametes/gametophytes from the heterogametic sex (e.g., in sperm from an XY male that are X-bearing,  $p_X^m$ , and Y-bearing,  $p_Y^m$ ), and the frequencies of the sex-determining factors in gametes/gametophytes from the heterogametic sex (e.g., the frequency of Y in sperm,  $q$ ). Assuming selection and meiotic drive are weak relative to recombination, the differences in the frequencies of  $A$  in each type of gamete/gametophyte are small, as is the bias in the sex-determining factor from the heterogametic sex, and we can solve for the mean frequency of  $A$  across all types ( $p_A$ ), the difference in the frequencies of  $A$  between two of the three types, and the bias in the frequency of the sex-determining factor, to first order in selection. Linear stability analysis can then be used to determine the stability of this equilibrium. Without haploid selection or meiotic drive our results reduce to those of van Doorn and Kirkpatrick (2007). However, with haploid selection or meiotic drive a stable polymorphism at locus **A** no longer requires sexually antagonistic selection.



## Sex chromosome turnover

The spread of a rare mutant  $m$  at the **M** locus in such a population is determined by the leading eigenvalue,  $\lambda$ , of the system described by the equations for the next generation frequency of haploid genotypes with the mutation. Below we present the results for invasion into an  $XY$  system assuming no competition among gametes/gametophytes from females, no meiotic drive in females, and linear arrangement **XAM**. Equivalent results for invasions into a  $ZW$  system can be derived by consistently switching the roles of males and females (van Doorn and Kirkpatrick 2010).

### Neo- $Y$ and neo- $W$

A rare, dominant neo- $Y$  ( $k = 0$ ) or neo- $W$  ( $k = 1$ ) is always expected to invade an ancestral  $XY$  system when the average growth rate of the mutant haplotypes ( $Am$  and  $am$ ) is positive,  $(g_A + g_a)/2 > 0$ . When the growth rates of the mutant haplotypes without recombination ( $R = 0$ ) are negative,  $g_i^* < 0 \forall i \in \{A, a\}$ , where  $g_i < g_i^*$ , the new sex-determining allele does not invade.

Otherwise, the new sex-determining allele increases in frequency on one **A** background and declines on the other, and invasion requires

$$R \left[ \frac{p_X^f w_a^m (1 - \alpha^m)}{g_A^*} + \frac{(1 - p_X^f) w_A^m \alpha^m}{g_a^*} \right] w_{Aa}^m < \nu^m, \quad (1)$$

for the neo- $Y$ , and

$$R \left[ \frac{(1 - \bar{p}^m) w_a^m}{g_A^*} + \frac{\bar{p}^m w_A^m}{g_a^*} \right] w_{Aa}^f < \nu^f \quad (2)$$

for the neo- $W$ . Here  $\bar{p}^m = (p_X^m + p_Y^m)/2$  is the average frequency of  $A$  in gametophytes produced by males,  $w_j^i$  is the relative viability fitness of sex  $i \in \{m, f\}$  depending on their haploid or diploid genotype at the **A** locus

(with  $j \in \{A, a, AA, Aa, aa\}$ ) and  $\nu^i = p_X^f p_{\mathbf{X}^i}^m w_a^m w_{AA}^i + p_X^f (1 - p_{\mathbf{X}^i}^m) w_a^m w_{Aa}^i + (1 - p_X^f) p_{\mathbf{X}^i}^m w_a^m w_{Aa}^i + (1 - p_X^f) (1 - p_{\mathbf{X}^i}^m) w_a^m w_{aa}^i$ , with  $\mathbf{X}^m = Y$  and  $\mathbf{X}^f = X$ , is the mean relative fitness of resident individuals of sex  $i$ . Although male meiotic drive does not explicitly appear in Equation 2, it does affect the average frequency of  $A$  in gametes/gametophytes from males,  $\bar{p}^m$ , and thus can play a role in neo-W invasion.

Equations (1) and (2) show that the new sex-determining allele is expected to invade for any recombination rate,  $R$ , when the net flow of double recombinants is from the less fit to the more fit  $\mathbf{A}$  background (making the terms inside the square brackets in Equations 1 and 2 negative). When the net flow of double recombinants is from the more fit to the less fit haplotype, the new sex-determining allele can still invade when the rate of recombination between it and the selected locus,  $R$ , is small enough.

Assuming weak selection and meiotic drive, we can explicitly solve for the invasion fitness of the new sex-determining allele,  $m$ , into the ancestral  $XY$  system, giving

$$\lambda_{m,XY} \approx 1 + V_A S_A C_m, \quad (3)$$

where  $V_A = p_A(1 - p_A)$  is the variance at the  $\mathbf{A}$  locus. We will consider haploid selection and meiotic drive separately. With haploid selection and no drive we have  $S_A = s^f (t^m)^2 / [rR (s^f + s^m)^2]$ , where  $s^f$  and  $s^m$  are the respective selection coefficients for  $A$  in diploid females and males,  $t^m$  is the selection coefficient for  $A$  in gametes/gametophytes from males, and we've assumed equal dominance coefficients in the two sexes. With drive and no haploid selection we replace  $t^m$  with  $\alpha^m - 1/2$ .

For the neo- $Y$  we have  $C_Y = (r - R)s^f$  with haploid selection and  $C_Y = 4(r - R)s^f$  with drive. The neo- $Y$  can therefore invade whenever it is in tighter linkage with the selected locus than the ancestral sex-determining locus,  $r > R$ , provided locus  $\mathbf{A}$  is polymorphic ( $V_A > 0$ ), there is diploid selection in females ( $s^f \neq 0$ ), and there is either haploid selection in gametes/gametophytes from males ( $t^m \neq 0$ ) or meiotic drive in males ( $\alpha^m \neq$

1/2). This reduces to the weak-linkage results of van Doorn and Kirkpatrick (2007) when we do not assume equal dominance coefficients in the two sexes, there is no haploid selection or meiotic drive, and there is free recombination between locus **A** and one of the sex-determining regions.

For the neo-*W* we have  $C_W = [(2r(1 - R) - R)s^f + (1 - 2r)Rs^m]/2$  with haploid selection and  $C_W = 4[r(1 - 2R)s^f + (1 - 2r)Rs^m]/2$  with drive. In this case, even when the novel sex-determining locus is in looser linkage with the selected locus than the ancestral sex-determining locus is,  $r < R$ , a neo-*W* can invade. For example,  $R = 1/2$  the neo-*W* invades if there is any linkage between the ancestral sex-determining and selected loci ( $r < 1/2$ ), there is selection among gametes/gametophytes in males ( $t^m \neq 0$ ), and there is selection for or against *A* in both males and females ( $s^m s^f > 0$ ) that is stronger in males than in females ( $|s^m| > |s^f|$ ). With meiotic drive and  $R = 1/2$ , all that is required for neo-*W* invasion is  $r < 1/2$  and  $s^m s^f > 0$ . Our results reduce to the weak-linkage results of (van Doorn and Kirkpatrick 2010, Equation 3) when we do not assume equal dominance coefficients in the two sexes and there is no haploid selection or meiotic drive.

### Offspring-controlled neo-ESD

The growth rate of a rare, dominant offspring-controlled neo-ESD region that produces males or females with equal probability ( $k = 1/2$ ) is

$$\lambda_{ESD,XY} \approx 1 + \frac{1}{2} \frac{(\lambda_{Y,XY} - 1) + (\lambda_{W,XY} - 1)}{2} \Big|_{R=1/2} \quad (4)$$

Thus with  $k = 1/2$  the neo-ESD gets half of the advantages of a neo-*W* and half that of a neo-*Y*, but only has an effect one half of the time (the other half of the time it produces the same sex as the ancestral system would have). Recombination between the selected locus and the novel sex-determining locus, *R*, doesn't enter into the  $k = 1/2$  results because sex is essentially randomized each generation, preventing associations from building up between allele

A and sex.

## Maternally-controlled neo-ESD

One might think that when the sex of zygotes is under the control of mothers, there would be strong selection to balance the sex ratio among zygotes. However, we find that, as with offspring control, assuming weak selection the invasion fitness of a sex-determiner that is maternally controlled can be written

$$\lambda_{k,XY} \approx 1 + V_A S_A C_k, \quad (5)$$

where  $C_k$  is a term that depends on  $k$ . Of particular interest is  $k = 1/2$  (i.e., when the mother perfectly balances the sex ratio of her offspring). When both recombination rates are small we have  $C_{1/2} \approx R(s^m - s^f)/8 = \lim_{r \rightarrow 0} C_W/4$ . This implies that, at least under tight linkage, the invasion of maternal ESD is independent of  $R$  (because  $S_A \propto R^{-1}$ ) and can invade whenever a neo- $W$  can.

## Discussion

2 Brief results summary.

DRAFT (improve): In Úbeda et al. (2015), the new sex determining locus  
4 spreads because it arises in linkage with a locus that experiences drive. They  
assume that drive occurs predominantly in one sex, e.g., during spermatogenesis  
6 or a 'killer' sperm. A driving allele is maintained at an intermediate  
frequency by selection, e.g., because it causes male sterility when homozygous  
8 (because all male sperm are killed). Y chromosomes that arise in linkage with  
the driving allele spread because they allow drive to occur more often, thus  
10 genetic sex determination with a sex ratio bias evolves. Thus Úbeda et al.  
(2015) also find that genetic sex determiners can invade, despite causing sex  
12 ratios to become biased. Finally, they show that autosomal 'restorers' that

negate the effects of meiotic drive can invade and restore an equal sex ratio.

14 OTHER RESTORERS. Indeed, alleles that negate the effect of sex-linked  
meiotic drivers and restore equal sex ratios have been identified (Stalker  
16 1961, Smith 1975). A similar process occurs with cytoplasmic male sterility  
alleles (that cause biased sex ratios) and nuclear ‘restorer’ genotypes (Frank  
18 1989). When sex ratio bias occurs due to haploid selection, a natural class  
of sex ratio ‘restorers’ exist because haploid selection often occurs in a con-  
20 text that is determined by the diploid parents. For example, the intensity of  
pollen competition can be manipulated by altering style length (Travers and  
22 Shea 2001, Lankinen and Skogsmyr 2001, Ruane 2009), delaying stigma re-  
ceptivity (Galen et al. 1986, Lankinen and Madjidian 2011) and/or delaying  
24 pollen tube growth in the pistil (Herrero 2003). Where the X and Y have  
fitness differences, Hough et al. (2013) and Otto et al. (2015) demonstrated  
26 that mothers should generally evolve to balance sex ratios by reducing the  
intensity of haploid competition.

28 FROM THESIS: However, reducing competition among haploids also re-  
duces the potential for harmful deleterious mutations to be purged. When  
30 deleterious mutations are included, the optimal intensity of haploid selection  
can reflect a balance between maximizing offspring fitness and equalizing sex  
32 ratios.

As part of a collaborative project (Otto et al. 2015), I considered the  
34 evolution of the haploid ‘selective arena’ in cases where the X chromosome  
harbours a polymorphism that affects haploid fitness. Mothers again primar-  
36 ily evolve to restore equal sex ratios. However, modifying haploid selection  
also affects the X-linked genotypes that are inherited by offspring. Specif-  
38 ically, increasing the intensity of haploid selection increases the proportion  
of daughters (all progeny of X-bearing sperm/pollen are female) that inherit  
40 the allele with high haploid fitness. If this allele has high fitness in daugh-  
ters, mothers can be selected to increase the intensity of haploid selection;  
42 otherwise, decreased selection among haploids is favoured. Thus, because

altering haploid selection intensity affects the alleles that are inherited by  
44 daughters, mothers can favour slightly biased sex ratios. In addition, I found  
that stronger sex ratio biases can be favoured by paternal manipulations of  
46 the haploid ‘selective arena’ because fathers are strongly selected to maximize  
their own siring success (above selection to equalize the sex ratio).

48 **Discuss patterns that might be looked for:**

In broadcast spawning animal species (e.g., corals, many fish) and species  
50 where sperm typically requires greater longevity, haploid selection may be  
stronger because transcripts shared during spermatogenesis may become de-  
52 pleted (Immler et al. 2014). **also, mating systems (e.g., fewer alleles are  
available during haploid competition in monogamous species), selfing rates,  
54 and estimates of pollen limitation could be used as indicators of the intensity  
of haploid selection**

56 We have results where polygenic sex determination is sometimes stable, may  
be worth mentioning:

58 “Polygenic sex determination has been reported in many plants (e.g.  
Shannon & Holsinger 2007), fishes (Vandeputte et al. 2007; Ser et al. 2010;  
60 Liew et al. 2012), crustaceans (e.g. Battaglia 1958; Battaglia & Malesani  
1959; Voordouw & Anholt 2002), bivalves (Haley 1977; Saavedra et al. 1997),  
62 gastropods (Yusa 2007a,b), and polychaetes (Bacci 1965, 1978; Premoli et al.  
1996).” From Vuilleumier et al. 2007: “Polymorphism for sex-determining  
64 genes within or among populations has been reported in many species in-  
cluding houseflies, midges, woodlice, platyfish, cichlid fish, and frogs (Gor-  
66 don, 1944; Kallman, 1970; Thomp-son, 1971; Macdonald, 1978; Bull, 1983;  
Rigaud et al., 1997; Caubet et al., 2000; Lande et al., 2001; Ogataet al.,  
68 2003; Lee et al., 2004; Mank et al., 2006).”

(Check with Jim Bull that it’s ok before including this speculation:) Fi-  
70 nally, Hamilton (1967) pointed out that biased sex ratios can affect popu-  
lation size because the number of offspring in each generation is typically  
72 determined by the number of females. Population density can, in turn, af-

fect the intensity of pollen/sperm competition in future generations because  
74 fewer males are available to donate pollen/sperm in a particular area. Thus,  
a feedback could occur between population densities and haploid selection,  
76 which has not yet been investigated.

## References

- 78 Arunkumar, R., E. B. Josephs, R. J. Williamson, and S. I. Wright. 2013.  
Pollen-specific, but not sperm-specific, genes show stronger purifying se-  
80 lection and higher rates of positive selection than sporophytic genes in  
*Capsella grandiflora*. *Molecular biology and evolution* 30:2475–2486.
- 82 Bachtrog, D., J. E. Mank, C. L. Peichel, M. Kirkpatrick, S. P. Otto, T.-L.  
Ashman, M. W. Hahn, J. Kitano, I. Mayrose, R. Ming, N. Perrin, L. Ross,  
84 N. Valenzuela, J. C. Vamasi, and Tree of Sex Consortium. 2014. Sex  
determination: why so many ways of doing it? *PLoS Biol* 12:e1001899.
- 86 Beukeboom, L. W., and N. Perrin. 2014. The evolution of sex determination.  
Oxford University Press, Oxford, UK.
- 88 Borg, M., L. Brownfield, and D. Twell. 2009. Male gametophyte development:  
a molecular perspective. *Journal of Experimental Botany* 60:1465–1478.
- 90 Bull, J. J. 1983. Evolution of sex determining mechanisms. The Benjamin  
Cummings Publishing Company.
- 92 Charlesworth, D., and J. E. Mank. 2010. The birds and the bees and the  
flowers and the trees: lessons from genetic mapping of sex determination  
94 in plants and animals. *Genetics* 186:9–31.
- Charnov, E. L., and J. Bull. 1977. When is sex environmentally determined?  
96 *Nature* 266:828–830.

- Clarke, H. J., T. N. Khan, and K. H. M. Siddique. 2004. Pollen selection  
 98 for chilling tolerance at hybridisation leads to improved chickpea cultivars.  
 Euphytica 139:65–74.
- 100 Conn, J. S., and U. Blum. 1981. Sex ratio of *Rumex hastatulus*: the effect of  
 environmental factors and certation. Evolution 35:1108–1116.
- 102 Conover, D. O., and S. W. Heins. 1987. Adaptive variation in environmental  
 and genetic sex determination in a fish. Nature 326:496–498.
- 104 Ezaz, T., S. D. Sarre, and D. O’Meally. 2009. Sex chromosome evolution  
 in lizards: independent origins and rapid transitions. Cytogenetic and  
 106 Genome Research 127:249–260.
- Field, D. L., M. Pickup, and S. C. H. Barrett. 2012. The influence of polli-  
 108 nation intensity on fertilization success, progeny sex ratio, and fitness in a  
 wind-pollinated, dioecious plant. International Journal of Plant Sciences  
 110 173:184–191.
- . 2013. Comparative analyses of sex-ratio variation in dioecious flow-  
 112 ering plants. Evolution 67:661–672.
- Frank, S. A. 1989. The Evolutionary Dynamics of Cytoplasmic Male Sterility.  
 114 American Naturalist 133:345–376.
- Galen, C., J. A. Shykoff, and R. C. Plowright. 1986. Consequences of stigma  
 116 receptivity schedules for sexual selection in flowering plants. American  
 Naturalist pages 462–476.
- 118 Gossmann, T. I., M. W. Schmid, U. Grossniklaus, and K. J. Schmid. 2014.  
 Selection-driven evolution of sex-biased genes Is consistent with sexual  
 120 selection in *Arabidopsis thaliana*. Molecular biology and evolution 31:574–  
 583.
- 122 Hamilton, W. D. 1967. Extraordinary sex ratios. Science 156:477–488.



- Hedhly, A., J. I. Hormaza, and M. Herrero. 2004. Effect of temperature  
 124 on pollen tube kinetics and dynamics in sweet cherry, *Prunus avium*  
 (Rosaceae). American journal of botany 91:558–564.
- Herrero, M. 2003. Male and female synchrony and the regulation of mating  
 126 in flowering plants. Philosophical Transactions of the Royal Society B:  
 128 Biological Sciences 358:1019–1024.
- Hillis, D. M., and D. M. Green. 1990. Evolutionary changes of heterogametic  
 130 sex in the phylogenetic history of amphibians. Journal of Evolutionary  
 Biology 3:49–64.
- Holleley, C. E., D. O’Meally, S. D. Sarre, J. A. Marshall Graves, T. Ezaz,  
 132 K. Matsubara, B. Azad, X. Zhang, and A. Georges. 2015. Sex reversal  
 134 triggers the rapid transition from genetic to temperature-dependent sex.  
 Nature 523:79–82.
- Hormaza, J. I., and M. Herrero. 1996. Male gametophytic selection as a plant  
 136 breeding tool. Scientia horticultrae 65:321–333.
- Hough, J., S. Immler, S. Barrett, and S. P. Otto. 2013. Evolutionarily stable  
 138 sex ratios and mutation load. Evolution 7:1915–1925.
- Immler, S., G. Arnqvist, and S. P. Otto. 2012. Ploidally antagonistic selection  
 140 maintains stable genetic polymorphism. Evolution 66:55–65.
- Immler, S., C. Hotzy, G. Alavioon, E. Petersson, and G. Arnqvist. 2014.  
 142 Sperm variation within a single ejaculate affects offspring development in  
 144 Atlantic salmon. Biology letters 10:20131040.
- Joseph, S., and M. Kirkpatrick. 2004. Haploid selection in animals. Trends  
 146 in Ecology & Evolution 19:592–597.

- Kozielska, M., F. J. Weissing, L. W. Beukeboom, and I. Pen. 2010. Segregation distortion and the evolution of sex-determining mechanisms. *Heredity* 104:100–112.
- Lankinen, A., and J. A. Madjidian. 2011. Enhancing pollen competition by delaying stigma receptivity: Pollen deposition schedules affect siring ability, paternal diversity, and seed production in *Collinsia heterophylla* (Plantaginaceae). *American journal of botany* 98:1191–1200.
- Lankinen, A., and I. Skogsmyr. 2001. Evolution of pistil length as a choice mechanism for pollen quality. *Oikos* 92:81–90.
- Li, J., R. B. Phillips, A. S. Harwood, B. F. Koop, and W. S. Davidson. 2011. Identification of the Sex Chromosomes of Brown Trout (*Salmo trutta*) and Their Comparison with the Corresponding Chromosomes in Atlantic Salmon (*Salmo salar*) and Rainbow Trout (*Oncorhynchus mykiss*). *Cytogenetic and Genome Research* 133:25–33.
- Lloyd, D. G. 1974. Female-predominant sex ratios in angiosperms, vol. 32. *Heredity*.
- Mank, J. E., D. E. L. Promislow, and J. C. Avise. 2006. Evolution of alternative sexdetermining mechanisms in teleost fishes. *Biological Journal of the Linnean Society* 87:83–93.
- Ming, R., A. Bendahmane, and S. S. Renner. 2011. Sex chromosomes in land plants. [dx.doi.org](https://doi.org/10.1093/aob/mbr244) 62:485–514.
- Mulcahy, D. L., M. Sari-Gorla, and G. B. Mulcahy. 1996. Pollen selection - past, present and future. *Sexual Plant Reproduction* 9:353–356.
- Myosho, T., H. Otake, H. Masuyama, M. Matsuda, Y. Kuroki, A. Fujiyama, K. Naruse, S. Hamaguchi, and M. Sakaizumi. 2012. Tracing the Emergence of a Novel Sex-Determining Gene in Medaka, *Oryzias luzonensis*. *Genetics* 191:163–170.

- 174 Ogata, M., Y. Hasegawa, H. Ohtani, M. Mineyama, and I. Miura. 2007. The  
ZZ/ZW sex-determining mechanism originated twice and independently  
176 during evolution of the frog, *Rana rugosa*. *Heredity* 100:92–99.
- Otto, S. P., M. F. Scott, and S. Immler. 2015. Evolution of haploid selection  
178 in predominantly diploid organisms. *Proceedings of the National ...*
- Pen, I., T. Uller, B. Feldmeyer, A. Harts, G. M. While, and E. Wapstra.  
180 2010. Climate-driven population divergence in sex-determining systems.  
*Nature* 468:436–438.
- 182 Pokorná, M., and L. Kratochvíl. 2009. Phylogeny of sexdetermining mech-  
anisms in squamate reptiles: are sex chromosomes an evolutionary trap?  
184 *Zoological Journal of the ...* 156:168–183.
- Ravikumar, R. L., B. S. Patil, and P. M. Salimath. 2003. Drought tolerance in  
186 sorghum by pollen selection using osmotic stress. *Euphytica* 133:371–376.
- Ruane, L. G. 2009. Post-pollination processes and non-random mating among  
188 compatible mates. *Evolutionary Ecology Research* 11:1031–1051.
- Ser, J. R., R. B. Roberts, and T. D. Kocher. 2010. Multiple interacting loci  
190 control sex determination in lake Malawi cichlid fish. *Evolution* 64:486–  
501.
- 192 Slancarova, V., J. Zdanska, B. Janousek, M. Talianova, C. Zschach, J. Zlu-  
vova, J. Siroky, V. Kovacova, H. Blavet, J. Danihelka, B. Oxelman, A. Wid-  
194 mer, and B. Vyskot. 2013. Evolution of sex determination systems with  
heterogametic males and females in *Silene*. *Evolution* 67:3669–3677.
- 196 Smith, D. A. S. 1975. All-female broods in the polymorphic butterfly *Danaus*  
*chrysippus* L. and their ecological significance. *Heredity* 34:363–371.
- 198 Stalker, H. D. 1961. The Genetic Systems Modifying Meiotic Drive in  
*Drosophila Paramelanica*. *Genetics* .

- 200 Stehlik, I., and S. Barrett. 2005. Mechanisms governing sex-ratio variation  
in dioecious *Rumex nivalis*. *Evolution* 59:814–825.
- 202 Stehlik, I., and S. C. H. Barrett. 2006. Pollination intensity influences sex  
ratios in dioecious *Rumex nivalis*, a wind-pollinated plant. *Evolution*  
204 60:1207–1214.
- Travers, S. E., and K. Shea. 2001. Selection on pollen competitive ability in  
206 relation to stochastic factors influencing pollen deposition. *Evolutionary  
Ecology Research* 3:729–745.
- 208 Úbeda, F., M. M. Patten, and G. Wild. 2015. On the origin of sex chromo-  
somes from meiotic drive. *Proceedings of the Royal Society B: Biological  
210 Sciences* 282:20141932.
- van Doorn, G. S., and M. Kirkpatrick. 2007. Turnover of sex chromosomes  
212 induced by sexual conflict. *Nature* 449:909–912.
- . 2010. Transitions Between Male and Female Heterogamety Caused  
214 by Sex-Antagonistic Selection. *Genetics* 186:629–645.
- Vibranovski, M. D., D. S. Chalopin, H. F. Lopes, M. Long, and T. L. Karr.  
216 2010. Direct evidence for postmeiotic transcription during *Drosophila  
melanogaster* spermatogenesis. *Genetics* 186:431–433.
- 218 Vicoso, B., and D. Bachtrog. 2015. Numerous transitions of sex chromosomes  
in Diptera. *PLoS Biol* 13:e1002078.
- 220 Vuilleumier, S., R. Lande, J. J. M. van Alphen, and O. Seehausen. 2007.  
Invasion and fixation of sex-reversal genes. *Journal of Evolutionary Biology*  
222 20:913–920.
- Werren, J. H., and P. D. Taylor. 1984. The effects of population recruitment  
224 on sex ratio selection. *The American Naturalist* 124:143–148.

- Wilson, D. S., and R. K. Colwell. 1981. Evolution of sex ratio in structured  
226 demes. *Evolution* 35:882–897.
- Yano, A., B. Nicol, E. Jouanno, E. Quillet, A. Fostier, R. Guyomard,  
228 and Y. Guiguen. 2012. The sexually dimorphic on the Y-chromosome  
gene ( sdY) is a conserved male-specific Y-chromosome sequence in many  
230 salmonids. *Evolutionary Applications* 6:486–496.
- Zheng, Y., X. Deng, and P. A. Martin-DeLeon. 2001. Lack of sharing of  
232 Spam1 (Ph-20) among mouse spermatids and transmission ratio distortion.  
*Biology of Reproduction* 64:1730–1738.