

Gametic Selection, Sex Ratio Bias, and Transitions Between Sex Determination Systems

Michael F Scott*¹ and Matthew M Osmond*², and Sarah P Otto²

* These authors contributed equally to this work

¹ Department of Botany, University of British Columbia, #3529 - 6270 University Boulevard, Vancouver, BC, Canada V6T 1Z4

² Department of Zoology, University of British Columbia, #4200 - 6270 University Boulevard, Vancouver, BC, Canada V6T 1Z4

email: mfcscott@biodiversity.ubc.ca, mmosmond@zoology.ubc.ca

Contributions:

Abstract

Sex determination systems are remarkably dynamic; many studied taxa display transitions of sex-determining genes between chromosomes or the evolution of new sex-determining systems. Here, we utilize population genetic models to study the spread of novel sex-determining systems where we also include haploid gametic selection, e.g., pollen or sperm competition. Haploid selected loci experience a form of sex-specific selection (because gametic competition occurs predominantly among haploids produced by males) and can also cause sex ratios at birth to become biased (because sex ratios are determined by the fertilization success of X- versus Y-bearing pollen/sperm). We find that the evolution of sex determination systems where mothers determine sex at birth (e.g., environmental sex determination where sex is determined at birth) is influenced by classic Fisherian sex ratio selection. (Maybe not true???) However, notably, we find that the spread of new genetic sex determination systems is not affected by sex ratio biases that are caused by gametic selection because sex ratios become biased after parental provisioning has occurred (even if pollen/sperm competition occurs within the mother). In addition, we find that linkage of an ancestral sex chromosome to a locus under haploid selection can favour transitions between male and female heterogamety (e.g., XY to ZW), which is not the case for any forms of diploid sex specific selection (e.g., sexually antagonistic selection). During these transitions, new sex-determining alleles spread despite breaking up favourable associations that build up between ancestral sex-determining loci and selected loci, reducing population mean fitness. Furthermore, a period of selection among haploids can favour the stable maintenance of polymorphic sex determination systems. Thus, our models offer several new insights to be explored as information about sex determination in non-model taxa accumulates.

Introduction

Animals and angiosperms exhibit extremely diverse sex determination systems, reviewed in Bull (1983), Charlesworth and Mank (2010), Beukeboom and Perrin (2014), Bachtrog et al. (2014). Among species with genetic sex determination of diploid sexes, some taxa have heterogametic males (XY) and homogametic females (XX), including mammals and most dioecious plants (Ming et al. 2011); whereas other taxa have homogametic males (ZZ) and heterogametic females (ZW), including Lepidoptera and birds. Within several taxa, the chromosome that harbours the master sex-determining region changes. For example, transitions of the master sex-determining gene between chromosomes or the evolution of new master sex-determining genes have occurred in Salmonids (Li et al. 2011, Yano et al. 2012), Diptera (Vicoso and Bachtrog 2015), and *Oryzias* (Myosho et al. 2012). In addition, many gonochoric/dioecious clades with genetic sex determination exhibit transitions between male (XY) and female (ZW) heterogamety, including eight of 26 teleost fish families (Mank et al. 2006), true fruit flies (Tephritids, Vicoso and Bachtrog 2015), amphibians (Hillis and Green 1990), the angiosperm genus *Silene* (Slancarova et al. 2013), Coleoptera and Hemiptera (Beukeboom and Perrin 2014, plate 2). Indeed, in some cases, both male and female heterogametic sex determination systems can be found the same species, including cichlid species (Ser et al. 2010) and *Rana rugosa* (Ogata et al. 2007).

Depending on the prominence of transitions to ESD in the manuscript include something like (currently quoted):

“Transitions have repeatedly occurred between environmental sex determination and genotypic sex determination, as exemplified by the distribution of temperature sex determination among reptiles: either temperature or genes provide the initial trigger in closely related species (Ewert and Nelson 1991; Pokorna and Kratochvil 2009; Ezaz et al. 2009) or even conspecific populations (Pen et al. 2010). Similar situations are found in fishes (e.g.,

Conover and Heins 1978a).”

We have results where polygenic sex determination is sometimes stable:

“Polygenic sex determination has been reported in many plants (e.g. Shannon & Holsinger 2007), fishes (Vandeputte et al. 2007; Ser et al. 2010; Liew et al. 2012), crustaceans (e.g. Battaglia 1958; Battaglia & Malesani 1959; Voordouw & Anholt 2002), bivalves (Haley 1977; Saavedra et al. 1997), gastropods (Yusa 2007a,b), and polychaetes (Bacci 1965, 1978; Premoli et al. 1996).”

Description of sex ratio adjustment and sexual antagonism theories:

Predominant theories in which new sex determination systems are favoured by selection involve fitness differences between sexes (e.g., sexually antagonistic selection) or sex ratio selection. van Doorn and Kirkpatrick (2007; 2010) show that new sex determination loci can be favoured if they arise in close linkage with a locus that experiences sexual antagonism. For example, linkage allows favourable associations to build up between a male-beneficial allele and a neo-Y chromosome. Such associations can favour a new master sex-determining gene on a new chromosome (van Doorn and Kirkpatrick 2007) and can also favour a transition between male and female heterogamety (e.g., a ZW to XY transition, van Doorn and Kirkpatrick 2010). However, any sexually-antagonistic loci that are linked to the ancestral sex-determination locus will develop similar, favourable associations and select against the spread of a new sex-determination system.

Sex ratio selection might be a particularly important force driving transitions between sex-determining systems (Beukeboom and Perrin 2014, Chapter 7). For example, feminizing mutations may invade when female biased sex ratios are favoured due to interdemic selection (Wilson and Colwell 1981, Vuilleumier et al. 2007 CITE) or local mate competition (Hamilton 1967, Reinhold 1996, Werren and Taylor 1984, Pen et al. 2010 CITE). In other situations, sex ratio selection may favour transitions in order to restore equal sex ratios. For example, Kozielska et al. (2010) consider systems in which

the ancestral sex chromosomes experience meiotic drive (e.g., driving X or Y chromosomes are inherited disproportionately often), which leads to biased sex ratios. They find that new, unlinked sex-determining loci (masculinizing or feminizing mutations) can then spread, restoring an even sex ratio.

For example, new sex-determining systems (particularly transitions between male and female heterogamety) can be favoured in order to restore equal sex ratios in populations that have a sex ratio bias (Bull 1983, Kozielska et al. 2010, Úbeda et al. 2015).

Sex ratios can also become biased due to meiotic drive; in a classic paper, Hamilton (1967) showed that X- or Y-linked alleles that experience meiotic drive will bias sex ratios. He assumed that driving alleles are under directional selection and spread to fixation but such alleles can also be maintained at intermediate frequencies by selection (Feldman and Otto 1989, Holman et al. 2015). When sex ratios are biased, other loci are expected to evolve to restore equal sex ratios. Indeed, alleles that negate the effect of sex-linked meiotic drivers and restore equal sex ratios have been identified (Stalker 1961, Smith 1975). A similar process occurs with cytoplasmic male sterility alleles (that cause biased sex ratios) and nuclear ‘restorer’ genotypes (Frank 1989).

When sex ratio bias occurs due to haploid selection, a natural class of sex ratio ‘restorers’ exist because haploid selection often occurs in a context that is determined by the diploid parents. For example, the intensity of pollen competition can be manipulated by altering style length (Travers and Shea 2001, Lankinen and Skogsmyr 2001, Ruane 2009), delaying stigma receptivity (Galen et al. 1986, Lankinen and Madjidian 2011) and/or delaying pollen tube growth in the pistil (Herrero 2003). Where the X and Y have fixed fitness differences, Hough et al. (2013) demonstrated that mothers should generally evolve to balance sex ratios by reducing the intensity of haploid competition. However, reducing competition among haploids also reduces the potential for harmful deleterious mutations to be purged. When deleterious mutations are included, the optimal intensity of haploid selection can reflect

a balance between maximizing offspring fitness and equalizing sex ratios.

As part of a collaborative project (Otto et al. 2015), I considered the evolution of the haploid ‘selective arena’ in cases where the X chromosome harbours a polymorphism that affects haploid fitness. Mothers again primarily evolve to restore equal sex ratios. However, modifying haploid selection also affects the X-linked genotypes that are inherited by offspring. Specifically, increasing the intensity of haploid selection increases the proportion of daughters (all progeny of X-bearing sperm/pollen are female) that inherit the allele with high haploid fitness. If this allele has high fitness in daughters, mothers can be selected to increase the intensity of haploid selection; otherwise, decreased selection among haploids is favoured. Thus, because altering haploid selection intensity affects the alleles that are inherited by daughters, mothers can favour slightly biased sex ratios. In addition, I found that stronger sex ratio biases can be favoured by paternal manipulations of the haploid ‘selective arena’ because fathers are strongly selected to maximize their own siring success (above selection to equalize the sex ratio).

It is not clear how the spread of new sex determination systems would be influenced by the combination of sex ratio biases and favourable associations between haploid selected loci and sex-determining regions.

We add haploid selection (and justification, see below)
Mention Hough study for sex ratio adjustment here or in discussion? Also discuss the fact that, in terms of recombination suppression, haploid selection among male gametes generates selection pressure similar to that of male specific selection.

What will be the result: where there is sex biases and sex-specific selection

FROM PREVIOUS PAPER:

To the extent that pollen and sperm success reflects differences in their haploid genotypes, selection among these gametes/gametophytes is qualitatively distinct from selection among diploid males. That is, diploids cannot be assigned fitness values that also account for the fitness of their haploid

gametes Immler et al. (2012). In plants, selection among haploid male gametophytes is thought to be pervasive Skogsmyr and Lankinen (2002), Moore and Pannell (2011), Marshall and Evans (2016); in *Arabidopsis*, 60-70% of all genes are expressed during the haploid phase Borg et al. (2009), and pollen expressed genes exhibit stronger signatures of purifying selection and positive selection Arunkumar et al. (2013), Gossmann et al. (2014). For agricultural breeding, pollen has been exposed to a variety of selection pressures *in vivo* and *in vitro*, including temperature Hedhly et al. (2004), Clarke et al. (2004), herbicides Frascaroli and Songstad (2001), metals Searcy and Mulcahy (1985), water stress Ravikumar et al. (2003), and pathogens Ravikumar et al. (2012), resulting in an increased frequency of resistant genotypes among the diploid sporophytic offspring. In animals, expression during the haploid sperm stage is traditionally thought to be suppressed Hecht (1998), although recent evidence suggests that the extent and selective importance of post-meiotic gene expression may be underestimated Zheng et al. (2001), Joseph and Kirkpatrick (2004), Vibranovski et al. (2010), Immler et al. (2014).

Here, suggest that the canonical view (no haploid expression in animals, genome highly chromatinized in sperm and not expressed) might be based on model organisms, such as mice, where sperm is sufficiently short-lived that transcripts provisioned during spermatogenesis may be sufficient without further haploid transcription (although note that the Vibranovski lab results are in mice showing some transcription does occur). In broadcast spawning animal species (e.g., corals, many fish) and species where sperm typically requires greater longevity, expression of the haploid genotype may be required (Immler paper indicates this, but not that strongly - as I remember). We can use this suggestion in discussion to speculate in what species the processes we study might be looked for (i.e., animals with multiple matings, broadcast spawning and/or long-lived sperm and outcrossing/non-pollen-limited plants).

FROM PREVIOUS PAPER:

The maintenance of polymorphism at loci that experience sex specific selection in both haploid and diploid phases was considered by Immler et al. Immler et al. (2012), demonstrating that polymorphisms can be maintained by sexually antagonistic selection or overdominance as well as by conflicting selection pressures in haploids and diploids (haploid-diploid conflict or ploidy antagonistic selection) or a combination of these selective regimes.

NOTE RE: DRIVE. I expect drive (that occurs specifically in one sex, e.g., during spermatogenesis) to behave almost exactly like haploid selection. That is, I think that a XY-linked driver that is maintained by selection (e.g., because it causes sterility when homozygous, which is common in known drive systems) will only favour invasion of a more tightly linked neo-Y (worsening sex ratio biases) and could favour invasion of a neo-W. This may run counter to generic expectations from new sex chromosome systems evolving to balance the sex ratio. So, do you think it would significantly enhance the paper to model drive explicitly or just discuss it as being similar???

Discussion

2 FROM THESIS: Generally, any sex-linked gene that harbours genetic variation in haploid fitness should cause sex ratios to become biased. Sex ratio bias
4 caused by pollen competition has previously been discussed in the context of Y-linked deleterious mutations, which are thought to build up after recombination suppression evolves (Lloyd 1974, Stehlik and Barrett 2005). Sex
6 ratios can also become biased due to meiotic drive; in a classic paper, Hamilton (1967) showed that X- or Y-linked alleles that experience meiotic drive
8 will bias sex ratios. He assumed that driving alleles are under directional selection and spread to fixation but such alleles can also be maintained at
10 intermediate frequencies by selection (Feldman and Otto 1989, Holman et al. 2015). When sex ratios are biased, other loci are expected to evolve to restore
12 equal sex ratios. Indeed, alleles that negate the effect of sex-linked meiotic

14 drivers and restore equal sex ratios have been identified (Stalker 1961, Smith
1975). A similar process occurs with cytoplasmic male sterility alleles (that
16 cause biased sex ratios) and nuclear ‘restorer’ genotypes (Frank 1989).

When sex ratio bias occurs due to haploid selection, a natural class of sex
18 ratio ‘restorers’ exist because haploid selection often occurs in a context that
is determined by the diploid parents. For example, the intensity of pollen
20 competition can be manipulated by altering style length (Travers and Shea
2001, Lankinen and Skogsmyr 2001, Ruane 2009), delaying stigma receptivity
22 (Galen et al. 1986, Lankinen and Madjidian 2011) and/or delaying pollen
tube growth in the pistil (Herrero 2003). Where the X and Y have fixed
24 fitness differences, Hough et al. (2013) demonstrated that mothers should
generally evolve to balance sex ratios by reducing the intensity of haploid
26 competition. However, reducing competition among haploids also reduces the
potential for harmful deleterious mutations to be purged. When deleterious
28 mutations are included, the optimal intensity of haploid selection can reflect
a balance between maximizing offspring fitness and equalizing sex ratios.

30 As part of a collaborative project (Otto et al. 2015), I considered the
evolution of the haploid ‘selective arena’ in cases where the X chromosome
32 harbours a polymorphism that affects haploid fitness. Mothers again primar-
ily evolve to restore equal sex ratios. However, modifying haploid selection
34 also affects the X-linked genotypes that are inherited by offspring. Specif-
ically, increasing the intensity of haploid selection increases the proportion
36 of daughters (all progeny of X-bearing sperm/pollen are female) that inherit
the allele with high haploid fitness. If this allele has high fitness in daugh-
38 ters, mothers can be selected to increase the intensity of haploid selection;
otherwise, decreased selection among haploids is favoured. Thus, because
40 altering haploid selection intensity affects the alleles that are inherited by
daughters, mothers can favour slightly biased sex ratios. In addition, I found
42 that stronger sex ratio biases can be favoured by paternal manipulations of
the haploid ‘selective arena’ because fathers are strongly selected to maximize

44 their own siring success (above selection to equalize the sex ratio).

Several aspects of the relationship between haploid selection (e.g., pollen
46 or sperm competition) and sex ratios remain to be explored. For example,
new sex-determining systems (particularly transitions between male and fe-
48 male heterogamety) can be favoured in order to restore equal sex ratios in
populations that have a sex ratio bias (Bull 1983, Kozielska et al. 2010, Úbeda
50 et al. 2015). Based on the results of Chapter ??, we would expect that sex
ratio biases would occur via associations between sex-determining loci and
52 loci that experience haploid selection. However, these associations should
also select against transitions between sex-determining systems, as has been
54 found with sexually antagonistic selection (van Doorn and Kirkpatrick 2007;
2010). It is not clear how the spread of new sex determination systems would
56 be influenced by the combination of sex ratio biases and favourable associ-
ations between haploid selected loci and sex-determining regions. Finally,
58 Hamilton (1967) pointed out that biased sex ratios can affect population size
because the number of offspring in each generation is typically determined by
60 the number of females. Population density can, in turn, affect the intensity
of pollen/sperm competition in future generations because fewer males are
62 available to donate pollen/sperm in a particular area. Thus, a feedback could
occur between population densities and haploid selection, which has not yet
64 been investigated.

References

- 66 Arunkumar, R., E. B. Josephs, R. J. Williamson, and S. I. Wright. 2013.
Pollen-specific, but not sperm-specific, genes show stronger purifying se-
68 lection and higher rates of positive selection than sporophytic genes in
Capsella grandiflora. *Molecular biology and evolution* 30:2475–2486.
- 70 Bachtrog, D., J. E. Mank, C. L. Peichel, M. Kirkpatrick, S. P. Otto, T.-L.
Ashman, M. W. Hahn, J. Kitano, I. Mayrose, R. Ming, N. Perrin, L. Ross,

- 72 N. Valenzuela, J. C. Vamosi, and Tree of Sex Consortium. 2014. Sex
determination: why so many ways of doing it? *PLoS Biol* 12:e1001899.
- 74 Beukeboom, L. W., and N. Perrin. 2014. The evolution of sex determination.
Oxford University Press, Oxford, UK.
- 76 Borg, M., L. Brownfield, and D. Twell. 2009. Male gametophyte development:
a molecular perspective. *Journal of Experimental Botany* 60:1465–1478.
- 78 Bull, J. J. 1983. Evolution of sex determining mechanisms. The Benjamin
Cummings Publishing Company.
- 80 Charlesworth, D., and J. E. Mank. 2010. The birds and the bees and the
flowers and the trees: lessons from genetic mapping of sex determination
82 in plants and animals. *Genetics* 186:9–31.
- Clarke, H. J., T. N. Khan, and K. H. M. Siddique. 2004. Pollen selection
84 for chilling tolerance at hybridisation leads to improved chickpea cultivars.
Euphytica 139:65–74.
- 86 Feldman, M. W., and S. P. Otto. 1989. More on recombination and selec-
tion in the modifier theory of sex-ratio distortion. *Theoretical Population*
88 *Biology* 35:207–225.
- Frank, S. A. 1989. The Evolutionary Dynamics of Cytoplasmic Male Sterility.
90 *American Naturalist* 133:345–376.
- Frascaroli, E., and D. D. Songstad. 2001. Pollen genotype selection for a
92 simply inherited qualitative factor determining resistance to chlorsulfuron
in maize. *Theoretical and Applied Genetics* 102:342–346.
- 94 Galen, C., J. A. Shykoff, and R. C. Plowright. 1986. Consequences of stigma
receptivity schedules for sexual selection in flowering plants. *American*
96 *Naturalist* pages 462–476.

- Gossmann, T. I., M. W. Schmid, U. Grossniklaus, and K. J. Schmid. 2014.
 98 Selection-driven evolution of sex-biased genes Is consistent with sexual
 selection in *Arabidopsis thaliana*. *Molecular biology and evolution* 31:574–
 100 583.
- Hamilton, W. D. 1967. Extraordinary sex ratios. *Science* 156:477–488.
- 102 Hecht, N. B. 1998. Molecular mechanisms of male germ cell differentiation.
Bioessays 20:555–561.
- 104 Hedhly, A., J. I. Hormaza, and M. Herrero. 2004. Effect of temperature
 on pollen tube kinetics and dynamics in sweet cherry, *Prunus avium*
 106 (Rosaceae). *American journal of botany* 91:558–564.
- Herrero, M. 2003. Male and female synchrony and the regulation of mating
 108 in flowering plants. *Philosophical Transactions of the Royal Society B:
 Biological Sciences* 358:1019–1024.
- 110 Hillis, D. M., and D. M. Green. 1990. Evolutionary changes of heterogametic
 sex in the phylogenetic history of amphibians. *Journal of Evolutionary*
 112 *Biology* 3:49–64.
- Holman, L., T. A. R. Price, N. Wedell, and H. Kokko. 2015. Coevolutionary
 114 dynamics of polyandry and sex-linked meiotic drive. *Evolution* 69:709–720.
- Hough, J., S. Immler, S. Barrett, and S. P. Otto. 2013. Evolutionarily stable
 116 sex ratios and mutation load. *Evolution* 7:1915–1925.
- Immler, S., G. Arnqvist, and S. P. Otto. 2012. Ploidally antagonistic selection
 118 maintains stable genetic polymorphism. *Evolution* 66:55–65.
- Immler, S., C. Hotzy, G. Alavioon, E. Petersson, and G. Arnqvist. 2014.
 120 Sperm variation within a single ejaculate affects offspring development in
 Atlantic salmon. *Biology letters* 10:20131040.

- 122 Joseph, S., and M. Kirkpatrick. 2004. Haploid selection in animals. Trends
in Ecology & Evolution 19:592–597.
- 124 Kozielska, M., F. J. Weissing, L. W. Beukeboom, and I. Pen. 2010. Segrega-
tion distortion and the evolution of sex-determining mechanisms. Heredity
126 104:100–112.
- Lankinen, A., and J. A. Madjidian. 2011. Enhancing pollen competition
128 by delaying stigma receptivity: Pollen deposition schedules affect siring
ability, paternal diversity, and seed production in *Collinsia heterophylla*
130 (Plantaginaceae). American journal of botany 98:1191–1200.
- Lankinen, A., and I. Skogsmyr. 2001. Evolution of pistil length as a choice
132 mechanism for pollen quality. Oikos 92:81–90.
- Li, J., R. B. Phillips, A. S. Harwood, B. F. Koop, and W. S. Davidson. 2011.
134 Identification of the Sex Chromosomes of Brown Trout (*Salmo trutta*)
and Their Comparison with the Corresponding Chromosomes in Atlantic
136 Salmon (*Salmo salar*) and Rainbow Trout (*Oncorhynchus mykiss*). Cyto-
genetic and Genome Research 133:25–33.
- 138 Lloyd, D. G. 1974. Female-predominant sex ratios in angiosperms, vol. 32.
Heredity.
- 140 Mank, J. E., D. E. L. Promislow, and J. C. Avise. 2006. Evolution of alter-
native sexdetermining mechanisms in teleost fishes. Biological Journal of
142 the Linnean Society 87:83–93.
- Marshall, D. L., and A. S. Evans. 2016. Can selection on a male mating char-
144 acter result in evolutionary change? A selection experiment on California
wild radish, *Raphanus sativus*. American journal of botany 103:553–567.
- 146 Ming, R., A. Bendahmane, and S. S. Renner. 2011. Sex chromosomes in land
plants. dx.doi.org 62:485–514.

- 148 Moore, J. C., and J. R. Pannell. 2011. Sexual selection in plants. *Current*
Biology 21:R176–R182.
- 150 Myosho, T., H. Otake, H. Masuyama, M. Matsuda, Y. Kuroki, A. Fujiyama,
K. Naruse, S. Hamaguchi, and M. Sakaizumi. 2012. Tracing the Emergence
152 of a Novel Sex-Determining Gene in Medaka, *Oryzias luzonensis*. *Genetics*
191:163–170.
- 154 Ogata, M., Y. Hasegawa, H. Ohtani, M. Mineyama, and I. Miura. 2007. The
ZZ/ZW sex-determining mechanism originated twice and independently
156 during evolution of the frog, *Rana rugosa*. *Heredity* 100:92–99.
- Otto, S. P., M. F. Scott, and S. Immler. 2015. Evolution of haploid selection
158 in predominantly diploid organisms. *Proceedings of the National ...*
- Ravikumar, R. L., G. N. Chaitra, A. M. Choukimath, and C. D. Soregaon.
160 2012. Gametophytic selection for wilt resistance and its impact on the
segregation of wilt resistance alleles in chickpea (*Cicer arietinum* L.). *Eu-*
162 *phytica* 189:173–181.
- Ravikumar, R. L., B. S. Patil, and P. M. Salimath. 2003. Drought tolerance in
164 sorghum by pollen selection using osmotic stress. *Euphytica* 133:371–376.
- Ruane, L. G. 2009. Post-pollination processes and non-random mating among
166 compatible mates. *Evolutionary Ecology Research* 11:1031–1051.
- Searcy, K. B., and D. L. Mulcahy. 1985. Pollen selection and the game-
168 tophytic expression of metal tolerance in *Silene dioica* (Caryophyllaceae)
and *Mimulus guttatus* (Scrophulariaceae). *American journal of botany*
170 72:1700–1706.
- Ser, J. R., R. B. Roberts, and T. D. Kocher. 2010. Multiple interacting loci
172 control sex determination in lake Malawi cichlid fish. *Evolution* 64:486–
501.

- 174 Skogsmyr, I., and A. Lankinen. 2002. Sexual selection: an evolutionary force
in plants? *Biological Reviews* 77:537–562.
- 176 Slancarova, V., J. Zdanska, B. Janousek, M. Talianova, C. Zschach, J. Zlu-
vova, J. Siroky, V. Kovacova, H. Blavet, J. Danihelka, B. Oxelman, A. Wid-
178 mer, and B. Vyskot. 2013. Evolution of sex determination systems with
heterogametic males and females in *Silene*. *Evolution* 67:3669–3677.
- 180 Smith, D. A. S. 1975. All-female broods in the polymorphic butterfly *Danaus*
chrysippus L. and their ecological significance. *Heredity* 34:363–371.
- 182 Stalker, H. D. 1961. The Genetic Systems Modifying Meiotic Drive in
Drosophila Paramelanica. *Genetics* .
- 184 Stehlik, I., and S. Barrett. 2005. Mechanisms governing sex-ratio variation
in dioecious *Rumex nivialis*. *Evolution* 59:814–825.
- 186 Travers, S. E., and K. Shea. 2001. Selection on pollen competitive ability in
relation to stochastic factors influencing pollen deposition. *Evolutionary*
188 *Ecology Research* 3:729–745.
- Úbeda, F., M. M. Patten, and G. Wild. 2015. On the origin of sex chromo-
190 somes from meiotic drive. *Proceedings of the Royal Society B: Biological*
Sciences 282:20141932.
- 192 van Doorn, G. S., and M. Kirkpatrick. 2007. Turnover of sex chromosomes
induced by sexual conflict. *Nature* 449:909–912.
- 194 ———. 2010. Transitions Between Male and Female Heterogamety Caused
by Sex-Antagonistic Selection. *Genetics* 186:629–645.
- 196 Vibranovski, M. D., D. S. Chalopin, H. F. Lopes, M. Long, and T. L. Karr.
2010. Direct evidence for postmeiotic transcription during *Drosophila*
198 *melanogaster* spermatogenesis. *Genetics* 186:431–433.

- 200 Vicoso, B., and D. Bachtrog. 2015. Numerous transitions of sex chromosomes
in Diptera. *PLoS Biol* 13:e1002078.
- 202 Yano, A., B. Nicol, E. Jouanno, E. Quillet, A. Fostier, R. Guyomard,
and Y. Guiguen. 2012. The sexually dimorphic on the Y-chromosome
gene (sdY) is a conserved male-specific Y-chromosome sequence in many
204 salmonids. *Evolutionary Applications* 6:486–496.
- Zheng, Y., X. Deng, and P. A. Martin-DeLeon. 2001. Lack of sharing of
206 Spam1 (Ph-20) among mouse spermatids and transmission ratio distortion.
Biology of Reproduction 64:1730–1738.