

Quadratic Residue Modulo $N = pq$, p and q are distinct odd primes

$$\mathbb{Z}_{15}^* = \{1, 2, 4, 7, 8, 11, 13, 14\} \quad \phi(15) = \phi(3)\phi(5) \\ = 2 \cdot 4 \\ = 8$$
$$\mathbb{QR}_{15} = \{1, 4\}$$

$$\mathbb{Z}_{15}^* \simeq \mathbb{Z}_3^* \times \mathbb{Z}_5^*$$
$$x \mapsto (x \bmod 3, x \bmod 5)$$
$$1 \mapsto (1, 1)$$
$$4 \mapsto (1, 4)$$

$$\mathbb{Z}_3^* = \{1, 2\}$$
$$\mathbb{QR}_3 = \{1\}$$
$$\mathbb{Z}_5^* = \{1, 2, 3, 4\}$$
$$\mathbb{QR}_5 = \{1, 4\}$$

Theorem: $\mathbb{QR}_N \simeq \mathbb{QR}_p \times \mathbb{QR}_q$

$$\text{let } x_p = x \bmod p$$

$$x_q = x \bmod q$$

if $x_p \in \mathbb{QR}_p$ and $x_q \in \mathbb{QR}_q$, then $x = (x_p, x_q) \in \mathbb{QR}_N$

if $x \in \mathbb{QR}_N$ then $x_p \in \mathbb{QR}_p$ and $x_q \in \mathbb{QR}_q$

proof: If $x_p \in \mathbb{QR}_p$ and $x_q \in \mathbb{QR}_q$, there exists a and b s.t.
 $a^2 = x_p \bmod p$ and $b^2 = x_q \bmod q$.

Hence $(x_p, x_q) = (a^2, b^2)$ is a \mathbb{QR}_N .

If $(x_p, x_q) \in \mathbb{QR}_N$, there exist a and b s.t.

$$(a, b) \cdot (a, b) = (a^2, b^2) = (x_p, x_q), \text{ hence}$$

$$x_p \in \mathbb{QR}_p \text{ and } x_q \in \mathbb{QR}_q$$

Theorem: If $x \in \mathbb{QR}_N$, then x has **four** square roots.

proof: $x = (x_p, x_q)$

Any square root of x is of form (a, b)

such that a is square root of x_p

b is square root of x_q

There are two square roots of x_p .

two square roots of x_q .

In total, four square roots of (x_p, x_q)

Example: What is square root of 4?

$4 \rightarrow (1, 4)$

↓
square roots of 1 in \mathbb{Z}_3^\times is 1, 2

↓
square roots of 4 in \mathbb{Z}_5^\times is 2, 3

square root of $(1, 4)$ is

7 ← $(1, 2)$

13 ← $(1, 3)$

2 ← $(2, 2)$

8 ← $(2, 3)$


7, 13, 2, 8 are square roots of 4

none of these square roots are \mathbb{QR}_N .

$p=3, q=5$ ($p \equiv 3 \pmod{4}$ but $q \not\equiv 3 \pmod{4}$)

In Homework 8, Question 5, when $p \equiv q \equiv 3 \pmod{4}$, exactly one of the square root is a \mathbb{QR} .

Corollary: $|\mathbb{Q}R_N| = \frac{1}{N} |\mathbb{Z}_N^*|$



Algorithm to check if an element is QRN

Input: x, N

Output: QR is x is quadratic residue, QNR o/w.

Algorithm: Compute $J_p(x_p) = x_p^{\frac{p-1}{2}}$

$$J_q(x_q) = x_q^{\frac{q-1}{2}}$$

If $J_p(x_p) = J_q(x_q) = 1$ then output QR
o/w QNR.

Running Time: Polynomial

Algorithm to find square roots of QR_N

Input: N, x where $x \in \text{QR}_N$

Output: Square roots of $x \bmod N$

Algorithm: Compute p and q such that $N = pq$

Compute square roots of x_p : a_1, a_2

Compute square roots of x_q : b_1, b_2

Output $(a_1, b_1), (a_1, b_2), (a_2, b_1), (a_2, b_2)$
(after using CTR to convert (a, b) to $c \in \mathbb{Z}_N^*$)

$p \equiv 3 \bmod 4$, a_1, a_2 are $\pm x_p^{\frac{p+1}{4}}$

$q \equiv 3 \bmod 4$, b_1, b_2 are $\pm x_q^{\frac{q+1}{4}}$

Running Time: Polynomial if $p \equiv q \equiv 3 \bmod 4$.

Theorem

1. If factoring is easy, then it is easy to find square root modulo N .
 2. If square root modulo N is easy, then it is easy to factor N .
-

2. Given N and x ,

Suppose you can find all square roots of $x \bmod N$
 x_1, x_2, x_3, x_4 .

How to use these square roots to find p, q .

Example: $N=15$

square roots of 4 is $7, -7, 2, -2$