

Joint inference of adaptive and demographic history from temporal population genomic data

Vitor A. C. Pavinato^{1,2,3}, Stéphane De Mita^{4,5}, Jean-Michel Marin^{2,2}, Miguel de Navascués^{1,5*1,6*}

¹CBGP, INRAE, CIRAD, IRD, ~~Montpellier SupAgro~~ Institut Agro, Université de Montpellier, Montpellier, France

²UMR Institut Montpelliérain Alexander Grothendieck, Université de Montpellier, France

³Entomology Dept., CFAES, The Ohio State University, Wooster, USA

⁴UMR Interactions Arbres-Microorganismes, INRAE, France

⁵INRAE, CIRAD, Institut Agro, IRD, PHIM, Université Montpellier, Montpellier, France

⁶Human Evolution, Department of Organismal Biology, Uppsala University, Uppsala, Sweden

^{*}corresponding author: miguel.navascues@inrae.fr

September 2, 2022

Abstract

Disentangling the effects of selection and drift is a long-standing problem in population genetics. ~~Recently, simulations shows~~ Simulations show that pervasive selection may bias the inference of demography. Ideally, models for the inference of demography and selection should account for the interaction between these two forces. With simulation-based likelihood-free methods such as Approximate Bayesian Computation (ABC), demography and selection parameters can be jointly estimated (~~ABC~~). We propose ~~a to use the~~ ABC-Random Forests framework to jointly infer demographic and selection parameters from temporal population genomic data (e.g., experimental evolution, monitored populations, ancient DNA). Our framework allowed the separation of demography (census size, N) from the genetic drift (effective population size, N_e), and the estimation of genome-wide parameters of selection. Selection parameters informed us about the adaptive potential of a population (the ~~scale scaled~~ mutation rate of beneficial mutations, θ_b), the realized ~~potential, as adaptation~~ (the number of ~~strong beneficial under selection, and fitness diversity as the population genetic load~~ mutation under strong selection), and population fitness (genetic load). We applied this approach to a dataset of feral populations of honey bees (*Apis mellifera*) collected in California, and we estimated parameters consistent with the biology and the recent history of this species.

Keywords— Temporal data, Population genomics, Machine learning, Adaptation

Introduction

One aim of population genomics is to understand how demography and natural selection shape the genetic diversity of populations. A classical approach ~~is to assume~~ assumes that demography (migration, population subdivision, population size changes) leaves a genome-wide signal; ~~whereas~~. In contrast, selection leaves a ~~localizes~~ localized signal close to where the causal mutation is located. Many methods follow this approach to infer demography or selection (reviewed by Beichman et al., 2018; Casillas and Barbadilla, 2017). ~~Methods for demographic inference assume~~ Demographic inference assumes that most of the genome evolves without the influence of selection and that any deviation from the mutation-drift equilibrium observed in the data was caused by demographic events (Beichman et al., 2018). ~~For selection, the majority~~ Many of the methods search for locus-specific signals ~~left by the beneficial mutation of selection~~ left on nearby neutral mutations (Tajima, 1989; Fay and Wu, 2000; Kim and Nielsen, 2004) (low genetic diversity and high differentiation) to localize the ~~beneficial~~ region affected by selection mutation, assuming ~~constant population size~~ (Nielsen, 2005; Pool et al., 2010) a specific demography (constant population size in early methods; Nielsen, 2005; Pool et al., 2010).

Conducting demographic and selection inference separately may have some shortcomings. First, there is the assumption that the signal left by demography is little affected by selection because the selection is rare. However, linked selection can affect neutral and weakly selected sites that are far from the mutation targeted by selection (Sella et al., 2009; Neher, 2013) and selection can be pervasive (Sella et al., 2009; Lange and Pool, 2018). In addition, some methods for selection scans are not robust to misspecifications of demographic history. Consequently, an unspecified bottleneck or population increase, for example, can inflate the ~~type I error~~ false positive rate of genome scans (Jensen et al., 2005, 2007; Schrider et al., 2016). These findings highlight the necessity of inferential methods that jointly accounts for the multiple evolutionary forces that act on populations (Lin et al., 2011; Li et al., 2012; Bank et al., 2014).

It is ~~difficult to obtain~~ often difficult to calculate the likelihood of models including demography and selection (but see Vitalis et al., 2014). Methods that rely on simulations provide ~~alternatives to the use of~~ easier alternatives to using likelihood functions (Csilléry et al., 2010; Schrider and Kern, 2018). One of the first works that proposed such a strategy addressed the inference of local adaptation (Bazin et al., 2010). With coalescent simulations of an island model, Bazin et al. (2010) estimated demographic parameters and inferred the number of loci under selection. In their simulations, the selection was modeled as ~~differential~~ locus-specific migration rates in which a selected locus had lower migration rates than ~~neutrally-evolved~~ neutral loci. However, locus-specific migration rates or effective population size (as in Roux et al., 2016; Fraïsse et al., 2021) represent crude approximations of the selection process. Forward-in-time simulation allows more realistic models of selection. These were used to make inferences on N_e in the presence of selection by Sheehan and Song (2016) (~~for selective sweep~~ selective sweeps and balancing selection) and Johri et al. (2020) (background selection). However, these works rely on ~~simulations of few independent loci not more than 50Kbp which prevents the modelling~~ independent simulation loci or genomic regions which prevents the modeling of genome-wide effects of selection as the reduction of effective population size due to the variance of reproductive success of individuals (~~e.g. Santiago and Caballero, 1995~~) (Santiago and Caballero, 1995) or the combined effects of mutations on individual fitness. Nevertheless, this strategy brought new insights into the dynamics of selection. ~~For instance~~, Laval et al. (2019) estimated the number of past selective sweeps ~~that occurred~~ in the human genome in the past ~~10~~ 100,000 years, their intensity, and their age. Besides some limitations, these works exemplify the power of likelihood-free methods ~~for the inference of~~ to infer the complex interaction between demography and selection.

Most population genetic studies use samples collected at ~~one time~~ a one-time point to infer the neutral processes (mutation, recombination, random genetic drift) and selection throughout the history of populations. Temporal data allows a better understanding of recent evolutionary

processes (~~Feder et al., 2021; Dehasque et al., 2020~~) (e.g. Feder et al., 2021; Dehasque et al., 2020) because they contain information about the allele frequency changes through time. By tracking the allele frequency changes over time, it is possible to estimate the relative role of selection and drift. Consequently, temporal data has the potential to give us a better understanding of the interaction between drift and selection (see for example, Buffalo and Coop, 2019, 2020).

Here, we propose ~~the use of~~ using ABC to jointly estimate demography and positive selection from temporal genomic data. In our framework, we use ~~agent-based~~ individual-based, forward-in-time simulations ~~which allowed the modelling,~~ which allow the modeling of the genome-wide ~~and linked selection,~~ linked selection and additive effects of beneficial mutations. Until recently, ~~the use of~~ such computationally demanding simulations in ABC inference ~~was unrealistic since a great number of~~ were unrealistic since many simulations are required to achieve accuracy in ABC (Frazier et al., 2018). However, with the introduction of Random Forests (ABC-RF), it ~~was~~ is now possible to reduce the computational burden as fewer simulations are required to achieve reliable estimates (Pudlo et al., 2016; Raynal et al., 2019). While many methods focus on the detection of targets of selection, our work addresses the inference of parameters that characterizes the genome-wide signal of demography and selection. Our genome-wide estimates ~~showed to be~~ were reasonably accurate for a wide ~~rate of adaptation~~ range of adaptation rates and strength of selection. We were able to separate the estimates of N_e ~~;~~ which reflect the short-term drift (a measure of genetic drift) from the population census size N . We also estimated the ~~rate of~~ influx of new beneficial mutations as measured by ~~genome-wide~~ the population scaled mutation rate of beneficial mutations. The separation between demography and drift ~~;~~ and the inference of genome-wide selection ~~were only possible with the use of~~ was only possible using latent variables. Latent variables emerged as properties of each simulation, and consequently, they better captured the emerging interaction between demography and selection than model parameters. We first evaluated the performance of an ABC-RF approach with forward-in-time simulations, ~~and finally.~~ Finally, we applied this framework to the analysis of a real time-series population genomics dataset of the feral population of honey bees (*Apis mellifera*) (Cridland et al., 2018). Our results were consistent with the ~~biology of the species~~ species' biology and with events that occurred recently in the history of the analyzed populations, taking into account the limitations of the current implementation of our approach.

Methods

Inference model

We assumed a closed population (no migration) of N diploid individuals that evolved under a Wright-Fisher model with selection. ~~Selection~~ The population census size N was constant, and selection only acted on *de novo* beneficial mutations ~~and~~ that were allowed to arise in the population since the first generation (generation one corresponds to the first burn-in generation). Every beneficial mutation had a selection coefficient of s higher than zero, and all were co-dominant. The values of the selection coefficients s were drawn from a gamma distribution with mean γ ~~;~~ and scale parameter 1. Beneficial mutations entered the population with a rate of μ_b per generation ~~that is~~ independent of the mutation selective strength. ~~Consequently~~ Consequently, we defined the scaled mutation rate of the beneficial mutations per generation θ_b as the product the population size N , the mutation rate of beneficial mutation μ_b and the genome size G , $\theta_b = 4N\mu_b G$. This rate determines the amount of new beneficial mutations that ~~arises~~ arise in the population every generation. It can also be viewed as the waiting time for the appearance of a new beneficial mutation in the population. Populations with high θ_b receive new beneficial mutations every generation (Karasov et al., 2010), but a population with low θ_b needs to wait more time for a new beneficial mutation to arise.

We ~~needed to divide the model in~~ divided the model into two periods: 1) the burn-in period,

that ~~which~~ is necessary to remove ~~,~~ from the simulations ~~,~~ any footprint of the initial ~~parameters~~ ~~sets~~ ~~simulation state~~; the duration of this period was defined as the time necessary to ~~contain all~~ ~~reach a point where the~~ most recent common ancestors (MRCA) for all genomic regions ~~in the~~ ~~simulation are more recent than the start of the simulation (i.e. the burn-in was run until this~~ ~~condition was fulfilled)~~; and 2) the inference period, where we defined the longitudinal samples of individuals; ~~the sampled genotypes were used to make inference of demography and selection.~~ ~~This.~~ These two periods were defined by their time ~~span and by~~ ~~span and~~ the population census size ~~with,~~ being N_0 and N as the ~~sample population~~ size of the burn-in and the inference period, ~~respectively. Population size is constant within each simulated period and changes between~~ ~~periods.~~

The ~~first~~ ~~First~~ sample of individuals was taken at t_1 , ~~that was the first generation after~~ the ~~immediate next generation after~~ burn-in ~~period ended, and ended;~~ the second was taken at t_2 , after τ generations from t_1 . Individuals were sampled following the sample plan II of Nei and Tajima (1981), where individuals were taken before ~~their~~ reproduction and permanently removed from the population. In this way, ~~their~~ genotypes did not contribute to the next generation ~~genotypes.~~

Each ~~individuals~~ ~~individual's~~ genome of size G (in base pairs) consisted of a single linkage group with ~~a~~ per base recombination rate per generation of e_0 . ~~In this genome, we modeled the~~ ~~act of selection~~ ~~r .~~ We modeled the selection effect in this genome by dividing it ~~in into~~ “neutral” and “non-neutral” regions. Non-neutral regions held both neutral and beneficial mutations. This division can be interpreted as a genomic architecture in which ~~there are genic regions~~ ~~that genic regions~~ have a combination of neutral (synonymous intron mutations) and selected (non-synonymous mutation) sites ~~,~~ ~~and with~~ intergenic regions (neutral mutations) ~~but also~~ ~~other characteristics of the genome that would make the action of selection heterogeneous in~~ ~~between.~~ However, this architecture allowed simulating the heterogeneous selection action along the genome.

We chose this simplification because it is ~~general and it was a~~ ~~a~~ ~~general and~~ straightforward way to define independent priors for the relative number of non-neutral to neutral regions ~~,~~ and for the number of beneficial mutations in non-neutral regions. The probability of beneficial mutation to arise in the simulation ~~,~~ ~~or their~~ ~~(i.e. the~~ mutation rate per generation ~~μ_b ,~~ ~~μ_b)~~ was determined by the product of the proportion of non-neutral regions P_R , the proportion of beneficial mutation in a non-neutral region P_B and the mutation rate per generation μ . Figure 1 shows a schematic representation of the model template (and see Table S1 for a summary of the notation).

Calculation of summary statistics and latent variables

The above model was used to simulate the dynamic of drift and selection in a closed population. In the two sample periods, individuals ~~from the whole population~~ were sampled and used for the calculation of the summary statistics for the ABC-RF framework. For each simulation, we calculated summary statistics that: 1) compared the two samples (*e.g.* ~~genetic differentiation~~ ~~genetic differentiation~~, F_{ST}), and 2) quantified the diversity within-sample (*e.g.* ~~expected heterozygosity~~ ~~expected heterozygosity~~, H_E). For the ~~later, calculations~~ ~~latter, statistics~~ were obtained for each ~~sample and the pooled sample (when the genetic data of populations were combined in~~ ~~one population) and all pooled samples~~. Some summary statistics were calculated genome-wide; ~~for.~~ For example, global F_{ST} , global H_E and the total number of polymorphic sites S ; others were calculated SNP-by-SNP as the H_E ; or they were calculated in windows as S , the nucleotide diversity π , and Tajima's D . For every simulation, we measured the mean, variance, kurtosis, skewness, and 5% and 95% quantiles among all locus-specific or window summary statistics. These statistics ~~informed~~ ~~inform~~ about the heterogeneity of genome-wide distribution of locus-specific or window summary statistics. We set three window sizes for the window summary




Figure 1: A schematic representation of the model used to simulate temporal population genomic data. (A) the population model ~~that~~ consisted of ~~:-~~1) the burn-in period, where the number of generations was determined by the time necessary to contain the MRCA for all genomic regions. 2) the ~~sampling~~ inference period between the two time points, where the inference of demography and selection was made. (B) the genomic architecture model ~~that~~ consisted of ~~:-~~1) a diploid genome of one linkage group ~~that was divided in~~ into neutral and non-neutral regions ~~that were~~ composed of neutral and a combination of neutral and beneficial mutations.

statistics: 500, 5,000, and 10,000 bp. ~~Those windows~~ Windows overlapped because each ~~window~~ was composed around every ~~single SNP~~ which put SNP, putting the targeted variation in the middle of the window with ~~the other~~ other surrounding SNPs in half ~~of~~ the window size on each side of the targeted SNP. The site-frequency spectrum was obtained as a global summary statistics with three different numbers of discrete classes (bin sizes): 10, 15, and 20 bins (the complete list of summary statistics can be found in Supplementary Methods, section ~~S2~~ S1.1 List of summary statistics).

For every simulation, we combined a vector of summary statistics with the vector of X model parameters and the vector of five latent variables. Latent variables ~~represents values that~~ emerged represent values from the simulation or ~~values that~~ emerged by combining a latent variable and a model parameter. In our inferential framework, for example, the effective population size N_e is a latent variable ~~and it was~~ calculated within each simulation. The ratio between the effective population size N_e and the population census size N , N_e/N , on the other hand, was derived by combining a latent variable and a model parameter for each simulation. The other three latent variables were: the number of beneficial mutations under strong selection P , the average selection coefficient of strongly selected mutations \bar{s} , and the average substitution load L .

The effective population size N_e measures the increase of inbreeding at each generation. In this definition, N_e is the size of an ideal population with the same amount of drift as the population under consideration. Defined in these terms, N_e is the inbreeding effective size (Santiago and Caballero, 1995; Walsh and Lynch, 2018). It was calculated in every generation i of the sampling period as:

$$N_{e,i} = \frac{4N}{\sigma_{k_i}^2 + 2}$$

$\sigma_{k_i}^2$ being the variance among parents of the number of gametes produced that contributed to offspring in generation i . The N_e for the whole inference period was obtained by calculating

the harmonic mean of $N_{e,i}$. The population size of N was kept constant for the whole period, as shown above, ~~and it represents a parameter of the simulation representing a simulation parameter~~. From the N_e we obtained the ratio N_e/N (it measures how the census size reflects the actual effective population size: we expect to have a reduction on N_e compared to N when beneficial mutations are more pervasive).

We also recorded the selection coefficient of all beneficial mutations ~~that arose present~~ in every generation i from t_1 to t_2 in each simulation. After, we calculated the fraction of beneficial mutations that were strongly selected (where $s > 1/N_e$ over all mutations that were segregating in the period). This fraction represented all beneficial mutations ~~that arose in present~~ between t_1 and t_2 , regardless if they were lost or fixed at any generation of the period ~~or~~ if their frequency ~~oscillated fluctuated~~ but never reached fixation. We decided ~~for on~~ it because any beneficial mutation ~~that arise~~ can impact the allele frequency trajectories of other mutations (neutral or beneficial). For these mutations, we also calculated the average across all selection coefficients. We also calculated, in every generation of this period, the substitution load L_i as the difference between the total fitness of the individual with the highest fitness $W_{\max i}$ and ~~the~~ mean total fitness of the population \bar{W}_i (it measures the overall diversity of beneficial mutations present in the inference period),

$$L_i = \begin{cases} 0, & \text{if } W_{\max i} = 0 \\ \frac{W_{\max i} - \bar{W}_i}{W_{\max i}}, & \text{otherwise} \end{cases}$$

The average substitution load was obtained by averaging all values of L_i .

Implementation

The model was simulated with the software SLiM v3.1 (Haller and Messer, 2017; Haller et al., 2019). ~~For the calculation of~~ To calculate the inbreeding effective size, we needed to activate an optional SLiM 3.1 behavior to track the pedigrees of each individual in the population. It allowed us to obtain the number of each parent gamete and the population variance of the number of gametes. For ~~the calculation of~~ calculating the generation substitution load, we used a SLiM built-in function that allowed us to obtain the ~~vector of fitness~~ fitness vector of all individuals in the population. The cached fitness was the sum of all fitness determined by each beneficial mutation.

Each simulation was produced by using different combinations of the model's parameters: 1) the mutation rate per bp per generation μ , 2) the per-base recombination rate per generation ~~$\epsilon_0 r$~~ , 3) the mean γ of a gamma distribution (with the shape ~~parameters~~ parameter equal to the mean), from which the selection coefficients s of each beneficial mutation in the simulation were sampled, 4) the number of non-neutral genomic regions P_R , 5) the parameter that determines the probability of beneficial mutation in non-neutral regions P_B , 6) the population census size of the burn-in period N_0 , and, finally, 7) the population size of the inferential period N .

We set SLiM to output genotypic data of samples of individuals as single nucleotide polymorphisms (SNPs), at t_1 and t_2 , in the VCF file format. Using bcftools (Li, 2011), custom R function (R Core Team, 2020) and EggLib ~~(De Mita and Siol, 2012)~~ (Siol et al., 2022), SLiM outputs were processed and summary statistics calculated. We implemented a pipeline in an R script that automates the sampling of the prior values, runs each simulation, manipulates the VCF files, calculates the summary statistics, and organizes the final reference table. This script was also produced to facilitate the model test with a few simulations and the job submission in ~~a~~ an HPC node(s). The main R and additional scripts are available on Zenodo (Pavinato et al., 2021). In this pipeline, for every simulation, a row of the reference table was produced by combining the model parameters ~~used to launch a SLiM simulation~~, latent variables, and ~~the~~ summary statistics.

ABC-RF

In this work, we ~~take advantage of the use of~~ use Random Forests (RF) in ~~the~~ ABC procedure, where the parameter estimation is a machine learning problem (Pudlo et al., 2016; Raynal et al., 2019).

The performance of this approach was evaluated through simulations. First, we assumed a target dataset consisting of two samples of 100 individuals sampled ten generations apart from the same population. A reference table ~~was generated with for that target data was produced by~~ simulating the whole-genome SNPs of diploid individuals using the model described above ~~and calculating the previous summary statistics~~. At each simulation, we sampled 100 individuals at each time point and recorded their genotypes. Only polymorphic SNPs were retained for each sample. In each simulation, each individual had a genome ~~of size size of~~ 100 Mb ~~that was~~ Mbp divided into 2,000 fragments of 50,000 bps. A number of these fragments were randomly set as either neutral or non-neutral, based on the ~~parameter probability~~ P_R . ~~In the inference period, 100 individual genotypes were sampled at t_1 and t_2 after $\tau = 10$ generations.~~ For all model parameters, values of each simulation were sampled from a log-uniform distribution with range: 1 to 2,000 for N_0 and N , 10^{-10} to 10^{-6} for μ , 5×10^{-10} to 5×10^{-7} for r , 10^{-5} to 1 for P_B , and 10^{-3} to 1 for γ . ~~And Furthermore,~~ uniform distribution with range 0 to 1 for P_R (Figure S1 shows the prior distribution for all model parameters and latent values).

The raw reference table produced by the pipeline was processed to remove missing data. Missing data were present in several summary statistics of simulations with low genetic diversity ~~that can be produced, for example, by low mutation rate, by small population size, by selection, or by the combination of these parameters~~. Missing data were also present in the entire row of a simulation if the combination of population size, mutation, and especially recombination rate produced simulations that were memory intense, which caused the simulation to crash. A final reference table containing 55,634 simulations with 405 summary statistics was used to train the ABC-RFs. Independent RFs were obtained for each parameter and latent variable using R package abcrf (Pudlo et al., 2016; Raynal et al., 2019). Each RF was obtained by growing 1,000 trees. The ~~RF-RFs~~ were grown with the default parameters. Average genetic load, L , and P were ~~logit transformed prior logit transformed before~~ the training. For these latent variables and for \bar{s} , simulations with $L = 0$, $P = 0$ or $\bar{s} = 0$ were also excluded from the training set, which reduced it to 36,026 simulations for L , and with 29,264 simulations for P and \bar{s} . ~~For We performed log transformation before training for~~ the other parameters and latent variables ~~we performed log transformation prior training and use and used~~ the reference table containing all simulations.

The performance of each trained Random Forest was evaluated with *out-of-bag* (OOB) estimates (Breiman, 2001). ~~These estimates were produced by the trained model~~ The trained model produced these estimates for the data used for training. Regression trees that compose the actual RF are grown using part of the data ~~selected randomly from the initial set of simulations~~. ~~As a consequence~~ Consequently, for each simulation, ~~there is a subset of trees that were was~~ grown without the data from that simulation. The estimate from that subset of trees is called ~~OOB estimate~~ the OOB estimate, and with it, ~~cross-validation of the trained model is done validated~~ without splitting the reference table ~~in into~~ the training and testing sets. We calculated the mean squared error (MSE) and the correlation coefficient (R^2) between the true and the OOB estimated values obtained with the function regAbcrf implemented in the R package abcrf. For neutral simulations of the latent variables L , P , and \bar{s} , we evaluated the performance with the MSE and the bias on the parameters estimated in the original parameter scale.

An additional 1,000 simulations were used to evaluate the method's robustness to heterogeneous recombination rates along the genome. The simulation model was identical to the previously described simulations, except that a recombination map was used with varying recombination rates along the genome. We used the already implemented genomic fragmentation of the genome in "neutral" and "non-neutral" regions, which split the genome into 2,000 blocks of 50 Kbp, to

define the positions at which the recombination rate changed. Each corresponding fragment had a recombination rate sampled from a log-uniform distribution with a range between $10^{\log_{10} r - 0.5}$ and $10^{\log_{10} r + 0.5}$, with r sampled from the prior distribution as described above. This range allowed the genome to have recombination rates spanning one order of magnitude. We evaluate the RF performance in these simulations by calculating the mean squared error (MSE) and the correlation coefficient (R^2) between the true parameter values and the RF estimates. For neutral simulations of the latent variables L , P , and \bar{s} , we evaluated the performance with the MSE and the bias.

Alternative estimates of N_e from temporal data

We compared the ~~the~~ ABC-RF N_e estimates with estimates obtained with the global F_{ST} between temporal genomic samples (Frachon et al., 2017). This estimator is defined as:

$$\hat{N}_e = \frac{\tau(1 - \hat{F}_{ST})}{4\hat{F}_{ST}}$$

where τ accounts for the time-interval, in generations, between the first and the last samples used to estimate the F_{ST} , and \hat{F}_{ST} is the ~~the~~ Weir and Cockerham's F_{ST} estimator (Weir and Cockerham, 1984). The N_e from the F_{ST} was calculated for all simulations used to train the random forest. We calculated the mean squared error (MSE) and the squared correlation coefficient of linear regression (R^2) between the observed (true) and the F_{ST} -based N_e estimated values of all simulations. We also evaluated the performance of each estimator by calculating the MSE for simulations within a specific range of values of θ_b (local MSE estimates). By comparing the changes in MSE values of each estimator as a function of θ_b we could better understand how the amount of selection affected each estimator.

Analysis of temporal genomic data of feral populations of *Apis mellifera*

~~To give an example of an application of our framework, we analyzed a subset of~~ We used our framework to analyze the whole-genome sequencing data of feral populations of honey bees from California (Cridland et al., 2018). Eight out of fourteen sites in this work were composed of samples from museum and contemporary collections of freely foraging honey bees: 1) Avalon site in Catalina Island, Los Angeles ~~county~~ County, 2) Arcata and Blue Lake sites in Humboldt county, 3) Placerita Canion Nature Area in Los Angeles ~~county~~ County, 4) Sky Valley and Idyllwild in Riverside County, 5) La Grange, Stanislaus county, 6) Stebbins ~~Could~~ Cold Canyon Reserve, Solano county and 7) UC Davis Campus, Yolo county (Table 1). This dataset contains pairs spanning 104 years (as in the Avalon site, Catalina Island, Los Angeles county) and pairs spanning only 15 years (as in the Placerita Canyon Nature Area, Southern California, and Idyllwild, in Riverside county). For the temporal samples from Riverside County, we only used the two samples collected in May 1999 in Idyllwild as the first sample, ~~and~~. We combined all samples collected in September 2014 (in Idyllwild and Sky Valley) as the second sample (Table 1). Publicly available whole genomes fastq files for the contemporary and museum samples are available from the Sequence Read Archive (PRJNA385500) as described by Cridland et al. (2018); we performed the data analysis from VCF files (the same files used in Cridland et al., 2018) kindly provided by J. Cridland.

Individual VCF files of each ~~site~~ population were combined with bcftools (Li, 2011), and a custom R script was used to convert ~~the dataset to an~~ each dataset to the input format required to run an EggLib custom implementation (in Pavinato et al., 2021). We ~~tagged samples from the same time point with the same label~~ first produce simulated data to train the RF to apply our model to this targeted dataset. A reference table was produced by simulating whole-genome

SNPs for diploid individuals of *Apis mellifera*, changing three model parameters specifically for this targeted dataset: the sample size for population time points t_1 and t_2 , and the size of the haploid genome. For each population, we set the simulation to sample the same number of sequenced individuals from the pool of simulated individuals (as detailed in Table 1). For the simulations, Avalon population, for example, at t_1 and t_2 we set the genome size of each individual to simulation to sample two and five individuals apart $\tau = 104$ generations (assuming one generation/year). Only polymorphic SNPs were retained for each sample. We set the haploid genome size to 250 Mb (similar to the most recent estimates of *A. mellifera* genome size (Elsik et al., 2014)). For the analysis of the *Apis mellifera* temporal genomic data we Mb (similar to the most recent estimates of *A. mellifera* genome size; Elsik et al., 2014). We measured the amount of missing data, but we present in the original VCF files (Cridland et al., 2018) for each population. We found a negligible amount ($< 1\%$) in most of the populations (except populations from Avalon and Placerita that had 10% of the total missing genotypes), and we decided not to simulate missing data for any population analyzed.

The simulation model to generate the reference tables for the ABC analysis of *A. mellifera* populations was similar to the model described above but required some modification to adjust it to the specificities of the species and samples available. The simulated genome was divided into 5,000 fragments of 50,000 bps. These fragments were randomly set as neutral or non-neutral according to the parameter P_R . Dominance coefficients were set to 0.5 for all beneficial mutations throughout the simulation. In the sampling period, for each site, the same number of individuals found in each dataset were sampled in the first (t_1) and second time point (t_2). We used a Normal distribution for μ with a mean of 3.4×10^{-9} with a standard deviation of 0.5, to have a prior distribution center around the estimated mutation rate for Hymenoptera (Liu et al., 2017). The per base recombination rate was set as Uniform, ranging from 10^{-8} to 10^{-4} . The genome was represented with a single linkage group. For the represented the genome. The population sizes N_0 and N were taken from a Uniform prior distribution ranging from 1 to 10,000 individuals. Other prior probability distribution distributions of the parameters were set with the same prior as described above. Sample sizes and times were adjusted to match each population's population (see table 1). We used the same summary statistics as described above, except this time, we only calculated described above. However, we calculated only one window size of 10Kbp for summary statistics calculated in windows and one bin size of 10 bins for the site-frequency spectrum. The raw reference table containing the vector of parameters, latent variables, and summary statistics produced by the pipeline was processed to remove missing data, and a final reference table containing 162 summary statistics for each population pair was used to train the ABC-RFs. (see We visually assessed the model goodness-of-fit by performing a principal component analysis on the summary statistics of each population training reference table and projecting the corresponding PC of the target population reference table on the PCA plot. We consider a good model fit when the target population data point falls within the cloud of population simulated data points.

Like the ABC analyses described above, independent RFs were obtained for each parameter and latent variable using R package abcrf (Pudlo et al., 2016; Raynal et al., 2019). Each RF was obtained by growing 1,000 trees. The RFs were grown with the default parameters. Average genetic load, L , and P were logit-transformed before the training. For these latent variables and for \bar{s} , simulations with $L = 0$, $P = 0$ or $\bar{s} = 0$ were also excluded from the training set. We performed log transformation before training for the other parameters and latent variables and used the reference table containing all simulations. As before, we evaluate the RF performance by calculating the mean squared error (MSE) and the correlation coefficient (R^2) between the true and the OOB estimated values obtained with the function regAbcrf implemented in the R package abcrf. For neutral simulations of the latent variables L , P , and \bar{s} , we evaluated the performance with the MSE and the bias on the parameters estimated in the original parameter scale. See Table 1 for the number of simulations of each reference table). The RFs were grown

Table 1: Populations and number of simulations in the reference table.

Location	Date	N Sample Sizes	Simulations
Avalon , Catalina Island, Los Angeles county	1910/2014	2 5 /5	13,953
Blue Lake and Arcata, Humboldt county	1966/2015	6 5 /6	14,216
Placerita Canyon Nature Area, Los Angeles county	1999/2014	5 5 /6	14,125
Idyllwild and Sky Valey, Riverside county	1999/2014	2 5 /8	13,930
La Grange, Stanislaus county	1976/2014	2 5 /6	13,956
Stebbins Cold Canyon Reserve, Los Angeles county	1996/2014	5 5 /5	14,121
UC Davis Campus, Yolo county	1968/2015	2 5 /6	13,970

Names highlighted in bold letters ~~corresponds~~ correspond to the population code we used in this work. For sample sizes, the first value indicates the size of the first (older) sample and the second value the size of the second (contemporary) sample.

Results

~~ABC-RF framework for joint~~ Joint inference of adaptive and demographic history

The ~~ABC-RF framework jointly estimated~~ proposed framework allowed us to estimate parameters informative about adaptive and demographic history in temporal population genomics settings. Independent random forests estimated the population scaled beneficial mutation rate θ_b , the population census size N , and the effective population size N_e (Figure 2). Trained RFs performed well in predicting N and N_e with small MSE and higher R^2 (Figure 2 b and c). But, the trained RF for θ_b had a lower performance than the trained RFs for demographic parameters, with high MSE and low R^2 (Figure 2a ~~and b~~). Still, the estimates were robust for intermediate to higher values of θ_b . For the results of other model parameters and latent variables informative about demography and selection (see Figure S2 and section S2 Supplementary Results). Similar values of MSE and R^2 on true vs. RF estimated values (Figure S5 a, b, and c) indicated similar performance RF for θ_b , see S3 N, and N_e on simulations with heterogenous recombination rates (see Figure S4 for an example of how r could vary across the genome). For the results of other model parameters and latent variables for simulation with heterogenous recombination rate, see Figure S5 and Table S2, section S2 Supplementary Results.

The ~~automated~~ automatic selection of informative summary statistics is an important feature of ABC-RF. For each tree of a random forest, summary statistics were selected given its ability to split the data. How many times ~~a summary statistics was~~ summary statistics were selected in each RF informs us of their importance for ~~the inference of~~ predicting a given parameter. For the prediction of θ_b values, the RF picked more frequently statistics that reflect the heterogeneity of the genome, such as the 5% quantile of Tajima's D calculated in the second sample, with the kurtosis and skewness of F_{ST} and D_a calculated globally (Figure ~~S4~~ S6 e). The population size was trained with a combination of within and between sample summary statistics: F_{ST} and D_a , with their respective derived statistics ~~were~~ frequently selected (Figure ~~S5-S7~~ c). For N_e , summary statistics that inform about the cumulative divergence between samples as F_{ST} and D_a , were frequently selected (Figure ~~S5-S7~~ d).




Figure 2: Out-of-bag estimates of ABC-RF trained for the joint inference of demography and selection, and \hat{N}_e estimates from the temporal F_{ST} to compare with the ABC-RF -based \hat{N}_e estimates. (a) population ~~census size N_{scaled}~~ mutation rate of beneficial mutations θ_b ; (b) ~~effective~~ population census size $N_e N$; (c) effective ~~population scaled mutation rate of beneficial mutations θ_b size N_e~~ ; and (d) N_e from temporal F_{ST}

Comparison with F_{ST} method to estimate N_e

425 We compared our ABC-RF N_e estimates ~~with estimates obtained with~~ the temporal F_{ST} estimates (Frachon et al., 2017). The F_{ST} - ~~N_e -based \hat{N}_e~~ \hat{N}_e was more affected by the amount of selection in larger populations ~~Consequently, the~~ when selection is more efficient. The F_{ST} - N_e -based \hat{N}_e showed higher overall MSE ~~and lower R^2~~ compared to the ~~ABC-RF-based~~ ABC-based estimates (Figure 2c and d). ~~When the beneficial mutations were less frequent (either because of the mutation rate of the beneficial mutation, and because of larger population size), the ABC-RF and the temporal F_{ST} \hat{N}_e performed well and similarly regardless, regardless of the strength of selection, with the ABC-RF-based estimator with less error when the beneficial mutations were less frequent (low θ_b). However, the ABC-based estimator had less local MSE than the temporal F_{ST} -based estimator. However, when~~ When the frequency of selection ~~started to increase~~ is high, 435 the N_e estimator based on the temporal F_{ST} had dramatically higher error (Figure 3).

Analysis of temporal genomic data of feral populations of *Apis mellifera*

440 The projection of each population target data point (in black) into the cloud of the training data points (in grey) in the PCA plots revealed that each population model could capture some dimension of the observed genetic diversity (Figures S8-S14). However, some PCs showed the observed data point outside the simulated data cloud of points, indicating some model inadequacies, possibly because we did not include gene flow or admixture in our simulations. For




Figure 3: Local MSE of N_e estimates as a function of θ_b . The lines corresponds to the MSE on N_e estimates from ABC-RF and from temporal F_{ST} . Dashed lines correspond to local MSE estimated from pseudo-observed data (POD) with heterogeneous recombination rates along the genome.

the analysis of feral *A. mellifera* populations, we first grew independent RF for each parameter in each population. ~~All populations had the same~~ Despite the differences in time intervals between samples, all populations had a similar performance of the ABC-RF estimator for N_e , as they showed similar values of MSE and R^2 (Figure S8S15). For N , trained RF for Humboldt, Stebbins and Placerita performed similarly well, ~~and they had with~~ the lowest MSE and higher R^2 (Figure S7S16). For θ_b , Riverside had trained RF with the worst performance (Figure S6S17). Overall, both MSE and R^2 obtained with OOB estimates from simulated data for ~~Apis~~ *A. mellifera* dataset were comparable to these parameters obtained with OOB estimates for the simulated data ~~for in~~ the evaluation of the method.

Trained ~~RF~~ RFs for N and N_e were able to predict these parameters in all populations, as the inference of the mean posterior value and the posterior distribution differentiated from the mean prior value and distribution (Figure 4 b and c). For N , posterior distribution were wider than for N_e . Trained RF for θ_b , for all populations had a similar posterior mean, except for the Avalon population that had a peak at a lower value (Figure 4 a). ~~But~~ However, the posterior distributions were wider and ~~they~~ followed the prior distribution, making it difficult to ~~accurately~~ predict the posterior mean and variance in all populations accurately. It is possible to see together with the posterior mean estimates that the ABC-RF estimates for θ_b were concentrated in lower values (Table S2S3) in all populations. N_e were also lower, and N_e and N were similar. For the results of OOB estimates of other model parameters and latent variables ~~, and for and~~ posterior estimates for these parameters ~~see S3~~, see section S2 Supplementary Results.




Figure 4: Joint inference of demography and selection for feral *A. mellifera* populations. (a) the scale mutation rate of selected mutations θ_b , (b) the population census size N , (c) the effective population size N_e . Dashed and filled lines correspond to the prior and posterior distributions, respectively. See Table S3, Supplementary Results for mean and 95% credibility intervals.

Discussion

Separating demography from drift, and the inference of θ_b

With temporal population genomics data, we can see the evolution in “action” as opposed to single time-point population genomics data (Feder et al., 2021). Consequently, temporal data have more information about the ongoing process, which make them better for the understanding of making them interesting for understanding the complex short-term effects of the interaction between demography and selection (Buffalo and Coop, 2019; Dehasque et al., 2020; Williams and Pennings, 2020). When samples from more than two time points are available, the footprints correlations among allele frequency changes allow to separate the effects of drift and selection in allele frequencies changes can be separated (e.g., Buffalo and Coop, 2020; Feder et al., 2014) (e.g., Buffalo and Coop, 2020; Feder et al., 2014). Our results showed that two samples collected in at different time points were are sufficient for the inference of the genome-wide footprint of adaptive evolution and to separate the demography (population census size N) from drift (effective population size N_e).

It is important to stress that N_e , calculated as a latent variable, captures the feedback dynamics between drift and linked selection. Selection, either positive or negative, causes a deviation of N_e from N . The impact of selection on the genome can extend far from the target of selection since individuals that carry beneficial mutations have more chance to reproduce, and their beneficial mutations are more likely to be in the next-generation offspring (Walsh and Lynch, 2018). In this complex dynamic, with many loci under selection which creates a dynamic that cannot be easily described analytically, latent variables actually summarized the by-product obtained from simulations can summarize the by-product of drift and selection interactions. With our approach, \hat{N}_e quantifies the drift due both to demographic and selection dynamics processes, unaffected by the bias of outlier loci.

This genome-wide reduction in N_e is not captured when loci are assumed to evolve independently (as in Sheehan and Song, 2016, for example) (as in Sheehan and Song, 2016; Laval et al., 2019, for example). In contrast, the complexity of linked selection and the genome-wide effect of selection are taken into account by using individual-based simulations with using individual-based simulations with

the whole genome in an ABC approach.

Estimates of genetic load or other genome-wide parameters about selection are obtained when annotated genomic data is available, ~~(Henn et al., 2015)~~ (Henn et al., 2015), or by conducting experiments on crossing populations ~~(for the genetic load Plough (2016))~~ ~~(for the genetic load; Plough, 2016)~~. However, we ~~were able to obtain~~ obtained estimates of selection parameters only ~~with using~~ polymorphism data. ~~A similar approach was used by Laval et al. (2019) to estimate the number of selective sweeps affecting the human genome in the past 10,000 years. But they did not consider the feedback dynamics of selection and demography.~~ Differently, Buffalo and Coop (2020) measured the genome-wide signature of selection by estimating the covariance of allele frequencies at consecutive time points. ~~This~~ The allele frequency covariance matrix allowed the quantification of the genome-wide contribution of selection to the observed allele frequency changes, even when selection involved many loci of small effect. In this work, we estimated the population scale mutation rate of beneficial mutations θ_b , which informs about the diversity of beneficial mutations that existed in the population between the two time points ~~and the~~ potential speed of adaptation at the genome level (Hermisson and Pennings, 2017). These estimates reflect the potential number of beneficial mutations ~~present~~ between the two time points regardless of their impact as determined by their selection coefficients.

The variable importance plot of each parameter shows us the global importance of each summary ~~statistics~~ statistic in the trained Random Forests. For N_e , N , and θ_b summary statistics calculated from the distribution of locus-specific summary statistics -skewness, kurtosis, mean, variance, 5% and 95% quantiles were more frequently used. Summary statistics derived from the distribution of locus-specific calculated from all segregating loci in the genome inform about the heterogeneity that selection and drift produce genome-wide. For example, a *de novo* a beneficial mutation entered the simulation and was selected, ~~it~~ it left a signal of lower diversity around the region it was located. The genome, after selection, contained spots where diversity was high and where it was low, and this heterogeneity was captured by the distribution of locus-specific H_E , more specifically, the lower tail of the distribution where the ~~values of diversity~~ diversity values of the statistic were lower. The ~~use of the~~ covariance matrix of allele frequencies through Buffalo and Coop (2020) ~~can be considered time~~ (Buffalo and Coop, 2020) ~~can be used~~ as a summary statistic ~~for the the genome-wide heterogeneity that to capture additional information about the selection and drift left on the genome. It would be interesting to include when more than two temporal samples are available. Including~~ this matrix as summary statistics for further development of the method would be interesting.

Comparison with F_{ST} method to estimate N_e

We compared the N_e obtained with ABC-RF framework to the N_e obtained with F_{ST} estimator ~~(Skoglund et al., 2014; Frachon et al., 2017)~~ (Frachon et al., 2017). Overall, the F_{ST} -based N_e estimator performed poorly compared to the ~~ABC-RF-based~~ ABC-RF-based estimator. The lower performance ~~were was~~ caused by N_e values that were underestimated when beneficial mutations were more frequent (higher θ_b). Consequently, the N_e estimates from the temporal F_{ST} were strongly affected by selection. ~~When~~ Both estimators performed similarly when the selection was infrequent or rare, ~~both estimators performed similarly well, but with~~ but the ABC-RF estimator ~~with lower MSE had lower MSE than the F_{ST} one~~. Positive selection can increase the variance of allele frequency between samples taken ~~in at~~ different time points. When selection is infrequent or rare, drift determines most ~~of~~ allele frequency changes between samples, ~~but~~, Still, when selection is pervasive, selection dominates, which ~~cause~~ causes dramatic and rapid changes in allele frequency, increasing the variance between samples. N_e estimator based on the F_{ST} depends on the differences in allele frequencies between samples, ~~consequently;~~ consequently, it is naturally biased by strong and frequent selection. We can assume that the N_e estimator from ABC-RF was insensitive to the amount of selection since we trained the ABC-RF

with N_e values from the simulation. In our simulations, N_e was a latent variable that captured the deviation that selection imposed on the number of individuals able to reproduce (selected for); ~~it was not a biased by unaccounted factors~~ unaccounted factors did not bias it.

The amount of selection for $\theta_b \sim O(1)$ ~~$\theta_b > 1$~~ could be unrealistic in some organisms, but plausible in virus (Feder et al., 2014) and many arthropod species, with large N_e , which have larger population sizes ~~(except in eusocial insects that have vertebrate-like population sizes~~ Romiguier et al. (2014)). ~~In larger populations sizes selection acts also (except in eusocial insects that have ver~~
The selection also acts on weaker and milder beneficial mutations in larger populations. In those organisms, it might be unreasonable to assume mutation-drift equilibrium given the pervasive role of selection. Consequently, ~~any attempt~~ attempts to estimate demography parameters as N_e without properly accounting for the pervasive role of selection could be biased.

Analysis of temporal genomic data of feral populations of *Apis mellifera*

Overall, the performance of the ABC-RF for selection and demography inference was similar across populations despite the differences in sample size and age. For θ_b , Avalon and Humboldt populations had posterior probability distributions ~~very~~ similar to the prior, indicating that the analysis provides no additional information on this parameter. These two ~~population~~ populations also present low effective population size estimates, ~~which can reduce the signal of selection reducing the selection signal~~. For the rest of the populations, the posterior probability distribution of θ_b is tilted toward the higher values but without a clear peak differentiating the distribution from the prior. Still, lower θ_b values could be excluded. ~~This favours an interpretation in which~~ It favors the interpretation that selection was acting during the study period ~~but~~ without providing a precise ~~estimate of the parameter~~. ~~The information about the parameter estimate~~. The presence of selection in these analyses comes mainly from the heterogeneity of the polymorphism along the genome, ~~thus~~. Thus, for a thorough interpretation of the results, it is important to discuss other processes that have not been ~~modelled but that~~ modeled but could affect this signal. The studied bee populations in California show a mixture of Eastern and Western European ancestry, with some populations presenting African ancestry in the most modern samples Cridland et al. (2018). Different levels of African admixture along the genome could create some heterogeneity and affect the inference. However, in Placerita and Riverside, the populations with higher African ancestry ~~at present~~, present similar estimates of θ_b that populations with little or no African admixture. ~~Also, Humboldt population changes from having~~ The Humboldt population changed from predominately Western European ~~ancestry to having predominately to~~ Eastern European ancestry, ~~which means meaning~~ that there was substantial gene flow into the population. These results suggest that admixture does not dramatically affect the inference of selection but also highlights the importance of incorporating admixture in the future development of the approach. Other processes, such as recombination and mutation rate, might be heterogeneous along the genome. Our analysis of simulations with heterogeneous recombination rate suggest that the approach is robust to those. However, more complex models also seem necessary to fully capture the observed genetic diversity (see Figures S7-S13, section S2 Supplementary Results). Including additional factors (admixture, heterogeneity of recombination and mutation rate, and other forms of selection) could be key to obtaining models that fit the data better. Further developments of this approach should take them into account.

Our ABC-RF approach estimated N_e with the same order of magnitude of other N_e estimates obtained for hymenopterans (Zayed, 2004). Lower values of N_e might reflect the presence of admixture, either African admixture or admixture that occurred with domesticated lineages facilitated by changes in beekeeping practices (Cridland et al., 2018). Northern populations, especially from Humboldt County, shared similarities with bees from reared colonies (with higher Eastern European ancestry). Southern populations, as shown by Cridland et al. (2018), showed

a higher level of admixture with African lineages. Populations from the southernmost ~~eites~~
590 ~~(from cities~~ (Riverside County, Placerita, and Avalon, Los Angeles ~~county~~County) showed higher
genetic diversity ~~;~~ ~~but they than the others but~~ did not show the highest values of N_e . On the
other hand, the population of Stanislaus County had the highest value of N_e , possibly because
it had lower levels of admixture with domesticated lineages compared to the population from
Riverside, Placerita, Avalon, and Los Angeles ~~county~~counties.

595 We observed that N_e and N had similar estimates. We were aware that our simulation
model did not account for key characteristics of eusocial insect reproductive biology: the mo-
nopolization of reproduction by the queen and the division of labor. In ~~Honey~~honey bees, a
queen mates with more than one male (a process called polyandry)(~~Estoup et al., 1994~~), which
leads to a biased breeding sex ratio (~~Estoup et al., 1994~~). Assuming that only queens can repro-
600 duce in the colony, polyandry increases the variance in the number of parents ~~that contribute~~
~~contributing~~ to the offspring gene pool, which leads to a decrease in the N_e compared ~~respect~~ to
 N (Nomura and Takahashi, 2012). In our simulations, we only simulated ~~monogamous mating;~~
~~therefore~~~~panmictic random mating.~~ ~~Therefore,~~ the difference between estimates of N_e and N
only reflects the ~~action of selection~~~~selection action~~. Therefore N must be interpreted with
605 caution as it is probably reflecting more the total number of female breeders per generation
rather than the size of the population. Individual-based forward simulators ~~such~~ as SLiM allows
setting different mating schemes. It is possible to simulate the haplodiploidy, the cast system,
diocry, and sex ratio found in honey bees. These ~~modifications in the simulation could potentially~~
~~simulation modifications could~~ allow us to estimate N and other parameters that could ~~reflect~~
610 ~~better the biology of the species~~~~but it better reflect the species' biology, but that~~ was not the
focus of this work.

One possible explanation for the similarities between N_e and N estimates, thus, relies on
cast specialization and concentration of reproduction to one of few females in the colony. These
came to a cost of reduced N_e , which reduces the efficacy of selection (either positive or negative).
615 Bees are the few insect groups that show very small N_e potentially linked with the evolution
of eusociality (Romiguier et al., 2014). Knowing that lower N_e reduces the effectiveness of
selection, it is plausible to think that lower N_e is restricting the effects of mutation affecting
fitness to stronger beneficial mutations. Since these mutations are less frequent than weak or
mild mutations, their effects on N_e were small, which explains why N_e and N had values in
620 the same range. Low N_e and low θ_b pointed to a biological system limited where adaptation is
limited by the influx of adaptive mutations (Rousset et al., 2020).

Our ABC-RF framework also estimated the per-site mutation rate per generation μ (Sup-
plementary Results, S18). ~~For all populations, the~~ ~~The~~ mean posterior μ ~~for all populations~~
exceeds the mean ~~prior~~ μ ~~of the species~~. The higher estimated values we obtained might ~~reflect~~
625 ~~the be due to the higher true mutation rate but also reflect~~ recent admixture events between
these populations. ~~We did not model gene flow and admixture.~~ Modeling admixture could help
us correctly separate the effects of selection and drift since the introgression of African genes
might have biased some estimates of selection parameters.

Perspectives and Limitations

630 Our model is ~~very simplistic~~~~relatively simple~~, as it only considered the impact of beneficial
mutations, neglecting the effect of background selection and standing variation. Background se-
lection can mimic directional selection because they cause a similar pattern of diversity reduction
around the target of selection ~~-. However, it was recently shown that background selection only~~
~~mimics the classical sweep in simplistic models, where the deleterious mutation is localized in a~~
635 ~~specific region~~ (Stephan, 2010) ~~has been discussed for a long time. Much less has been discussed~~
~~about the patterns of background selection on temporal data and their differences with selective~~
~~sweeps.~~ Cvijović et al. (2018) showed that neutral alleles linked to less deleterious backgrounds

could quickly rise to high frequencies due to purifying selection, which could mimic the temporal signal of a selective sweep. However, if background selection is not localized to specific regions of the genome (Schrider, 2020). For more realistic scenarios, where the concentration of deleterious mutations varies across the genome, background selection does not behave as a classical hard sweep, the signal of selective sweeps will be distinct from the effects of background selection (Schrider, 2020). In an attempt to jointly accommodate the effect of demography and selection on the inference of N_e , Johri et al. (2020) modeled the effect of background selection and developed an ABC-based approach that jointly estimated the distribution of fitness effects and N_e . In their simulations, they modeled deleterious mutations and the classical hard sweep with the inclusion of beneficial mutations. They showed an unbiased estimate of N_e regardless of the presence of positive and negative selection. Future developments should include a more realistic genomic architecture where both negative and positive mutations can co-occur and explore different concentrations of deleterious mutations. In addition to that, further developments should explore not only scenarios of *de novo* mutations, but also selection acting on standing variation. This can be easily achieved with our pipeline and allows for a more general treatment of the selection of soft sweeps. The model can also be expanded to more complex demographic scenarios, including changes in population size and genetic exchange with external sources (migration). Including such admixtures will be key in the future development of this approach since it is also a source of heterogeneity in the genome and, thus, might influence the performance of the method.

Conclusion

We show that an ABC-RF -based approach is able to jointly infer adaptive and demographic history from temporal population genomics data. This approach allows the quantification of the genome-wide footprint of selection expressed in the scaled mutation rate of beneficial mutations. The ABC-RF N_e is robust to varying degrees of strength of selection and frequency of beneficial mutations. Our ABC-RF -based approach can be applied to temporal population genomics datasets to gain insight about the adaptive and demographic history of natural populations.

Acknowledgements

This project has received funding from the LabEx AGRO (convention ANR-10-LABX-0001-01), CEMEB (convention ANR-10-LABX-0004), and NUMEV (convention ANR-10-LABX-20) through the AAP Inter-LabEx (ABCSelection). This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 791695 (TimeAdapt). S. De Mita was funded by INRAE (Projet Innovant EFPA). We are thankful for the Genotoul bioinformatics platform Toulouse Occitanie (Bioinfo Genotoul, <https://doi.org/10.15454/1.5572369328961167E12>), for the High-Performance Computing Center University of Montpellier (MESO@LR-Platform) and for the Ohio Super Computer Center (OSC Center (1987)) for providing computing resources. The authors would like to thank Julie Cridland for sharing the processed data, and Andrew P. Michel for providing suggestions to improve an early version of the manuscript.

References

Bank, C., Ewing, G. B., Ferrer-Admettla, A., Foll, M., and Jensen, J. D. (2014). Thinking too positive? Revisiting current methods of population genetic selection inference. *Trends in*

Genetics, 30(12):540–546.

Bazin, E., Dawson, K. J., and Beaumont, M. A. (2010). Likelihood-free inference of population structure and local adaptation in a Bayesian hierarchical model. *Genetics*, 185(2):587–602.

685 Beichman, A. C., Huerta-Sanchez, E., and Lohmueller, K. E. (2018). Using genomic data to infer historic population dynamics of nonmodel organisms. *Annual Review of Ecology, Evolution, and Systematics*, 49(1):433–456.

Breiman, L. (2001). Random Forests. *Machine Learning*, 45(1):5–32.

690 Buffalo, V. and Coop, G. (2019). The linked selection signature of rapid adaptation in temporal genomic data. *Genetics*, 213(3):1007–1045.

Buffalo, V. and Coop, G. (2020). Estimating the genome-wide contribution of selection to temporal allele frequency change. *Proceedings of the National Academy of Sciences of the United States of America*, 2:201919039.

695 Casillas, S. and Barbadilla, A. (2017). Molecular population genetics. *Genetics*, 205(3):1003–1035.

Center, O. S. (1987). Ohio supercomputer center.

Cridland, J. M., Ramirez, S. R., Dean, C. A., Sciligo, A., and Tsutsui, N. D. (2018). Genome sequencing of museum specimens reveals rapid changes in the genetic composition of honey bees in California. *Genome Biology and Evolution*, 10(2):458–472.

700 Csilléry, K., Blum, M. G. B., Gaggiotti, O. E., and Francois, O. (2010). Approximate Bayesian Computation (ABC) in practice. *Trends in Ecology and Evolution*, 25(7):410–418.

Cvijović, I., Good, B. H., and Desai, M. M. (2018). The effect of strong purifying selection on genetic diversity. *Genetics*, 209(4):1235–1278.

705 De Mita, S. and Siol, M. (2012). EggLib: processing, analysis and simulation tools for population genetics and genomics. *BMC Genetics*, 13:27.

Dehasque, M., Ávila-Arcos, M. C., Díez-del Molino, D., Fumagalli, M., Guschanski, K., Lorenzen, E. D., Malaspinas, A.-S., Marques-Bonet, T., Martin, M. D., Murray, G. G. R., Papadopoulos, A. S. T., Therkildsen, N. O., Wegmann, D., Dalén, L., and Foote, A. D. (2020). Inference of natural selection from ancient DNA. *Evolution Letters*, 29:753–15.

710 Elsik, C. G., Worley, K. C., Bennett, A. K., Beye, M., Camara, F., Childers, C. P., de Graaf, D. C., Debyser, G., Deng, J., Devreese, B., Elhaik, E., Evans, J. D., Foster, L. J., Graur, D., Guigo, R., HGSC production teams, Hoff, K. J., Holder, M. E., Hudson, M. E., Hunt, G. J., Jiang, H., Joshi, V., Khetani, R. S., Kosarev, P., Kovar, C. L., Ma, J., Maleszka, R., Moritz, R. F. A., Munoz-Torres, M. C., Murphy, T. D., Muzny, D. M., Newsham, I. F.,
715 Reese, J. T., Robertson, H. M., Robinson, G. E., Rueppell, O., Solovyev, V., Stanke, M., Stolle, E., Tsuruda, J. M., Vaerenbergh, M. V., Waterhouse, R. M., Weaver, D. B., Whitfield, C. W., Wu, Y., Zdobnov, E. M., Zhang, L., Zhu, D., Gibbs, R. A., and Honey Bee Genome Sequencing Consortium (2014). Finding the missing honey bee genes: lessons learned from a genome upgrade. *BMC Genomics*, 15:86.

720 Estoup, A., the, M. S. P. o., and 1994 (1994). Precise assessment of the number of patriline and of genetic relatedness in honeybee colonies. *Proceedings of the Royal Society B*, 258:1–7.

Fay, J. C. and Wu, C.-I. (2000). Hitchhiking under positive darwinian selection. *Genetics*, 155(3):1405–1413.

Feder, A. F., Kryazhimskiy, S., and Plotkin, J. B. (2014). Identifying signatures of selection in genetic time series. *Genetics*, 196(2):509–522.

Feder, A. F., Pennings, P. S., and Petrov, D. A. (2021). The clarifying role of time series data in the population genetics of HIV. *PLoS Genetics*, 17(1):e1009050.

Frachon, L., Libourel, C., Villoutreix, R., Carrère, S., Glorieux, C., Huard-Chauveau, C., Navascués, M., Gay, L., Vitalis, R., Baron, E., Amsellem, L., Bouchez, O., Vidal, M., LE CORRE, V., Roby, D., Bergelson, J., and Roux, F. (2017). Intermediate degrees of synergistic pleiotropy drive adaptive evolution in ecological time. *Nature Ecology & Evolution*, 1(10):1551–1561.

Fraïsse, C., Popovic, I., Mazoyer, C., Spataro, B., Delmotte, S., Romiguier, J., LOIRE, E., Simon, A., Galtier, N., Duret, L., Bierne, N., Vekemans, X., and Roux, C. (2021). DILS: Demographic inferences with linked selection by using ABC. *Molecular Ecology Resources*.

Frazier, D. T., Martin, G. M., Robert, C. P., and Rousseau, J. (2018). Asymptotic properties of approximate Bayesian computation. *Biometrika*, 105(3):593–607.

Haller, B. C., Galloway, J., Kelleher, J., Messer, P. W., and Ralph, P. L. (2019). Tree-sequence recording in SLiM opens new horizons for forward-time simulation of whole genomes. *Molecular Ecology Resources*, 19(2):552–566.

Haller, B. C. and Messer, P. W. (2017). SLiM 2: flexible, interactive forward genetic simulations. *Molecular Biology and Evolution*, 34(1):230–240.

Henn, B. M., Botigué, L. R., Bustamante, C. D., Clark, A. G., and Gravel, S. (2015). Estimating the mutation load in human genomes. *Nature Reviews Genetics*, 16(6):333–343.

Hermisson, J. and Pennings, P. S. (2017). Soft sweeps and beyond: understanding the patterns and probabilities of selection footprints under rapid adaptation. *Methods in Ecology and Evolution*, 8(6):700–716.

Jensen, J. D., Kim, Y., DuMont, V. B., Aquadro, C. F., and Bustamante, C. D. (2005). Distinguishing between selective sweeps and demography using DNA polymorphism data. *Genetics*, 170(3):1401–1410.

Jensen, J. D., Thornton, K. R., Bustamante, C. D., and Aquadro, C. F. (2007). On the utility of linkage disequilibrium as a statistic for identifying targets of positive selection in nonequilibrium populations. *Genetics*, 176(4):2371–2379.

Johri, P., Charlesworth, B., and Jensen, J. D. (2020). Toward an evolutionarily appropriate null model: jointly inferring demography and purifying selection. *Genetics*, 215(1):173–192.

Karasov, T., Messer, P. W., and Petrov, D. A. (2010). Evidence that adaptation in drosophila is not limited by mutation at single sites. *PLoS Genetics*, 6(6):e1000924.

Kim, Y. and Nielsen, R. (2004). Linkage disequilibrium as a signature of selective sweeps. *Genetics*, 167(3):1513–1524.

Lange, J. D. and Pool, J. E. (2018). Impacts of recurrent hitchhiking on divergence and demographic inference in *Drosophila*. *Genome Biology and Evolution*.

Laval, G., Patin, E., Boutillier, P., and Quintana-Murci, L. (2019). A genome-wide Approximate Bayesian Computation approach suggests only limited numbers of soft sweeps in humans over the last 100,000 years. *bioRxiv*, 5:2019.12.22.886234.

765 Li, H. (2011). Improving SNP discovery by base alignment quality. *Bioinformatics*, 27(8):1157–1158.

Li, J., Li, H., Jakobsson, M., Li, S., Sjödin, P., and Lascoux, M. (2012). Joint analysis of demography and selection in population genetics: where do we stand and where could we go? *Molecular Ecology*, 21(1):28–44.

770 Lin, K., Li, H., Schlötterer, C., and Futschik, A. (2011). Distinguishing positive selection from neutral evolution: boosting the performance of summary statistics. *Genetics*, 187(1):229–244.

Liu, H., Jia, Y., Sun, X., Tian, D., Hurst, L. D., and Yang, S. (2017). Direct determination of the mutation rate in the bumblebee reveals evidence for weak recombination-associated mutation and an approximate rate constancy in insects. *Molecular Biology and Evolution*,
775 34(1):119–130.

Neher, R. A. (2013). Genetic draft, selective interference, and population genetics of rapid adaptation. *Annual Review of Ecology, Evolution, and Systematics*, 44(1):195–215.

Nei, M. and Tajima, F. (1981). Genetic drift and estimation of effective population size. *Genetics*, 98(3):625–640.

780 Nielsen, R. (2005). Molecular signatures of natural selection. *Annual Review of Genetics*, 39(1):197–218.

Nomura, T. and Takahashi, J. (2012). Effective population size in eusocial Hymenoptera with worker-produced males. *Heredity*, 109(5):261–268.

Pavinato, V., De Mita, S., and Navascués, M. (2021). Tracking-selection.
785 doi:10.5281/zenodo.4599736.

Plough, L. V. (2016). Genetic load in marine animals: a review. *Current Zoology*, 62(6):567–579.

Pool, J. E., Hellmann, I., Jensen, J. D., and Nielsen, R. (2010). Population genetic inference from genomic sequence variation. *Genome Research*, 20(3):291–300.

Pudlo, P., Marin, J.-M., Estoup, A., Cornuet, J. M., Gautier, M., and Robert, C. P. (2016).
790 Reliable ABC model choice via random forests. *Bioinformatics*, 32(6):859–866.

R Core Team (2020). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.

Raynal, L., Marin, J.-M., Pudlo, P., Ribatet, M., Robert, C. P., and Estoup, A. (2019). ABC random forests for Bayesian parameter inference. *Bioinformatics*, 35(10):1720–1728.

795 Romiguier, J., Lourenco, J., Gayral, P., Faivre, N., Weinert, L. A., Ravel, S., Ballenghien, M., Cahais, V., Bernard, A., Loire, E., Keller, L., and Galtier, N. (2014). Population genomics of eusocial insects: the costs of a vertebrate-like effective population size. *Journal of Evolutionary Biology*, 27(3):593–603.

800 Rousselle, M., Simion, P., Tilak, M.-K., Figuet, E., Nabholz, B., and Galtier, N. (2020). Is adaptation limited by mutation? A timescale-dependent effect of genetic diversity on the adaptive substitution rate in animals. *PLoS Genetics*, 16(4):e1008668.

Roux, C., Fraïsse, C., Romiguier, J., Anciaux, Y., Galtier, N., and Bierne, N. (2016). Shedding light on the grey zone of speciation along a continuum of genomic divergence. *PLOS Biol*, 14(12):e2000234–22.

805 Santiago, E. and Caballero, A. (1995). Effective size of populations under selection. *Genetics*, 139(2):1013–1030.

Schrider, D. R. (2020). Background selection does not mimic the patterns of genetic diversity produced by selective sweeps. *Genetics*, 216(2):499–519.

810 Schrider, D. R. and Kern, A. D. (2018). Supervised machine learning for population genetics: a new paradigm. *Trends in Genetics*, 34(4):301–312.

Schrider, D. R., Shanku, A. G., and Kern, A. D. (2016). Effects of linked selective sweeps on demographic inference and model selection. *Genetics*, 204(3):1207–1223.

Sella, G., Petrov, D. A., Przeworski, M., and Andolfatto, P. (2009). Pervasive natural selection in the Drosophila genome? *PLoS Genetics*, 5(6):e1000495.

815 Sheehan, S. and Song, Y. S. (2016). Deep learning for population genetic inference. *PLoS Computational Biology*, 12(3):e1004845.

Siol, M., Coudoux, T., Ravel, S., and De Mita, S. (2022). EggLib 3: A python package for population genetics and genomics. *Molecular Ecology Resources*.

820 Skoglund, P., Sjödin, P., Skoglund, T., Lascoux, M., and Jakobsson, M. (2014). Investigating population history using temporal genetic differentiation. *Molecular Biology and Evolution*, 31(9):2516–2527.

Stephan, W. (2010). Genetic hitchhiking versus background selection: the controversy and its implications. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365(1544):1245–1253.

825 Tajima, F. (1989). Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics*, 123(3):585–595.

Vitalis, R., Gautier, M., Dawson, K. J., and Beaumont, M. A. (2014). Detecting and measuring selection from gene frequency data. *Genetics*, 196(3):799–817.

830 Walsh, B. and Lynch, M. (2018). *Evolution and selection of quantitative traits*. Oxford University Press.

Weir, B. S. and Cockerham, C. C. (1984). Estimating F-Statistics for the analysis of population structure. *Evolution*, 38(6):1358.

835 Williams, K.-A. and Pennings, P. (2020). Drug resistance evolution in HIV in the Late 1990s: hard sweeps, soft sweeps, clonal interference and the accumulation of drug resistance mutations. *G3: Genes/Genomes/Genetics*, 10(4):1213–1223.

Zayed, A. (2004). Effective population size in Hymenoptera with complementary sex determination. *Heredity*, 93(6):627–630.