

Multiplicity and Koszul Homology

Lemma 0.1. *Let M be a finitely generated R -module and let I be an ideal of R . Then*

$$\sqrt{\operatorname{Ann}(M/IM)} = \sqrt{\langle I, \operatorname{Ann} M \rangle}.$$

Proof. To prove the equality on radicals, it suffices to show that a prime \mathfrak{p} of R contains $\operatorname{Ann}(M/IM)$ if and only if it contains $\langle I, \operatorname{Ann} M \rangle$. Recall that for any finitely generated R -module N , we have $V(\operatorname{Ann} N) = \operatorname{Supp} N$, or equivalently, $\mathfrak{p} \supseteq \operatorname{Ann} N$ if and only if $N_{\mathfrak{p}} \neq 0$. Thus since M is finitely generated (and hence M/IM is finitely generated too), we have

$$\begin{aligned} \mathfrak{p} \supseteq \operatorname{Ann}(M/IM) &\iff M_{\mathfrak{p}}/I_{\mathfrak{p}}M_{\mathfrak{p}} \neq 0 \\ &\iff M_{\mathfrak{p}} \neq 0 \text{ and } I_{\mathfrak{p}} \subseteq \mathfrak{p}_{\mathfrak{p}} \\ &\iff \mathfrak{p} \supseteq \operatorname{Ann} M \text{ and } I \subseteq \mathfrak{p} \\ &\iff \mathfrak{p} \supseteq \langle \operatorname{Ann} M, I \rangle \end{aligned}$$

□

Let $A = (A, \mathfrak{m}, \mathbb{k})$ be a noetherian local ring, let $\mathbf{x} = x_1, \dots, x_r$ be a sequence contained in \mathfrak{m} , and let M be a finitely generated A -module such that $\ell(M/\mathbf{x}M) < \infty$ (equivalently, we have $\mathfrak{m} = \sqrt{\operatorname{Ann}(M/\mathbf{x}M)}$). We set $K = K(\mathbf{x}, M)$ to be the Koszul complex with respect to \mathbf{x} and M and we denote its homology by $H_i(\mathbf{x}, M)$. Recall that the A -module $H_i(\mathbf{x}, M)$ is finitely generated and annihilated by $\langle \mathbf{x}, \operatorname{Ann} M \rangle$, hence they have finite length (indeed, we have $\mathfrak{m} = \sqrt{\operatorname{Ann}(M/\mathbf{x}M)} = \sqrt{\langle \mathbf{x}, \operatorname{Ann} M \rangle}$). We may therefore define the **Euler-Poincaré characteristic**

$$\chi(\mathbf{x}, M) = \sum_{i=0}^r (-1)^i \ell(H_i(\mathbf{x}, M)).$$

On the other hand, we the Hilbert-Samuel polynomial $P_{\mathbf{x}}(M)$ has degree $\leq r$, and we have

$$P_{\mathbf{x}}(M, n) = e_{\mathbf{x}}(M, r) \frac{n^r}{r!} + Q(n)$$

with $\deg Q < r$ and where $e_{\mathbf{x}}(M, r) = \Delta^r P_{\mathbf{x}}(M)$ is the Hilbert-Samuel multiplicity.

Theorem 0.2. *We have $\chi(\mathbf{x}, M) = e_{\mathbf{x}}(M, r)$.*

Proof. We prove this in several steps:

Step 1: To ease notation in what follows, we set $Q = \langle \mathbf{x} \rangle$. We first equip A with the standard Q -filtration $A = (Q^n)$ and view it as a filtered ring. Similarly, we equip M with the Q -filtration $M = (Q^n M)$ and view it as a filtered A -module. We now equip K with a Q -filtration as follows: for each $n \in \mathbb{N}$, let K^n be the R -subcomplex of K whose component in homological degree i

$$K_i^n = \begin{cases} Q^{n-i} K_i & \text{if } 0 \leq i < n \\ K_i & \text{else} \end{cases}$$

Thus for example, we have

$$\begin{aligned} K^0 &= M + \sum Me_i + \sum Me_{i,j} + \cdots \\ K^1 &= QM + \sum Me_i + \sum Me_{i,j} + \cdots \\ K^2 &= Q^2M + \sum QMe_i + \sum Me_{i,j} + \cdots \\ &\vdots \end{aligned}$$

Notice that

$$\begin{aligned} K^0/K^1 &= M/QM \\ K^1/K^2 &= QM/Q^2M + \sum (M/QM)e_i \\ K^2/K^3 &= Q^2M/Q^3M + \sum (QM/Q^2M)e_i + \sum (M/QM)e_{i,j} \\ &\vdots \end{aligned}$$

In particular, we clearly have

$$\begin{aligned} \mathrm{gr}(K) &= \bigoplus_{n=0}^{\infty} K^n/K^{n+1} \\ &= \mathrm{gr}(M) + \sum \mathrm{gr}(M)e_i + \sum \mathrm{gr}(M)e_{i,j} \\ &= K(\mathbf{x}, \mathrm{gr}(M)). \end{aligned}$$

Finally, we have

$$\begin{aligned} \chi(\mathbf{x}, M) &= \sum_{i=0}^r (-1)^i \ell(H_i(\mathbf{x}, M)) \\ &= \sum_{i=0}^r (-1)^i \ell(H_i(K/K^n)) \\ &= \sum_{i=0}^r (-1)^i \ell(K_i/K_i^n) \\ &= \sum_{i=0}^r (-1)^i \ell\left(\bigoplus_{\binom{r}{i}} M/\mathbf{x}^{n-i}M\right) \\ &= \sum_{i=0}^r (-1)^i \binom{r}{i} \ell(M/\mathbf{x}^{n-i}M) \\ &= e_{\mathbf{x}}(M, r). \end{aligned}$$

□

0.1 Extra

Let $(R, \mathfrak{m}, \mathbb{k})$ be a local noetherian ring of dimension n , let M be a nonzero finitely generated R -module of dimension d , and let $\mathbf{r} = r_1, \dots, r_d$ be a system of parameters for M . By definition, this means \mathbf{r} is a sequence contained in \mathfrak{m} such that $M/\mathbf{r}M$ has finite length, or equivalently, such that

$$\mathfrak{m} = \sqrt{\mathrm{Ann}(M/\mathbf{r}M)} = \sqrt{I+J} = \sqrt{Q},$$

where I set $I = \langle \mathbf{r} \rangle$, $J = \mathrm{Ann} M$, and $Q = I + J$ (so Q is \mathfrak{m} -primary). There's a beautiful formula (I think due to Serre) which expresses the Hilbert multiplicity of M with respect to Q as an alternating sum of lengths of Koszul homology modules. To explain this, first let me recall how the Hilbert multiplicity of M with respect to Q is defined: let (M_i) be any Q -stable filtration of M (for example, we can pick $M_i = I^i M = Q^i M$). Then the Hilbert-Samuel function with respect (M_i) is the function $f_{Q, (M_i)} = f: \mathbb{N} \rightarrow \mathbb{N}$ defined by

$$f(i) = \mathrm{length}_R(M/M_i) = \mathrm{length}_{R/Q}(M/M_i).$$

Note that each M/M_i is an R/Q -module since (M_i) is a Q -filtration, and since Q is \mathfrak{m} -primary, the lengths above are all finite. For i sufficiently large, we have $f(i) = P(i)$ where $P = P_{Q, M}$ is a polynomial whose lead coefficient is $e/d!$. Here, $e = e_{Q, M}$ is called the **Hilbert multiplicity** of M with respect to Q . It depends on the choice of Q (which itself depends on the choice of \mathbf{r} assuming M is fixed), however it doesn't depend on the choice of stable Q -filtration (M_i) .

The Euler-Poincaré characteristic with respect to \mathbf{r} and M is the alternating sum:

$$\chi(\mathbf{r}, M) = \sum_{i \geq 0} (-1)^i \mathrm{length}(H_i(\mathbf{r}, M)),$$

where $H(\mathbf{r}, M)$ is the homology of the Koszul complex $\mathcal{K}(\mathbf{r}, M) = \mathcal{K}(\mathbf{r}) \otimes_R M$. Note that

1. $H(\mathbf{r}, M)$ is a graded R/Q -module. In particular we have

$$H_0(\mathbf{r}, M) = M/\mathbf{r}M \text{ and } H_i(\mathbf{r}, M) = 0$$

for all $i \neq 0, 1, \dots, d$.

2. If \mathbf{r} is an R -sequence, then we have

$$H(\mathbf{r}, M) = \text{Tor}_R(R/I, M)$$

since $\mathcal{K}(\mathbf{r})$ is an R -free resolution of R/I in this case. If in addition we have $M = R/J$ and I and J are prime ideals of R , then

$$\chi(\mathbf{r}, M) = \chi(R/I, R/J) = \sum_{i \geq 0} (-1)^i \text{Tor}_i(R/I, R/J)$$

is called the **intersection multiplicity** of R/I and R/J . If $\dim(R/I) + \dim(R/J) = \dim R$, then it is an open conjecture that $\chi(R/I, R/J) > 0$.

3. If $H_1(\mathbf{r}, M) = 0$, then \mathbf{r} is an M -sequence, and this implies $H_i(\mathbf{r}, M) = 0$ for all $i > 0$.