

Fibers

Definition 0.1. Let S be an R -algebra and let \mathfrak{p} be a prime ideal of R . We define the **fiber of S over \mathfrak{p}** to be the $\kappa(\mathfrak{p})$ -algebra $\kappa(\mathfrak{p}) \otimes_R S$ where $\kappa(\mathfrak{p}) = K(R/\mathfrak{p})$ denotes the quotient field of R/\mathfrak{p} . In particular, if \mathfrak{m} is a maximal ideal of R , then the fiber of S over \mathfrak{m} is the R/\mathfrak{m} -algebra $R/\mathfrak{m} \otimes_R S \simeq S/\mathfrak{m}S$. If R is an integral domain with fraction field K , then **generic fiber of S** is the K -algebra $K \otimes_R S$.

Remark 1. Let $\iota: A \rightarrow B$ be an inclusion of \mathbb{k} -algebras where \mathbb{k} is a field. Geometrically speaking, the inclusion map $\iota: A \rightarrow B$ of \mathbb{k} -algebras corresponds to the morphism $\pi: Y \rightarrow X$ of affine \mathbb{k} -schemes, where $X = \operatorname{Spec} A$, $Y = \operatorname{Spec} B$, and where π is defined by

$$\pi(\mathfrak{q}) = A \cap \mathfrak{q}$$

for all primes \mathfrak{q} of B . If $\iota: A \rightarrow B$ is an integral extension, then π is surjective (this is referred to as the **lying over** property for integral extensions). Note that π is continuous with respect to the Zariski topology, for if $U := D(a)$ is an open subset of X where $a \in A$, then

$$\pi^{-1}(U) = \pi^{-1}(D(a)) = D(\iota(a)) := V.$$

In other words, we have $a \notin A \cap \mathfrak{q}$ if and only if $a \notin \mathfrak{q}$ for all primes \mathfrak{q} of B . Now, given a prime \mathfrak{p} of A , the fiber of $\pi: Y \rightarrow X$ over \mathfrak{p} , denoted $Y_{\mathfrak{p}}$, is the pullback of $\pi: Y \rightarrow X$ with respect to the morphism $\varepsilon_{\mathfrak{p}}: \operatorname{Spec}(\kappa(\mathfrak{p})) \rightarrow X$ where $\varepsilon_{\mathfrak{p}}$ is the morphism which corresponds to the \mathbb{k} -algebra homomorphism $A \rightarrow \kappa(\mathfrak{p})$. In particular, $Y_{\mathfrak{p}}$ is an affine \mathbb{k} -scheme and the \mathbb{k} -algebra which corresponds to $Y_{\mathfrak{p}}$ is $\kappa(\mathfrak{p}) \otimes_A B$, which is precisely how we defined the fiber of B over \mathfrak{p} in the first place.

Example 0.1. Let $R = \mathbb{k}[\mathbf{a}] = \mathbb{k}[a_1, a_2, a_3]$ and let $S = R[x]/f = R[x_1, x_2]/f$ where $f = a_1x_1^2 + a_2x_1x_2 + a_3x_2^2$. Also for $\alpha = (\alpha_1, \alpha_2, \alpha_3) \in \mathbb{k}^3$, we set

$$\mathfrak{m}_{\alpha} = \langle a_1 - \alpha_1, a_2 - \alpha_2, a_3 - \alpha_3 \rangle \quad \text{and} \quad f_{\alpha} = \alpha_1x_1^2 + \alpha_2x_1x_2 + \alpha_3x_2^2.$$

Then the fiber of S over \mathfrak{m}_{α} is the \mathbb{k} -algebra $S_{\alpha} := \mathbb{k}[\mathbf{a}, x]/f_{\alpha}$. Geometrically speaking, the inclusion map $\iota: R \rightarrow S$ of \mathbb{k} -algebras corresponds to the projection $\pi: Y \rightarrow X$ of affine \mathbb{k} -schemes, where $X = \operatorname{Spec} R$ and $Y = \operatorname{Spec} S$. The fiber of π over \mathfrak{m}_{α} is given by

$$\pi^{-1}(\{\mathfrak{m}_{\alpha}\}) = V(f_{\alpha}) = \operatorname{Spec}(S_{\alpha}).$$

Example 0.2. Let $R = \mathbb{k}[t]$, let $S = R[x]/\langle x^2 - t \rangle$, and let $\mathfrak{p}_{\tau} = \langle t - \tau \rangle$ where $\tau \in \mathbb{k}$. Then for $\tau \neq 0$, the fiber of S over \mathfrak{p}_{τ} is $\mathbb{k}[x]/\langle x^2 - \tau \rangle \cong \mathbb{k} \times \mathbb{k}$. The fiber over \mathfrak{p}_0 is $S_0 := \mathbb{k}[x]/\langle x^2 \rangle$. Finally, the fiber over the zero ideal $\langle 0 \rangle$ is $\mathbb{k}(t)[x]/\langle x^2 - t \rangle$, a field of degree 2 over the residue field $\kappa(\langle 0 \rangle) = \mathbb{k}(t)$. We see that for each prime \mathfrak{p} , the fiber over \mathfrak{p} is a vector space of dimension 2 over its residue field $\kappa(\mathfrak{p})$. In fact, S is a free R -module on the generators $(1, x)$. Thus $S \otimes_R N = N \oplus N$ for any R -module N , and it follows that S is flat.

Proposition 0.1. Let $\varphi: A \rightarrow B$ be a ring homomorphism and let \mathfrak{p} be a prime ideal of A . Let $f: Y \rightarrow X$ be the corresponding map of affine schemes where $Y = \operatorname{Spec} A$ and $X = \operatorname{Spec} B$. Then \mathfrak{p} is in the image of f if and only if the fiber of B over \mathfrak{p} is nonzero.

Proof. First note that if \mathfrak{q} is a prime of B that lies over \mathfrak{p} , then $\mathfrak{q}_{\mathfrak{q}}$ is a prime of $B_{\mathfrak{q}}$ that lies over $\mathfrak{p}_{\mathfrak{p}}$. Conversely, if \mathfrak{r} is a prime of $B_{\mathfrak{q}}$ that lies over $\mathfrak{p}_{\mathfrak{p}}$, then it must have the form $\mathfrak{r} = \mathfrak{q}_{\mathfrak{p}}$ for some prime \mathfrak{q} of B . Thus, by localizing at \mathfrak{p} if necessary, we may assume that $A = (A, \mathfrak{p}, \mathbb{k})$ is a local ring. Now if \mathfrak{q} is a prime of B that lies over \mathfrak{p} , then $\bar{\mathfrak{q}} := \mathfrak{q}/\mathfrak{p}B$ is a prime of $\bar{B} := B/\mathfrak{p}B$ which implies \bar{B} is nonzero. Conversely, if $\bar{B} \neq 0$, then there exists a prime \mathfrak{r} of \bar{B} , which must have the form $\mathfrak{r} = \bar{\mathfrak{q}} := \mathfrak{q}/\mathfrak{p}B$ for some prime \mathfrak{q} of B which necessarily lies over \mathfrak{p} . \square