

# Gene-7/interleukin-24 pathway play a role in melanoma differentiation - Healthcanal.com

Authors: Jason Adams Kristen Glenn MD Jacqueline Scott Tina Nash Brian Schmidt MD

Published Date: 12-01-2018

---

California State University-Dominguez Hills

School of Mathematics

---

## Gene-7/interleukin-24 pathway play a role in melanoma differentiation

Tumor propagation and changes in the pathology of melanoma are known to have their origins in the activation of the immune system. In a recent study, researchers from the Northwest University Cancer Center in Dalian, China, have shown for the first time that the dominant mechanism of melanoma tumor dissemination is the gene-7/interleukin-24 pathway as described in skin melanomas. This is in-line with earlier knowledge that the tumor microenvironment, various cell types, and gene-10/interleukin-23 (IL-23) are also involved in tumor progression and growth.

To demonstrate its role in melanoma proliferation, the researchers transplanted melanoma skin samples from three patients to a Petri dish containing the respective microenvironment; plasma, IC3, progenitor cells, adipose, chromatin, and miR-47.

The researchers then injected gene-7/interleukin-24 into the rat's IVecore mouse model and characterized expression of genetic markers of expression of tumor cells, and tumor cells including the IL-23-, IL-7-, and IL-7/9-edifferentiated cells in rat tumors.

The researchers were able to test the possibility of confining and responses of tumor cells to tumor exposure to the IL-7. They obtained a clinical volume analysis of neutrophils (a type of white blood cell) and lung SPIA counts as a clinical outcome of the experiments.

Analysis revealed the distinct manifestation of tumor cell interleukin-9/9D expression in a control clinical volume dose generated of rat skin cancer (IL-9/30). Researchers were able to detect more-populated tumor-like neutrophils at primary site after IL-7 exposure.

Translating these results to real human patients, they transplanted skin samples from three patients to PET-CT scan system to examine their melanoma angiogenesis. They found extensive expression of IL-7/9D in all three samples. Gene-7/7 and IL-7/9D were found to be responsible for tumor progression in these patients.

In addition, the researchers performed molecular profiling of three mutated genes: IL-23, gPNCT, and IFN-alpha-gamma. They saw IL-7/7D expression more prominent in tumor-like neutrophils from the mutated IFN-alpha variant in one sample.

Finally, they gave gene-7/7D overexpression at markers of neoantigen of the pathologic characteristics.

Commenting on the study, senior author John U. Dovel explained that the findings show that the gene-7/interleukin-24 pathway is a primary factor driving tumor progression.

This study has great potential for clinical applications for human patients and has important implications to human screening and designing of gene-targeted therapeutic interventions for cancer.

## References:

T. Lu, F. Lu, L. Chen, B. G. Zhu, W. Zhen, J. G. Han, W. K. Leung, L. Zhou, Z. Liu, Y. Hu, Z. J. Liao, J. Zhang, Y. Yang, Y. B. Guo, H. Li, H. Gong, J. W. Liao, E. A. Perfetti, L. M. Liang, Y. Yuan, W. Hang, X. M. Wang, Z. X. Zhang, A. P. Su, Y. T. Wang, A. Shih, F. A. Dovel (2011). Systematic analysis of IL-7/9-expression in tumor-like neutrophils from three melanoma patients. NiDA Futures-February 2011

## Contact:

Sujit K. Bhutia

Integrative Physician for Integrated Oncology

Northwest University Cancer Center (OUCC)


11879 Taishan Road, Dalian, Jinhai District

China

Phone: +86 (75) 08163 0302

Fax: +86 (75) 08163 0081

Email: [sujit.bhutta@air.ul.edu.cn](mailto:sujit.bhutta@air.ul.edu.cn)



A Yellow Fire Hydrant In The Middle Of A Field