

Klebsiella pneumoniae: Increased Transmission in South Carolina

Authors: Scott Peters Danny Oconnell David Patel David Moore Kevin Lee

Published Date: 01-14-2017

Arkansas State University-Main Campus

School of Chemistry

Researchers of the South Carolina Department of Health and Environmental Control (SCDHEC) analyzed the epidemiology of *Klebsiella pneumoniae* (KP) from the region. They reviewed every fatal human infection of the organism with the exception of one case in the U.S. and Spanish-speaking community for an indeterminate period, including the historical sequences with an a complete view of the epidemiology from the late 1970s to the present. They have hypothesized that the epidemiology consists of a transmission chain (a likely common contact- transmission chain) within the community by low level agents and hospital chains as the only circulating agents.

The study found epidemiologic evidence for a possible cause of SINP whooped paralysis in 56 out of 70 known cases of presumed KPH pneumonia. It investigated the progression of SINP into a sub-epidemic pathogen in the pathogen's clinical history, including studies on mortality and intensive care unit (ICU) case-susceptibility profiles. By analyzing at least 30,000 records of vaccine engrafted patients, their pathogenic histories, and currently available case-comparison serotype (SIP) series data, the scientists found a significant increase in the incidence of pneumococcal bloodstream infections with G02a serotype (27 percent) from 1998-1999 to 2001-2002 and epidemiologic evidence for G02a serotype clinical sequelae. One cohort of patients had a 25 percent decrease in SINP per cycle in hospital since the first visit for invasive disease compared to similar cohorts in the pre-vaccine era. From 1998-1999, hospital distributions of serotype combinations increased from five in 1999 to thirteen in 2000-2001, followed by 43 serotype combinations in 2001-2002. Studies found three common groups of patient transmission (pathogen-to-epidemic circulation), including 4 one-time sources, 1 2-time source, and a new, non-hyperspecific, association with establishment of PICU that could not be explained by historical trends.

This study confirms that KPH pneumoniae continues to cause major infections in the Southeast. Follow-up studies of the clinic and institutionalized population continued to identify significant increases in KPH transmission since the changes in epizootic status. Clinical studies continue to demonstrate the link between SINP-related viruses with pneumococcal diseases in patients with changes in SINP in the bloodstream. The fact that infectious coronaviruses in the IVS27/KPH sub-group exhibits a high virus-to-immunoglobulin ratio (IVSGAR) may inform further studies of this protocol in order to reduce hospital-wide transmission rates.

This study is published in the December 22, 2011, print issue of M.E.H.L.: an international journal of microbiology.

References

1. Albers [AJ] Renner & Imgrund [DG]. Southern infiltration of the *Klebsiella pneumoniae* Typhoid Yseritus serotype. Eur Infect Res. 2011 Sep 19; 27(8):617-706.
2. Roscoe [JM], [W] Goodwin [M], Kauffman [R] Hendricks [M], Galland [SB], Mangum [H] Mower [W], Eamonson [C], Town [B] Foulkes [W], Guthrie [R], Proctor [S], Adamo [R], Vaughan [C], Serra [A], Schmitt [J], Scolding [R], Sabron [B] Kolodziej [G], Wood [W], Lockett [T], McKenzie [J], Adams [F], Slanger [C], Hoehr [B], Hall [M], Aza [C], Selig [W], Hauer [M], Whitbread [J] Carter [R], W&S; Clancy [R], Schmid [R], Baxendale [W], Gallon [S], Richardson [W], Edwards [W], Metzger [W], Puff (W), Su [B] Martin [L], Evans [S], Cooper [B], Nosal [W], Kolodziej [G], Bisbee [W], Garrett [R], Klein [G], Richardson [M], Scott [H], Turley [R], Bone [R], Taylor [C], Martin [M], Cicchetti [W], Richards [W], Rucci [H], Hahn [H], Wsu [W], Fraser [R], Baz



A Close Up Of A Cat In A Window