

Volume 124 · Supplement 2 · December 2018

BCPT

Basic & Clinical Pharmacology & Toxicology

**2018 Americas Conference on Medical Imaging
and Clinical Research (AMICR 2018)
23–25 December 2018
Panama**

WILEY
Blackwell

BASIC & CLINICAL PHARMACOLOGY & TOXICOLOGY

www.bcpt.org

Editor:

Kim Brøsen, Odense, Denmark

Associate Editors:

Jens Lykkesfeldt, Copenhagen, Denmark; Ulf Simonsen, Aarhus, Denmark

Assistant Editors:

Troels Bergmann, Odense, Denmark
Anders A. Jensen, Copenhagen, Denmark
Niels Jessen, Aarhus, Denmark
Anton Pottegård, Odense, Denmark
Pernille Tveden-Nyborg, Copenhagen, Denmark
Petrine Wellendorph, Copenhagen, Denmark

MiniReview Editors:

Jesper Hallas, Odense, Denmark; Lisbeth Ehlert Knudsen, Copenhagen, Denmark; Eeva Moilanen, Tampere, Finland

Editorial Manager:

Henrik Horneberg, Odense, Denmark

Editorial Office:

Basic & Clinical Pharmacology & Toxicology, University of Southern Denmark, Clinical Pharmacology and Pharmacy, Winslowparken 19, 2nd floor, DK-5000 Odense C, Denmark. Tel. +45 6550-4237, email: hhorneberg@health.sdu.dk

Board of Directors:

Hans Bräuner-Osborne, Copenhagen, Denmark
Ulrik Gether, Copenhagen, Denmark
Rickard Malmström, Stockholm, Sweden
Jesper Bo Nielsen, Odense, Denmark
Karin Kramer Nielsen, Copenhagen, Denmark
Helle Northeved, Copenhagen, Denmark
Kristin Ólafsdóttir, Reykjavík, Iceland
Mimi Stokke Opdal, Oslo, Norway
Pesonen, Ullamari, Turku, Finland
Thomas Kongstad Petersen, Copenhagen, Denmark

Editorial Advisers:

Brünnner, Nils, Frederiksberg, Denmark
Cascorbi, Ingolf, Kiel, Germany
Chlopicki, Stefan, Krakow, Poland
Christen, Stephan, Berne, Switzerland
de Mey, Jo, Odense, Denmark
de Oliveira, Marcelo G, Campinas, Brazil
Dahl, Marja-Liisa, Stockholm, Sweden
Drago, Filippo, Catania, Italy
Du, Guan Hua, Beijing, P.R. China
Dusinska, Maria, Kjeller, Norway
Eriksson, Tommy, Lund, Sweden
Forsberg, Markus, Kuopio, Finland
Frøkjær, Sven, Copenhagen, Denmark
Garland, Chris, Bath, UK
Holme, Jørn, Oslo, Norway
Huupponen, Risto, Turku, Finland
Jacqz-Aigrain, Evelyne, Paris, France
Javitch, Jonathan A., New York, NY, USA
Karlsson, Mats, Uppsala, Sweden
Korpi, Esa, Helsinki, Finland

Liapi, Charis, Athens, Greece
Levy, Finn Olav, Oslo, Norway
Loft, Steffen, Copenhagen, Denmark
MacDonald, Ewen, Cuopio, Finland
Nicoletti, Ferdinando, Catania, Italy
Niemi, Mikko, Helsinki, Finland
Parnham, Michael J., Zagreb, Croatia
Patrignani, Paula, Chieti, Italy
Rosenkilde, Mette Marie, Copenhagen, Denmark
Schwartz, Thue, Copenhagen, Denmark
Spina, Eduardo, Messina, Italy
Stürmer, Til, Chapel Hill, NC, USA
Tamargo, Juan, Madrid, Spain
Tanus-Santos, Jose Eduardo, Ribeirão Preto, Brazil
Teixeira, Mauro M., Belo Horizonte, Brazil
Thürmann, Petra, Wuppertal, Germany
Tostes, Rita, São Paulo, Brazil
Vähäkangas, Kirsi, Kuopio, Finland
Vanhoutte, Paul M., Hong Kong
Villalón, Carlos M., Mexico D.F., Mexico
Zhang, Yongxiang, Beijing, China

Aims and Scope

Basic & Clinical Pharmacology & Toxicology is an independent journal, publishing original scientific research in all fields of toxicology, basic and clinical pharmacology. This includes experimental animal pharmacology and toxicology and molecular (-genetic), biochemical and cellular pharmacology and toxicology. It also includes all aspects of clinical pharmacology: pharmacokinetics, pharmacodynamics, therapeutic drug monitoring, drug/drug interactions, pharmacogenetics/-genomics, pharmacoepidemiology, pharmacovigilance, pharmacoconomics, randomized controlled clinical trials and rational pharmacotherapy. For all compounds used in the studies, the chemical constitution and composition should be known, also for natural compounds.

Copyright and Copying

Copyright © 2018 Nordic Association for the Publication of BCPT (former Nordic Pharmacological Society). Published by John Wiley & Sons Ltd. All rights reserved. No part of this publication may be reproduced, stored or transmitted in any form or by any means without the prior permission in writing from the copyright holder. Authorization to copy items for internal and personal use is granted by the copyright holder for libraries and other users registered with their local Reproduction Rights Organisation (RRO), e.g. Copyright Clearance Center (CCC), 222 Rosewood Drive, Danvers, MA 01923, USA (www.copyright.com), provided the appropriate fee is paid directly to the RRO. This consent does not extend to other kinds of copying such as copying for general distribution, for advertising and promotional purposes, for creating new collective works or for resale. Special requests should be addressed to: permissions@wiley.com

Disclaimer: The Publisher, Nordic Association for the Publication of BCPT (former Nordic Pharmacological Society) and Editors cannot be held responsible for errors or any consequences arising from the use of information contained in this journal; the views and opinions expressed do not necessarily reflect those of the Publisher, Nordic Association for the Publication of BCPT (former Nordic Pharmacological Society) and Editors, neither does the publication of advertisements constitute any endorsement by the Publisher, Nordic Association for the Publication of BCPT (former Nordic Pharmacological Society) and Editors of the products advertised. *Basic & Clinical Pharmacology & Toxicology* ISSN 1742-7843 (Online) is published by John Wiley & Sons Ltd, 9600 Garsington Road, Oxford OX4 2DQ, UK. Tel: +44 1865 776868; Fax: 44 1865 714591.

heated at 160 °C for 24 h, removed the upper layer of solution, collected the precipitate, washed by deionized water and ethanol to neutral, and dried at 160 °C for 6 h.

Results: The XRD results show that the resulting products are mainly depend on the pH value of solution. Pure phase of $\text{Bi}_5\text{O}_7\text{Br}$ can be obtained when the pH value is 11.0, and only BiOBr formed when the pH value is 1.0. The SEM and TEM images show that the synthesized $\text{Bi}_5\text{O}_7\text{Br}$ is a layered structural particles with four angles star-shape, and the average particle size is about 27 μm . According to the Uv-vis spectra test curves, the theoretical calculation results indicate that the band-gap width of $\text{Bi}_5\text{O}_7\text{Br}$ is 2.52 eV, which is much lower than that of TiO_2 (3.0 eV for rutile and 3.2 eV for anatase). After 5 and 20 minutes irradiation by visible light, the decolorization rate of Rhodamine solution by TiO_2 was 10 % and 22 % respectively, while that by $\text{Bi}_5\text{O}_7\text{Br}$ was 93 % and 99 % respectively.

Conclusions: Well dispersed $\text{Bi}_5\text{O}_7\text{Br}$ crystals with layered structure and tetragonal star shape have been obtained by simple hydrothermal synthesis. The $\text{Bi}_5\text{O}_7\text{Br}$ has good visible-light photocatalytic performance and is expected to be widely used in the fields of catalytic degradation, disinfection and sterilization.

Acknowledgements: This work was supported by the National Natural Science Foundation of China (51762023) and the Training Program of Outstanding Young Scientists in Jiangxi Province (20171BCB23070).

095 | Digital microfluidic biochip routing method considering contamination and washing capacity

Tingshen Lan¹; Xingquan Li²; Jianli Chen¹; Wenxing Zhu¹

¹Center for Discrete Mathematics and Theoretical Computer Science, Fuzhou University, Fuzhou, China; ²School of Mathematics and Statistics, Minnan Normal University, Zhangzhou, China

Objectives: As an emerging device in biochemistry, digital microfluidic biochips (DMFBs) can integrate a series of basic operations into a few square centimeter chips. By embedding a programmable electrode array on the chip, DMFBs can independently control the movement of each discrete sample droplet. For a complete bioassay on the DMFBs, it will be divided into a series of successive sub-problems to be performed separately. In the process of execution, most of electrodes would be shared by different sample droplets for the limited of the number of electrodes. Thus, contaminations caused by liquid residues among droplets are inevitable and lead to lethal errors in bioassays. To remove the contaminations, washing operation is essential to ensure the correctness of bioassay. However, existing works either ignored the continuity of sub-problems and clean the contaminations only within a single

sub-problem, or ignored the realistic capacity constraint of washing droplets and clean all contaminations only used a few of washing droplets. Moreover, simply deploying washing operations with droplet routing may increase the execution time of a bioassay, which is not feasible for timing-critical bioassay. To effectively remove contaminations and minimize the execution time, it is desirable to consider routing and washing all sub-problems simultaneously.

Methods: To effectively remove contaminations and minimize the execution time of a bioassay, we propose a contamination-aware routing method with realistic washing capacity constraint. Firstly, we present a top-down scheme to generate candidates of routing paths for each sub-problem, and then we construct a shortest-path model to select desirable routing solution for all sub-problems. With a decision diagram of droplets, we further propose an integer linear programming (ILP) formulation to compact the execution time of bioassay. Finally, a washing method considering realistic washing capacity is proposed to remove contaminations for all sub-problems.

Results: To evaluate the performance of simultaneous consideration of sub-problems, we compared the results obtained by simultaneous consideration with the results achieved by separately processing the sub-problems on the real-life benchmarks (e.g., In vitro diagnosis, Protein detection). Experimental results show that our proposed method significantly reduces 72% contaminations and saves 11% execution time of bioassays. Consequently, our proposed method improves both the sensitivity and efficiency of bioassays.

Conclusions: We presented a novel flow to solve the routing and washing problem with capacity constraint by simultaneous considering all sub-problems in DMFBs. Firstly, we presented an incremental iteration procedure to generate route candidates. Secondly, we proposed an optimal ILP to compact execution time. Finally, cleaning contamination by washing droplets with limited washing capacity is simultaneously considered for all sub-problems. Experimental results shown that our method can greatly reduce the number of contaminations and execution time of bioassays.

Acknowledgements: This work was supported by Empyrean and the National Natural Science Foundation of China (Grants No. 11501115, 61672005 and 11331003).