

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

Koebe's
algorithm

Rectangle
map

On random walks inside generalized quadrilaterals

Mikko Nummelin

email: `mikko.nummelin@tkk.fi`

Turku Analysis seminar, 2007-10-24

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms
Koebe's
algorithm

Rectangle
map

1 Riemann mapping theorem

2 Harmonic measure

3 Osculation algorithms
■ Koebe's algorithm

4 Rectangle map

Riemann mapping theorem

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

Koebe's
algorithm

Rectangle
map

- Riemann mapping theorem states that there always exists a conformal map from a Jordan-domain Ω , $0 \in \Omega$ onto the unit disk, fixing the origin:

$$w(\Omega) = \mathbb{D}(0; 1), \quad w(0) = 0.$$

Riemann mapping theorem

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms
Koebe's
algorithm

Rectangle
map

- Riemann mapping theorem states that there always exists a conformal map from a Jordan-domain Ω , $0 \in \Omega$ onto the unit disk, fixing the origin:

$$w(\Omega) = \mathbb{D}(0; 1), \quad w(0) = 0.$$

- Thus each Jordan domain is *conformally equivalent* to the unit disk.

Harmonic measure

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

**Harmonic
measure**

Osculation
algorithms

Koebe's
algorithm

Rectangle
map

- Suppose a Jordan region Ω , $0 \in \Omega$ and a continuous time random walk $\xi(t)$ starting from the origin.

Harmonic measure

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms
Koebe's
algorithm

Rectangle
map

- Suppose a Jordan region Ω , $0 \in \Omega$ and a continuous time random walk $\xi(t)$ starting from the origin.
- There is a probability measure defined on $\partial\Omega$ which tells the probability that random walk $\xi(t)$ exits on particular places on $\partial\Omega$. This probability measure is called *harmonic measure* of Ω respect to the origin and usually notated

$$\omega(0, \alpha, \Omega) = \int_{\alpha} \rho_0(\theta) d\theta, \quad \alpha \subset \partial\Omega, \theta \in [0, 2\pi).$$

Computing harmonic measure

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

**Harmonic
measure**

Osculation
algorithms

Koebe's
algorithm

Rectangle
map

- It can be proved via Kolmogorov's law of total probability and first step analysis that harmonic measure is conformally invariant.

Computing harmonic measure

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms
Koebe's
algorithm

Rectangle
map

- It can be proved via Kolmogorov's law of total probability and first step analysis that harmonic measure is conformally invariant.
- So, if

$$\gamma(\theta), \quad \theta \in [0, 2\pi)$$

is a parametrization of $\partial\Omega$ and

$$\mu(\theta), \quad \theta \in [0, 2\pi)$$

is the respective parametrization of $\partial\mathbb{D}(0; 1)$ under a conformal mapping $w : \Omega \rightarrow \mathbb{D}(0; 1)$, then the density function of harmonic measure can be computed as

$$\rho_0(\theta) = \frac{\mu'(\theta)}{2i\pi\mu(\theta)}.$$

Graphical features of the harmonic measure

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

**Harmonic
measure**

Osculation
algorithms

Koebe's
algorithm

Rectangle
map

- Nearby smooth sides appear as smooth tops in density function of the harmonic measure.

Graphical features of the harmonic measure

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

Koebe's
algorithm

Rectangle
map

- Nearby smooth sides appear as smooth tops in density function of the harmonic measure.
- Inward opening angles (convex polygons have only these) and far-away branches appear as zeroes or near-zero-regions in the density function. This similarity is due to the fact that in conformal mappings, far-away branches are "almost" the same as 0 radian inward opening angles at infinity.

Graphical features of the harmonic measure

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

Koebe's
algorithm

Rectangle
map

- Nearby smooth sides appear as smooth tops in density function of the harmonic measure.
- Inward opening angles (convex polygons have only these) and far-away branches appear as zeroes or near-zero-regions in the density function. This similarity is due to the fact that in conformal mappings, far-away branches are "almost" the same as 0 radian inward opening angles at infinity.
- Outward opening angles (nonconvex polygons have usually some of these), appear as singularities in the density function.

Graphical features - example

On random walks inside generalized quadrilaterals

Mikko Nummelin

Outline

Riemann mapping theorem

Harmonic measure

Osculation algorithms
Koebe's algorithm

Rectangle map





Figure: A "horseshoe"-like region with its conformal center located in the origin, pictured inside the unit disk (left) and density function of its harmonic measure (right). Note, how the only visible features in the density function are lower left corner (two smooth "hills" with steep "valley" down to zero between them) and nearby outward-pointing corner (singularity).

Graphical features - example 2

On random walks inside generalized quadrilaterals

Mikko Nummelin

Outline

Riemann mapping theorem

Harmonic measure

Osculation algorithms

Koebe's algorithm

Rectangle map





Figure: Same region with conformal center (starting point of the random walk) transformed. Note, how this changes absorption probabilities on the edges.

Osculation algorithms

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

**Osculation
algorithms**

Koebe's
algorithm

Rectangle
map

- In order to construct the harmonic measure, it is necessary to find the unit disk map:

$$|\mu(\theta)| = 1, \quad \theta \in [0, 2\pi).$$

Osculation algorithms

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

Koebe's
algorithm

Rectangle
map

- In order to construct the harmonic measure, it is necessary to find the unit disk map:

$$|\mu(\theta)| = 1, \quad \theta \in [0, 2\pi).$$

- Here we will compute it via *osculation algorithms* and concentrate on *logarithmic Koebe's algorithm*. It is not the best or most accurate algorithm for all situations, but it is quite stable.

Osculation algorithms

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

Koebe's
algorithm

Rectangle
map

- In order to construct the harmonic measure, it is necessary to find the unit disk map:

$$|\mu(\theta)| = 1, \quad \theta \in [0, 2\pi).$$

- Here we will compute it via *osculation algorithms* and concentrate on *logarithmic Koebe's algorithm*. It is not the best or most accurate algorithm for all situations, but it is quite stable.
- Osculation algorithms are, in general, designed to pick up a feature from approximation of boundary of a Jordan-region $\partial\Omega$ and decide a mapping function accordingly to find a conformal image of Ω closer to the unit disk.

Koebe's algorithm

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

**Koebe's
algorithm**

Rectangle
map

- In (quadratic) Koebe's algorithm the extremal point is taken as $z_e = \min(\vec{z})$.

Koebe's algorithm

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

**Koebe's
algorithm**

Rectangle
map

- In (quadratic) Koebe's algorithm the extremal point is taken as $z_e = \min(\vec{z})$.
- After the minimum is found, a rotation coefficient is computed:

$$z_R = \frac{-\overline{z_e}}{|z_e|}$$

Koebe's algorithm

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

**Koebe's
algorithm**

Rectangle
map

- In (quadratic) Koebe's algorithm the extremal point is taken as $z_e = \min(\vec{z})$.
- After the minimum is found, a rotation coefficient is computed:

$$z_R = \frac{-\overline{z_e}}{|z_e|}$$

- Pointwise multiplication

$$\vec{z} \rightarrow z_R \vec{z}, \quad z_e \rightarrow z_R z_e$$

which rotates the extremal (minimum) point to negative real axis, is applied.

Koebe's algorithm

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

**Koebe's
algorithm**

Rectangle
map

- In (quadratic) Koebe's algorithm the extremal point is taken as $z_e = \min(\vec{z})$.
- After the minimum is found, a rotation coefficient is computed:

$$z_R = \frac{-\overline{z_e}}{|z_e|}$$

- Pointwise multiplication

$$\vec{z} \rightarrow z_R \vec{z}, \quad z_e \rightarrow z_R z_e$$

which rotates the extremal (minimum) point to negative real axis, is applied.

- A linear fractional transformation

$$\vec{z} \rightarrow \frac{\vec{z} + |z_e|}{1 + |z_e| \vec{z}}$$

is applied.

Koebe's algorithm - continued

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

**Koebe's
algorithm**

Rectangle
map

- The latter transformation ensured that the Jordan-curve approximated by \vec{z} passes through the origin but does not enclose it inside. This guarantees existence of continuous, holomorphic square root.

Koebe's algorithm - continued

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

**Koebe's
algorithm**

Rectangle
map

- The latter transformation ensured that the Jordan-curve approximated by \vec{z} passes through the origin but does not enclose it inside. This guarantees existence of continuous, holomorphic square root.
- Holomorphic square root

$$\vec{z} \rightarrow \sqrt{\vec{z}}$$

is applied so that first nonzero element in \vec{z} is transformed into lower half plane and for following points the branch of square root minimizing the distance to previous point's image is chosen.

Koebe's algorithm - continued

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

**Koebe's
algorithm**

Rectangle
map

- The latter transformation ensured that the Jordan-curve approximated by \vec{z} passes through the origin but does not enclose it inside. This guarantees existence of continuous, holomorphic square root.
- Holomorphic square root

$$\vec{z} \rightarrow \sqrt{\vec{z}}$$

is applied so that first nonzero element in \vec{z} is transformed into lower half plane and for following points the branch of square root minimizing the distance to previous point's image is chosen.

- The previous transformation sent all points of \vec{z} nearer the edge of unit disk but not outside it.

Koebe's algorithm - fixing the origin

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

**Koebe's
algorithm**

Rectangle
map

- Finally origin's image is fixed back to its original location by linear fractional transformation

$$\vec{z} \rightarrow \frac{\vec{z} - \sqrt{|z_e|}}{1 - \sqrt{|z_e|}\vec{z}}$$

Koebe's algorithm - fixing the origin

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

**Koebe's
algorithm**

Rectangle
map

- Finally origin's image is fixed back to its original location by linear fractional transformation

$$\vec{z} \rightarrow \frac{\vec{z} - \sqrt{|z_e|}}{1 - \sqrt{|z_e|}\vec{z}}$$

- and optionally the rotation can also be reversed

$$\vec{z} \rightarrow \overline{z_R \vec{z}}.$$

for better visual effect.

Koebe's algorithm - summary

On random
walks inside
generalized
quadrilater-
als

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

**Koebe's
algorithm**

Rectangle
map

So, one iteration of Koebe's algorithm is:

$$z_e = \min(\vec{z})$$

$$z_R = \frac{-\overline{z_e}}{|z_e|}$$

$$\vec{z} \rightarrow z_R \vec{z}$$

$$\vec{z} \rightarrow \frac{\vec{z} + |z_e|}{1 + |z_e| \vec{z}}$$

$$\vec{z} \rightarrow \sqrt{\vec{z}}$$

$$\vec{z} \rightarrow \frac{\vec{z} - \sqrt{|z_e|}}{1 - \sqrt{|z_e|} \vec{z}}$$

$$\vec{z} \rightarrow \overline{z_R} \vec{z}$$

Logarithmic Koebe's algorithm

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

**Koebe's
algorithm**

Rectangle
map

- If, instead of taking square root in Koebe algorithm, a cube root or higher is taken, it accelerates convergence.

Logarithmic Koebe's algorithm

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

**Koebe's
algorithm**

Rectangle
map

- If, instead of taking square root in Koebe algorithm, a cube root or higher is taken, it accelerates convergence.
- As a limit formula

$$\ln(z) = \lim_{n \rightarrow \infty} n \left(\sqrt[n]{z} - 1 \right)$$

holds, this suggests replacing the root with a holomorphic logarithm, taking \vec{z} into left half plane.

Logarithmic Koebe's algorithm

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

Koebe's
algorithm

Rectangle
map

- If, instead of taking square root in Koebe algorithm, a cube root or higher is taken, it accelerates convergence.
- As a limit formula

$$\ln(z) = \lim_{n \rightarrow \infty} n (\sqrt[n]{z} - 1)$$

holds, this suggests replacing the root with a holomorphic logarithm, taking \vec{z} into left half plane.

- After taking the logarithm, image of the origin lies somewhere in left half plane. It can be transformed to -1 by scaling and then a *Cayley transformation*

$$\vec{z} \rightarrow \frac{1 + \vec{z}}{1 - \vec{z}}$$

maps the transformed area back into unit disk.

Logarithmic Koebe's algorithm - summary

So, one iteration of logarithmic Koebe's algorithm is:

$$z_e = \min(\vec{z})$$

$$z_R = \frac{-\overline{z_e}}{|z_e|}$$

$$\vec{z} \rightarrow z_R \vec{z}$$

$$\vec{z} \rightarrow \frac{\vec{z} + |z_e|}{1 + |z_e| \vec{z}}$$

$$\vec{z} \rightarrow \ln \vec{z}$$

$$\vec{z} \rightarrow \frac{-\vec{z}}{\ln(|z_e|)}$$

$$\vec{z} \rightarrow \frac{1 + \vec{z}}{1 - \vec{z}}$$

$$\vec{z} \rightarrow \overline{z_R} \vec{z}$$

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

**Koebe's
algorithm**

Rectangle
map

The Schwarz-Christoffel integration formula

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

Koebe's
algorithm

Rectangle
map

- Suppose that Ω is a polygon, having n internal angles of sizes $\alpha_1, \alpha_2 \dots \alpha_n$ respectively.

The Schwarz-Christoffel integration formula

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

Koebe's
algorithm

Rectangle
map

- Suppose that Ω is a polygon, having n internal angles of sizes $\alpha_1, \alpha_2 \dots \alpha_n$ respectively.
- According to the Riemann mapping theorem, Ω is conformally equivalent to the unit disk.

The Schwarz-Christoffel integration formula

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms
Koebe's
algorithm

Rectangle
map

- Suppose that Ω is a polygon, having n internal angles of sizes $\alpha_1, \alpha_2 \dots \alpha_n$ respectively.
- According to the Riemann mapping theorem, Ω is conformally equivalent to the unit disk.
- The function defined by

$$w(z) = A + B \cdot \int_0^z \prod_{k=1}^n \left(1 - \frac{\zeta}{\zeta_k}\right)^{\frac{-\alpha_k}{\pi}} d\zeta, \quad \sum_{k=1}^n \alpha_k = 2\pi.$$

where A , B and each ζ_k are parameters, maps unit disk onto Ω .

Rectangle map

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms
Koebe's
algorithm

Rectangle
map

- The special case of Schwarz-Christoffel formula

$$w(z) = \int^z \left(1 - \frac{\zeta}{\zeta_1}\right)^{-1/2} \left(1 - \frac{\zeta}{\zeta_2}\right)^{-1/2} \dots \\ \dots \left(1 - \frac{\zeta}{\zeta_3}\right)^{-1/2} \left(1 - \frac{\zeta}{\zeta_4}\right)^{-1/2} d\zeta$$

maps unit disk onto a rectangle. It can be used to construct module of a generalized quadrilateral or potential lines of a capacitor.

Rectangle map

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms
Koebe's
algorithm

Rectangle
map

- The special case of Schwarz-Christoffel formula

$$w(z) = \int^z \left(1 - \frac{\zeta}{\zeta_1}\right)^{-1/2} \left(1 - \frac{\zeta}{\zeta_2}\right)^{-1/2} \dots \\ \dots \left(1 - \frac{\zeta}{\zeta_3}\right)^{-1/2} \left(1 - \frac{\zeta}{\zeta_4}\right)^{-1/2} d\zeta$$

maps unit disk onto a rectangle. It can be used to construct module of a generalized quadrilateral or potential lines of a capacitor.

- Here we will answer the question: if a random walk starts at some place inside Ω , where is the boundary from which it is most likely the random walk to end to either side.

Gaier quadrilateral

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms
Koebe's
algorithm

Rectangle
map

- Consider the so-called Gaier polygon $[0, 2, 2 + i, 1 + i, 1 + 2i, 2i]$ as a generalized quadrilateral, where $[2i, 0]$ and $[2, 2 + i, 1 + i, 1 + 2i]$ are generalized left and right sides of it, respectively.

Gaier quadrilateral

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms

Koebe's
algorithm

Rectangle
map

- Consider the so-called Gaier polygon $[0, 2, 2 + i, 1 + i, 1 + 2i, 2i]$ as a generalized quadrilateral, where $[2i, 0]$ and $[2, 2 + i, 1 + i, 1 + 2i]$ are generalized left and right sides of it, respectively.
- Construct a rectangle map of it to $[0, 1, 1 + iM, iM]$.

Gaier quadrilateral

On random
walks inside
generalized
quadrilaterals

Mikko
Nummelin

Outline

Riemann
mapping
theorem

Harmonic
measure

Osculation
algorithms
Koebe's
algorithm

Rectangle
map

- Consider the so-called Gaier polygon $[0, 2, 2 + i, 1 + i, 1 + 2i, 2i]$ as a generalized quadrilateral, where $[2i, 0]$ and $[2, 2 + i, 1 + i, 1 + 2i]$ are generalized left and right sides of it, respectively.
- Construct a rectangle map of it to $[0, 1, 1 + iM, iM]$.
- Find the reverse image of line $[0.5, 0.5 + iM]$ on the Gaier polygon and we have the middle equipotential line. A random walk starting on it will be as likely to hit the left side first than the right side.

Equipotential surface of Gaier quadrilateral

On random walks inside generalized quadrilaterals

Mikko Nummelin

Outline

Riemann mapping theorem

Harmonic measure

Osculation algorithms
Koebe's algorithm

Rectangle map




Figure: Gaier polygon as a generalized quadrilateral with middle equipotential line calculated. Note how this is to the left from geometric middle line. This is due to the lower right hand side "cave" as an attractor.



Milton Abramowitz, Irene A. Stegun: *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, Dover, New York, 1964

<http://www.math.sfu.ca/~cbm/aands/>



Lars V. Ahlfors: *Complex analysis, An Introduction to the Theory of Analytic Functions of One Complex Variable*, New York, Toronto, London, McGraw-Hill Book Company, Inc., 1953



Glen D. Anderson, Mavina K. Vamanamurthy, Matti K. Vuorinen: *Conformal invariants, inequalities, and quasiconformal maps*, John Wiley & Sons, Inc., New York, 1997



Lehel Banjai: *Revisiting the crowding phenomenon in Schwarz-Christoffel mapping*, Institut für Mathematik, Universität Zürich, 2007

-  Daniel J. Bernstein: *Computing logarithm intervals using arithmetic-geometric mean iteration*, National Science Foundation, Preprint, 2003
<http://cr.yp.to/papers.html#logagm>
-  Christopher J. Bishop: *Conformal welding and Koebe's theorem*, Preprint, 2003
-  Tobin A. Driscoll: *Schwarz-Christoffel Toolbox*
<http://www.math.udel.edu/~driscoll/software/SC>
-  Tobin A. Driscoll, Lloyd N. Trefethen: *Schwarz-Christoffel mapping*, Cambridge monographs on applied and computational mathematics, 2002
-  Dieter Gaier: *Ermittlung des konformen Moduls von Vierecken mit Differenzenmethoden*, Numer. Math. 19, 179-194, Springer-Verlag, 1972.



Theodore W. Gamelin: *Complex analysis*, Springer-Verlag, New York, Inc., 2001



John B. Garnett, Donald E. Marshall: *Harmonic measure*, Cambridge University Press, 2005



Ville Heikkala, Mavina K. Vamanamurthy, Matti Vuorinen: *Generalized elliptic integrals*, Preprint, 2007.
[arXiv:math/0701436](https://arxiv.org/abs/math/0701436)



Louis Hill Howell: *Computation of conformal maps by modified Schwarz-Christoffel transformations*, PhD Thesis, Massachusetts Institute Of Technology, 1985.



Stig Larsson, Vidar Thomée: *Partial Differential Equations with Numerical Methods*, Springer-Verlag, 2003.



PlanetMath: *Automorphisms of unit disk*
<http://planetmath.org/encyclopedia/AutomorphismsOfU>



PlanetMath: *Schwarz-Christoffel transformation (circular version)*

[http://planetmath.org/encyclopedia/
SchwarzChristoffelTransformationCircularVersion.htm](http://planetmath.org/encyclopedia/SchwarzChristoffelTransformationCircularVersion.htm)



R. Michael Porter: *History and Recent Developments in Techniques for Numerical Conformal Mapping*, Preprint, 2005.

http://www.math.utu.fi/projects/madras/w_proc_mike



Roland A. Schinzing, Patricio A. A. Laura: *Conformal Mapping - Methods And Applications*, Dover Publications Inc, Mineola, New York, 2003



Theodore Theodorsen: *Theory of wing sections of arbitrary shape*, NACA Report No. 411, 1933

<http://naca.central.cranfield.ac.uk/reports/1933/naca-r411>



Lloyd N. Trefethen, Tobin A. Driscoll: *Schwarz-Christoffel mapping in the computer era*, Proceedings of the International Congress of Mathematicians, Vol. III, 533-542.



Guangyu Zou, Jing Hua, Xianfeng Gu, Otto Muzik: *An approach for intersubject analysis of 3D brain images based on conformal geometry*

<http://www.cs.sunysb.edu/~vislab/papers/ICIP-2873.p>