G11: The adapter on the early steps only

UHz6 question 3 · status: answered

THE REVIEWER'S POINT

UHz6, question 3

- Given the finding in Sec. 3, does restricting the adapter to the early steps change the failure modes in Fig. 7?

U-Q3

Reviews.pdf, page 3 (no text layer; passage located by OCR)

these failures

THE SUBMITTED WEAKNESS

Figure 7, limitations


a)
Cat and dog
Tiger and dog
Chess board and hourglass
Picnic basket and watermelon
Joint prompt, base model
Joint prompt, through the adapter
Adapter trained for 1,000 epochs
b)
One fused animal
Invented people
Grey mush, no animal in it
Flat line art of one animal
A third animal in the corner
Two animals of the wrong species

Figure 7: **Limitations of the Plurality LoRA:** a) The adapter can change the render under joint prompting and is inconsistently useful in this case. Each row compares the model conditioned on the joint prompt with and without the LoRA applied from the same noise sample. While the LoRA improves the render for a chessboard and hourglass, it disrupts the render for a cat and a dog, and a tiger and a dog. b) We show five examples of how a corrected cat–dog render can lead to inconsistent or unexpected results. The uncorrected hybrid (top left) is followed by five corrected failures showing intrusions (top middle and bottom right), counting errors (bottom middle), style-dependent loss of plurality (bottom left) or instability (top right).

56173_Restoring_plurality_in_c.pdf (submitted), page 9

failures, no count

counted per window

shown per window

THE REVISED ANSWER

Figure 10, failure types by window

steps 0 to 10, 0 to 25, all 50



Figure 10: Failure types as the adapter acts on fewer steps. Each image is labelled by Claude

paper/iclr-56173/build/iclr2027_conference.pdf (revised), page 16

Figure 12, the images behind it

the renders



Figure 12: The images behind Figure 10. Seven held-out pairs, each at the seed where

paper/iclr-56173/build/iclr2027_conference.pdf (revised), page 18

The reply we plan to send (replies/k4nd-and-uhz6.md: UHz6 Q3)

answered

Q3: restricting the adapter to the early steps

We tested this directly. Figure 10 counts each failure type on the 53 held-out pairs at seeds 9 to 16 with the adapter switched on for the first 10, the first 25 or all 50 of the 50 steps, and Figure 12 shows the images. The first 25 steps give the same result as all 50. With the first 10 only, the detector finds two animals far less often (Figure 11a). The cow and horse row of Figure 12 shows that the species substitution of the submitted Fig. 7 (Figure 14 in the revision) persists at every window, so it does not come from the late steps.

The submitted Figure 7 is Figure 14 in the revision. Every excerpt is a crop of a page render; the file and page are printed under it. Boxes mark text already on the page.