



Exercise Sheet 3

Machine Learning Basics

Deadline: 13.11.2024 23:59

Guidelines: You are expected to work in a group of 2-3 students. While submitting the assignments, please make sure to include the following information for all our teammates in your PDF/python script:

Name:

Student ID (matriculation number):

Email:

Your submissions should be zipped as **Name1_id1_Name2_id2_Name3_id3.zip** when you have multiple files. For assignments where you are submitting a single file, use the **same naming convention** without creating a zip. For any clarification, please reach out to us on the **CMS Forum**. These instructions are mandatory. If you are not following them, tutors can decide not to correct your exercise.

Please note:

- **Notational clarification:** In this assignment, x, y, z denote scalar values; $\mathbf{x}, \mathbf{y}, \mathbf{z}$ denote vectors; and X, Y, Z denote matrices.
- Ex 3.1-3.3 are written assignments, please submit a pdf (written using Latex) with the **names, matriculation IDs and emails** of all team members for this part. In case you are not familiar with Latex, clearly written handwritten submissions are also accepted, but we strongly encourage pdfs written using Latex.
- Ex 3.4 and 3.5 are programming assignments, you can write your code in the supplied notebooks and submit them. Don't forget to put in your **names, matriculation IDs and emails**.
- Submit the pdfs and notebooks together in a zip file in CMS. No need to submit any datasets.

Exercise 3.1 - Linear Regression

((1+0.25+0.25)+0.5+0.5 points)

Linear regression aims to model the relationship between the matrix of data points X and the label vector \mathbf{y} . Considering a dataset of N data points, with each having n features, this relationship is formulated using a linear equation of the form:

$$y^{(i)} = \mathbf{w}^T \mathbf{x}^{(i)} = b + w_1 x_1^{(i)} + \dots + w_n x_n^{(i)}; \quad (\text{for a single data point})$$

or,

$$\mathbf{y} = X\mathbf{w}; \quad (\text{in matrix notation for the entire dataset})$$

where \mathbf{w} represents the vector of regression parameters. Here, $X_{N \times (n+1)} = \begin{bmatrix} 1 & x_1^{(1)} & \cdots & x_n^{(1)} \\ 1 & x_1^{(2)} & \cdots & x_n^{(2)} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & x_1^{(N)} & \cdots & x_n^{(N)} \end{bmatrix} =$

$$\begin{bmatrix} \mathbf{x}_1^T \\ \vdots \\ \mathbf{x}_N^T \end{bmatrix}, \mathbf{y}_{N \times 1} = \begin{bmatrix} y^{(1)} \\ \vdots \\ y^{(N)} \end{bmatrix}, \text{ and } \mathbf{w}_{(n+1) \times 1} = \begin{bmatrix} b \\ w_1 \\ \vdots \\ w_n \end{bmatrix}.$$

Here, we include the bias term b as part of \mathbf{w} , so it has dimension $(n+1)$. The bias is also represented by the coefficient 1 in each data vector \mathbf{x}_i .

The optimal parameters that satisfy this equation given the data minimize the cost function:

$$f(\mathbf{w}) = \frac{1}{N} \|X\mathbf{w} - \mathbf{y}\|_2^2$$

- a) This least-squares formulation has a closed-form analytical solution called the normal equation. Derive the formulation of the normal equation, clearly stating any lemmas or assumptions used in your derivation.

Now, answer the following questions:

- i. What are the problems with using this normal equation method?
 - ii. Does this work when the data matrix X is not invertible? If yes, how so?
- b) Consider the following dataset \mathcal{D}_1 :

x	9.0	2.0	6.0	1.0	8.0
y	1.0	0.0	3.0	0.0	1.0

Using the normal equation derived above, find the optimal parameters \mathbf{w} given the dataset \mathcal{D}_1 .

- c) Let us assume we already have the parameters $\mathbf{w}_{\mathcal{D}_1}^T = [b_1 \ w_1]$ that minimize the mean-squared error (MSE) for \mathcal{D}_1 . Now, consider another dataset \mathcal{D}_2 :

x	$9.0+\gamma$	$2.0+\gamma$	$6.0+\gamma$	$1.0+\gamma$	$8.0+\gamma$
y	$1.0+\eta$	$0.0+\eta$	$3.0+\eta$	$0.0+\eta$	$1.0+\eta$

where $\gamma, \eta > 0$ and $w_1\gamma \neq \eta$. Let $\mathbf{w}_{\mathcal{D}_2}^T = [b_2 \ w_2]$ be the parameters that minimize the MSE for \mathcal{D}_2 . Which of the cases listed below hold in this case? Explain your reasoning.

- a) $w_1 = w_2, b_1 = b_2$
- b) $w_1 \neq w_2, b_1 = b_2$
- c) $w_1 = w_2, b_1 \neq b_2$
- d) $w_1 \neq w_2, b_1 \neq b_2$

Exercise 3.2 - Principal Component Analysis (PCA)-I

(0.5 + 0.5 points)

Please answer the following questions in two to three sentences:

- a) Why is normalization an important step in PCA?
- b) Give an example of an instance when PCA performs badly.

Exercise 3.3 - Principal Component Analysis (PCA)-II

(0.5 + 0.5 + 0.5 points)

Let's assume we've performed PCA on the toy dataset shown in Table 1:

Table 1: Dataset

Row	X1	X2	X3	X4
1	0.49	0.07	0.12	-1.19
2	-0.35	1.14	0.18	0.57
3	-0.44	0.29	-0.85	0.30
4	0.65	-0.42	-0.30	-0.22
5	1.15	-0.44	0.77	0.98
6	0.45	0.14	-0.02	0.86

And we've obtained the principal components as shown in Table 2:

Table 2: Principal Components of the dataset

PC1	PC2	PC3	PC4
0.69	-0.24	0.03	-0.7
-0.49	0.33	0.58	-0.55
-0.45	0.06	0.76	0.45
0.32	-0.9	-0.26	-0.02

Which correspond to the following eigenvalues:

[0.739, 0.685, 0.239, 0.004]

Answer the following questions:

- a) Why are there only 4 principal components?
- b) How much of the variance in the data is preserved by the first two principal components?
- c) How much of the variance in the data is preserved by the first and third principal components together?

Exercise 3.4 - Image denoising using PCA

(5 points)

See attached notebook

Exercise 3.5 - Polynomial Regression (Bonus)

(3 points)

See attached notebook