

Unit 2: Conditioning and Independence

MITx Statistics and Data Science

Seung Hyun Sung

2022 10 6

Note

Bayes' rule is the foundation for the field of inference. It guides on how to process data and make inferences about unobserved quantities or phenomena

[1] Conditioning and Bayes'Rule

Conditional Probabilities

- $P(A|B) = \frac{P(A \cap B)}{P(B)}$, defined only when $P(B) > 0$.
- $P(A \cap B|A) = \frac{P(A \cap B)}{P(A)}$
- Properties:
 - $P(A|B) \geq P(B)$, assuming $P(B) > 0$
 - If $A \cap B = \emptyset$, then $P(A \cup C|B) = P(A|B) + P(C|B)$

Three important tools

- Multiplication rule
$$P(A \cap B) = P(B)P(A|B) + P(A)P(B|A)$$
$$P(A^c \cap B \cap C^c) = P(A^c)P(B|A^c)P(C^c|A^c \cap B)$$
$$P(A_1 \cap A_2 \cap A_3) = P(A_1) \prod_{i=2}^n P(A_i|A_1 \cap \dots \cap A_{i-1})$$
- Total probability theorem
Mutually exclusive ($B_i \cap B_j = \emptyset$), $\{B_1, B_2, \dots, B_k\}$
$$P(B) = \sum_{i=1}^k P(A_i)P(B|A_i), \text{ which is a weighted average of } P(B|A). \text{ Note that } \sum_i P(A_i) = 1$$
- Bayes' rule
$$P(A_i|B) = \frac{A_i \cap B}{P(B)} = \frac{P(A_i)P(B|A_i)}{\sum_{j=1}^k P(A_j)P(B|A_j)} = \frac{P(A_i)P(B|A_i)}{P(B)}$$

Inference

- initial belief $P(A_i)$ on possible causes of an observed event B
$$A_i \xrightarrow{\text{model } P(B|A_i)} B$$
- draw conclusions about causes
$$B \xrightarrow{\text{inference } P(A_i|B)} A_i$$

[2] Independence

- Independence of two events
 - Definition of independence: $P(A \cap B) = P(A) \cdot P(B)$.
 - If $P(B|A) = P(B)$, occurrence of A provides no new information about B .
 - If A and B are independence, then A and B^c are independent.
- Conditional independence
 - Given C , the conditional independence is defined as independence under the probability law $P(\cdot|C)$.

$$P(A \cap B|C) = P(A|C)P(B|C)$$

- Independence does not imply conditional independence.

In the case below, A and B have no intersection in the condition of C . If A happens, B won't happen in the condition of C . This means that A and B are not independent.
- Independence of a collection of events
 - Event A_1, A_2, \dots, A_n are independent if $P(A_i \cap A_j \cap \dots \cap A_m) = P(A_i)P(A_j)\dots P(A_m)$, for any distinct indices i, j, \dots, m .
- Pairwise independence
 - Independent events must be pairwise independent, but the reverse may not be true.
 - E.g. two independent fair coin tosses. C : the two tosses had the same result. H_1 : first toss is H , H_2 : second toss is H , $P(H_1) = P(H_2) = 1/2$, $P(C) = 1/2$.
 - $P(H_1 \cap H_2) = 1/4 = P(H_1)P(H_2)$
 - $P(H_1 \cap C) = 1/4 = P(H_1)P(C)$
 - But, $P(C|H_1 \cap H_2) = 1 \neq P(C) = 1/2$

So H_1, H_2, C are pairwise independent, but not independent.
- Reliability
 - p_i : probability that unit i is “up” ; u_i : i th unit up, u_i are independent; F_i : i th unit down, F_i are independent.

$$\begin{aligned} P(\text{system up}) &= P(u_1 \cap u_2 \cap u_3) \\ &= P(u_1) \cap P(u_2) \cap P(u_3) \\ &= p_1 p_2 p_3 \end{aligned}$$

$$\begin{aligned} P(\text{system up}) &= P(u_1 \cup u_2 \cup u_3) \\ &= 1 - P(F_1 \cap F_2 \cap F_3) \\ &= 1 - P(F_1)P(F_2)P(F_3) \\ &= 1 - (1 - p_1)(1 - p_2)(1 - p_3) \end{aligned}$$

- In general, if a **serial** sub-system contains m components with success probabilities $p_1, p_2 \dots p_m$, then the probability of success of the entire sub-system is given by

$$P(\text{whole system succeeds}) = p_1 p_2 \dots p_m$$

If a parallel sub-system contains m components with success probabilities $p_1, p_2 \dots p_m$, then the probability of success of the entire sub-system is given by

$$P(\text{whole system succeeds}) = 1 - (1 - p_1)(1 - p_2) \dots (1 - p_m)$$