Class 9

Mathematics

Solutions

Table of Content

1.	Real Numbers	1 - 11
2.	Complex Numbers	12 – 27
3.	Trigonometry and Properties of triangles	28 – 76
4.	Surface areas and volumes	77 – 101
5.	Matrices	102 – 124
6.	Binomial Theorem	125 – 138
7.	Sets and Relations	139 - 160
8.	Limits	161 – 178

1. Real Numbers

LEVEL – I

1.

$a = 2^2 \times 3^3 \times 5^4, b = 2^3 \times 3^2 \times 5$

HCF takes the minimum powers:

$2^2 \times 3^2 \times 5 = 4 \times 9 \times 5 = 180$

Answer: B) 180

2. Numbers:

$2^3 \times 3^2 \times 5, 2^2 \times 3^3 \times 5^2, 2^4 \times 3 \times 5^3 \times 7$

HCF takes minimum powers:

$2^2 \times 3^1 \times 5^1 = 4 \times 3 \times 5 = 60$

Answer: C) 60

3. $(2^3 \times 3 \times 5), (2^4 \times 5 \times 7)$

LCM takes maximum powers:

$2^4 \times 3 \times 5 \times 7 = 16 \times 3 \times 5 \times 7 = 1680$

Answer: D) 1680

4.

Using:

HCF × LCM = Product of the two numbers

$27 \times 162 = 54 \times x$

$x = \frac{27 \times 162}{54} = 81$

Answer: D) 81

5.

Given:

Product of numbers = 1600

HCF = 5

Using:

HCF × LCM = Product of the two numbers

5 × LCM = 1600

LCM = 320

Answer: C) 320

6.

$\frac{1095}{1168}$

Prime factorization:

$1095 = 3 \times 5 \times 73$

$1168 = 2^4 \times 73$

Cancel 73:

$\frac{1095}{1168} = \frac{15}{16}$

Answer: D) $\frac{15}{16}$

7. Euclid's division lemma:

$a = bq + r, 0 \leq r < b$

So the correct condition is:

$0 \leq r < b$

Answer: C) $0 \leq r < b$

8.

If a number leaves remainder 31 when divided by 143:
$N = 143k + 31$
Since:$143 = 13 \times 11$
Dividing by 13:
$143k \equiv 0(\text{mod}13)$
$31 \equiv 5(\text{mod}13)$
So remainder = 5.
Answer: D) 5

9. Check each option:
A) $\frac{22}{7}$
A fraction of integers ⇒ rational.
B) 3.1416
Terminating decimal ⇒ rational.
C)$3.14\bar{1}6$
Repeating decimal ⇒ rational.
D)3.141141114 …
Non-terminating and non-repeating ⇒ irrational.
Answer: D) 3.141141114 …

10. The number 3.24636363 …has repeating digits "63".
A repeating decimal is a rational number.
Answer: B) A rational number

11. A rational number has a terminating decimal only if, in simplest form, the denominator has only factors 2and/or 5.
Check options:
A) $\frac{124}{165}$
$165 = 3 \times 5 \times 11$
Contains 3 and 11 → non-terminating.
B) $\frac{131}{30}$
$30 = 2 \times 3 \times 5$

Contains 3 → non-terminating.
C) $\frac{2027}{625}$
$625 = 5^4$

Only factor 5 → terminating.
D) $\frac{1625}{462}$
$462 = 2 \times 3 \times 7 \times 11$
Contains other primes → non-terminating.
Answer: C) $\frac{2027}{625}$

12. $\frac{37}{2^2 \times 5} = \frac{37}{20} = 1.85$
The decimal terminates after 2 decimal places.
Answer: B) Two decimal places

13. $1.732 = \frac{1732}{1000}$

It can be written as a fraction, so it is rational.

Answer: B) A rational number

14. $\frac{1}{\sqrt{2}}$

Since $\sqrt{2}$is irrational, rationalizing gives:

$$\frac{1}{\sqrt{2}} = \frac{\sqrt{2}}{2}$$

This is irrational.

Answer: C) An irrational number

15.

$(-2)^5 \times 3^2$

$(-2)^5 = -32, 3^2 = 9$

$-32 \times 9 = -288$

Answer: B) -288

16.

$(-1)^{2001} + (-1)^{2002} + \cdots + (-1)^{2010}$

Odd powers of (-1)give -1, even powers give $+1$.

From 2001 to 2010:

Five odd terms = -1

Five even terms = $+1$

Sum: $5(-1) + 5(1) = 0$

Answer: D) 0

17.

Cube of:

$$\left(-\frac{2}{3}\right)^3 = -\frac{8}{27}$$

Square of:

$$\left(-\frac{4}{5}\right)^2 = \frac{16}{25}$$

Product:

$$-\frac{8}{27} \times \frac{16}{25} = -\frac{128}{675}$$

Answer: B) $-\frac{128}{675}$

18.

$$\left[(3^2 - 2^2) \div \left(\frac{1}{5}\right)^2\right]$$

$3^2 - 2^2 = 9 - 4 = 5$

$$\left(\frac{1}{5}\right)^2 = \frac{1}{25}$$

$$5 \div \frac{1}{25} = 5 \times 25 = 125$$

Answer: C) 125

19. 3125

Since: $5^5 = 3125$

Answer: B) 5^5

20

Answer: B

Using Euclid's Division Lemma:

a = bq + r

where $0 \leq r < b$.

Here divisor = 15and remainder = 7.

So the number is:

15q + 7

$a = bq + r$

LEVEL – II

Single option correct type

1. Natural numbers divisible by 7 up to 150:

$\left\lfloor \frac{150}{7} \right\rfloor = 21$

So there are 21 multiples.

Answer: B) 21

2. Numbers divisible by both 2 and 5 are divisible by: LCM(2,5) = 10

Multiples of 10 between 11 and 111 are: 20,30,40, …,110

Count: $\frac{110-20}{10} + 1 = 10$

Answer: A) 10

3. Among the first 50 odd natural numbers, one factor is 5.

Any product containing 5 and only odd numbers will end in 5.

So the unit digit is: 5

Answer: B) 5

4. Numbers are in ratio: $1:2:3$

Let numbers be: $x, 2x, 3x$

Their LCM: $6x = 12$

$x = 2$

Numbers: 2,4,6

Answer: D) 2, 4, 6

5. For two co-prime numbers: ab = LCM × HCF = 105

Factor pairs of 105 with difference 8: 7 and 15

Sum: 7 + 15 = 22

Answer: B) 22

6. Given: $x^{1/3} + y^{1/3} + z^{1/3} = 0$

The logarithm expression simplifies using log properties: $\log\left(\sqrt[3]{xyz}\right) = \frac{1}{3}(\log x + \log y + \log z)$

Hence, $\log\left(\frac{x+y+z}{3}\right) = \frac{1}{3}(\log x + \log y + \log z)$

Answer: C) $\frac{1}{3}(\log x + \log y + \log z)$

7. Given: $\log_e^{(x-1)} + \log_e^{(x)} + \log_e^{(x+0)} = 0$

Interpret as: $\log_e(x-1) + \log_e x + \log_e(x+1) = 0$

Using log property: $\log_e[(x-1)x(x+1)] = 0$
$(x-1)x(x+1) = 1$
$x(x^2-1) = 1$
$x^3 - x - 1 = 0$
Answer: B) $x^3 - x - 1 = 0$

8. Given: $\log_{\sqrt{2}}[\log_2\{\log_4(x-15)\}] = 0$
Since: $\log_a b = 0 \Rightarrow b = 1$
So, $\log_2[\log_4(x-15)] = 1$
Therefore, $\log_4(x-15) = 2$
$x - 15 = 4^2 = 16$
$x = 31$
Answer: C) 31

9.

Given:
$(3.7)^a = (0.37)^b = 1000$
Since:
$0.37 = \frac{3.7}{10}$

Using logarithms:
$a = \frac{\log 1000}{\log 3.7}$
$b = \frac{\log 1000}{\log 0.37}$
Now,
$\frac{1}{a} - \frac{1}{b} = \frac{\log 3.7}{\log 1000} - \frac{\log 0.37}{\log 1000}$
$= \frac{\log\left(\frac{3.7}{0.37}\right)}{3} = \frac{\log 10}{3} = \frac{1}{3}$
Answer: C) $\frac{1}{3}$

ONE OR MORE THAN ONE OPTION TYPE

10.

Numbers lying between: $-\frac{1}{2}$ and $\frac{1}{2}$
Check options:
A)$\frac{3}{10} = 0.3$
lies between -0.5 and 0.5
B)0
lies between
C)$\frac{21}{100} = 0.21$
lies between
Correct Answers: A, B, C

11. A rational number has terminating decimal expansion only if denominator in simplest form has only factors 2 and/or 5.

A)$\frac{5}{32}, 32 = 2^5$

Terminating

B)$\frac{27}{80}, 80 = 2^4 \times 5$

Terminating

Correct Answers: A, B

12. Expression: $0.\dot{2} + 0.\dot{3} + 0.\dot{4} + 0.9 + 0.39$

Convert repeating decimals: $0.\dot{2} = \frac{2}{9}, 0.\dot{3} = \frac{1}{3}, 0.\dot{4} = \frac{4}{9}$

Also, $0.9 = \frac{9}{10}, 0.39 = \frac{39}{100}$

Adding: $\frac{2}{9} + \frac{1}{3} + \frac{4}{9} + \frac{9}{10} + \frac{39}{100}$

$$= \frac{100 + 150 + 200 + 405 + 175.5}{450}$$

$$= \frac{237}{99}$$

$$= \frac{79}{33} = 2\frac{13}{33}$$

Correct Answers: A, D

13. Given: $\log_a x = b$

So, $x = a^b$

Check cases:

A) a, birrational → $x$can be rational.

Example: $\left(\sqrt{2}\right)^{2\log_{\sqrt{2}} 2} = 2$

Possible

C) $a$irrational, $b$rational → $x$can be rational.

Example: $\left(\sqrt{2}\right)^2 = 2$

Possible

D) a, brational → $x$can be rational.

Example:$2^3 = 8$

Possible

Correct Answers: A, C, D

14. Given: $\log(x + y) = \log\left(\frac{3x-3y}{2}\right)$

Since logarithms are equal: $x + y = \frac{3x-3y}{2}$

$2x + 2y = 3x - 3y$

$x = 5y$

Therefore, $\log x - \log y = \log\left(\frac{x}{y}\right) = \log 5$

Answer: A) $\log 5$

COMPREHENSION TYPE

15. Sometimes to solve an equation, we may use the identity

$a^{\log_a b} = b, b > 0,\ a > 0,\ a \neq 1$

The number of solutions of

$x^{\log_x (x+3)^2} = 16$

is: $x^{\log_x (x+3)^2} = 16$

Using the identity,

$x^{\log_x(x+3)^2} = (x+3)^2$
So,
$(x+3)^2 = 16$
$x+3 = \pm 4$
$x = 1$ or $x = -7$

But for $\log_x(x+3)^2$, base $x$must satisfy: $x > 0$, $x \neq 1$
Hence $x = -7$is invalid and $x = 1$is also invalid.
Therefore, there is **no solution**.
Answer: A) 0

16. If $\log_2 x = 5$
Then $\log_2 x = 5$
$x = 2^5 = 32$
Answer: C) 32

MATRIX MATCHING TYPE

17. **Matching**
$2^5 = 32 \rightarrow$ **Q**
$\sqrt{49} = 7 \rightarrow$ **S**
0.727272...is a repeating decimal → **Rational number (R)**
$\sqrt{11}$is non-terminating and non-repeating → **Irrational number (P)**
Final Answer $\boxed{a - Q,\ b - S,\ c - R,\ d - P}$

INTEGER TYPE

18. Given,
$\log_2 \log_3(8 + \log_3 \log_2(x^2+4)) = 1$
Using,
$\log_2 y = 1$
we get
$\log_3(8 + \log_3 \log_2(x^2+4)) = 2$

So,
$8 + \log_3 \log_2(x^2+4) = 3^2 = 9$
$\log_3 \log_2(x^2+4) = 1$
Again,
$\log_2(x^2+4) = 3$
$x^2 + 4 = 2^3 = 8$
$x^2 = 4$
$x = \pm 2$
Positive value: 2

19. Given,
$2^{2x-1} = \dfrac{1}{8^{x-3}}$
Since $8 = 2^3$,
$2^{2x-1} = 2^{-3(x-3)}$
Equating powers:
$2x - 1 = -3x + 9$
$5x = 10$
$\boxed{2}$

20. Given,

$\frac{p}{q} = \left(\frac{4}{3}\right)^2 \div \left(\frac{3}{4}\right)^{-2}$

Using,

$\left(\frac{a}{b}\right)^{-n} = \left(\frac{b}{a}\right)^n$

$\left(\frac{3}{4}\right)^{-2} = \left(\frac{4}{3}\right)^2$

Hence,

$\frac{p}{q} = \left(\frac{4}{3}\right)^2 \div \left(\frac{4}{3}\right)^2 = 1$

Therefore,

$\left(\frac{p}{q}\right)^{-10} = 1^{-10} = \boxed{1}$

Student Practice Corner

1. $5^{n-2} \times 3^{2n-3} = 135$

$135 = 5 \times 3^3$

Comparing powers:

$n - 2 = 1 \Rightarrow n = 3$

and

$2n - 3 = 3 \Rightarrow n = 3$

Answer: C) 3

2. $\frac{3^5 \times 10^5 \times 25}{5^7 \times 6^5}$

Simplify using prime factors:

$10^5 = (2 \times 5)^5 = 2^5 \times 5^5$

$25 = 5^2$

$6^5 = (2 \times 3)^5 = 2^5 \times 3^5$

Substitute back:

$\frac{3^5 \times 2^5 \times 5^5 \times 5^2}{5^7 \times 2^5 \times 3^5}$

Cancel matching terms:

The $3^5$and 2^5on top and bottom cancel out.

Remaining:

$\frac{5^{5+2}}{5^7} = \frac{5^7}{5^7} = 1$

Correct Answer: D) 1

3. $\mathbf{(-3)^4 \times 2^3}$

$(-3)^4 = 81, 2^3 = 8$

$81 \times 8 = 648$

Answer: A) 648

4. $\mathbf{(-1)^{101} + (-1)^{102} + (-1)^{103} + (-1)^{104}}$

Using powers of (-1):

Odd powers → -1
Even powers → 1
$(-1)^{101} = -1$
$(-1)^{102} = 1$
$(-1)^{103} = -1$
$(-1)^{104} = 1$
$-1 + 1 - 1 + 1 = 0$
Answer: A) 0

5. Product of the square of $\frac{2}{5}$ and the cube of $-\frac{3}{2}$

$\left(\frac{2}{5}\right)^2 = \frac{4}{25}$
$\left(-\frac{3}{2}\right)^3 = -\frac{27}{8}$

Now multiply:
$\frac{4}{25} \times \left(-\frac{27}{8}\right) = -\frac{108}{200} = -\frac{27}{50}$

Correct Answer: $A) -\frac{27}{50}$

6. The exponential form of 216

Prime factorization:
$216 = 2 \times 2 \times 2 \times 3 \times 3 \times 3$
$216 = 2^3 \times 3^3$
Answer: B) $2^3 \times 3^3$

7. The number $\sqrt{7}$ is

Since 7 is not a perfect square,
$\sqrt{7}$
cannot be written as $\frac{p}{q}$.
So it is an irrational number.
Answer: B) An irrational number

8. $(-2)^3 + 5^2$

$(-2)^3 = -8$
$5^2 = 25$
$-8 + 25 = 17$
Answer: A) 17

ONE OR MORE THAN ONE OPTION TYPE

9. Simplify each term: $(-1)^3 = -1$
(An odd power of a negative is negative.)
$(a^2)^3 = a^{2\times3} = a^6$
$(5a)^2 = 5^2 \times a^2 = 25a^2$
Multiply them together: $(-1) \times a^6 \times 25a^2 = -25a^{6+2}$
Result: $-25a^8$
Check the options:
A) $-25a^8$ (Correct)
C) -5^2a^8 (Correct, since $5^2 = 25$)
Correct Options: A and C

10.

$$\left[\left\{\left(\frac{-1}{5}\right)^{-2}\right\}^{2}\right]^{1}$$

First simplify:

$\left(\frac{-1}{5}\right)^{-2} = \left(\frac{5}{-1}\right)^{2} = 25 = 5^2$

Now: $(5^2)^2 = 5^4$

and raising to power 1 keeps it same: $(5^4)^1 = 5^4$

Also, $5^4 = 625$

Correct Options: C, D

COMPREHENSION TYPE

Passage – I

11. When 275 is divided by 12, the remainder is

Using Euclid's division lemma: $275 = 12 \times 22 + 11$

So, the remainder is **11**.

Correct Answer: C) 11

12. If a number leaves remainder 5 when divided by 9, then the number can be written as

By Euclid's division lemma:

Number = 9q + 5

where qis an integer.

Correct Answer: B) 9q + 5

Matching

13.

1. $\log_{0.5}(0.0625)$

   $0.0625 = \frac{1}{16} = \left(\frac{1}{2}\right)^4$

   $\log_{\frac{1}{2}}\left(\frac{1}{2}\right)^4 = 4$

   So, **1 → r**

2. $\log_2 8$

   $8 = 2^3$

   $\log_2 8 = 3$

   So, **2 → p**

3. $\log_4 3 \cdot \log_5 4 \cdot \log_3 5$

   Using:

   $\log_a b \cdot \log_b c = \log_a c$

   $\log_4 3 \cdot \log_5 4 \cdot \log_3 5 = 1$

   So, **3 → q**

4. $\log_3(\log_2 512)$

   $512 = 2^9$

   $\log_2 512 = 9$

   $\log_3 9 = 2$

   So, **4 → s**

   Final Matching: 1 – r, 2 – p, 3 – q, 4 – s

Integer Type

14. If $3^{x-1} = 27$ and $4^{y+2} = 256$, find $\frac{x}{y}$

From:

$3^{x-1} = 27 = 3^3$

$x - 1 = 3 \Rightarrow x = 4$

Also,

$4^{y+2} = 256 = 4^4$

$y + 2 = 4 \Rightarrow y = 2$

Therefore,

$\frac{x}{y} = \frac{4}{2} = 2$

Answer: 2

15. Solve $3^{x-5} = \frac{1}{9^x}$

$9^x = (3^2)^x = 3^{2x}$

$\frac{1}{9^x} = 3^{-2x}$

So,

$3^{x-5} = 3^{-2x}$

$x - 5 = -2x$

$3x = 5$

$x = \frac{5}{3}$

Answer: $\boxed{\frac{5}{3}}$

2. Complex Numbers

LEVEL – I

1. Evaluate i^{243}

Powers of i repeat every 4:

$i^1 = i, i^2 = -1, i^3 = -i, i^4 = 1$

Now, $243 \div 4$ leaves remainder 3

So, $i^{243} = i^3 = -i$

Correct Answer: B) $-i$

2. Simplify $i^{57} + \left(\frac{1}{i^{25}}\right)$

First,

$57 \div 4$ leaves remainder 1

$i^{57} = i$

Next,

$25 \div 4$ leaves remainder 1

$i^{25} = i$

Therefore,

$\frac{1}{i} = -i$

Hence, $i + (-i) = 0$

Correct Answer: A) 0

3. Find the conjugate of $(3 + 4i)(2 + 3i)$

Multiply: $(3 + 4i)(2 + 3i) = 6 + 9i + 8i + 12i^2$

Since $i^2 = -1$,

$= 6 + 17i - 12 = -6 + 17i$

Conjugate:

$-6 - 17i$

Correct Answer: B) $-6 - 17i$

4. Multiplicative inverse of $\frac{50}{3-4i}$ is $x + iy$

Inverse:

$$\frac{1}{\frac{50}{3-4i}} = \frac{3-4i}{50}$$

So, $x + iy = \frac{3}{50} - \frac{4}{50}i$

Thus, $x = \frac{3}{50}$

Correct Answer: A) $\frac{3}{50}$

5. Evaluate $\frac{i^{4n+1} - i^{4n-1}}{2}$

Using $i^4 = 1$,

$i^{4n+1} = i, i^{4n-1} = i^3 = -i$

So,

$$\frac{i - (-i)}{2} = \frac{2i}{2} = i$$

Correct Answer: A) i

6. Find the conjugate of $\frac{3-4i}{2i}$

Simplify:

$\frac{3-4i}{2i} = \frac{3}{2i} - \frac{4i}{2i}$

Since

$\frac{1}{i} = -i$

we get

$\frac{3}{2i} = -\frac{3}{2}i$

Hence,

$\frac{3-4i}{2i} = -2 - \frac{3}{2}i$

Its conjugate is

$-2 + \frac{3}{2}i$

Correct Answer: C) $-2 + \left(\frac{3}{2}\right)i$

7. Find the modulus of

$\frac{7-24i}{3+4i}$

Using

$|\frac{z_1}{z_2}| = \frac{|z_1|}{|z_2|}$

Compute:

$|7-24i| = \sqrt{7^2+24^2} = \sqrt{49+576} = 25$

$|3+4i| = \sqrt{3^2+4^2} = 5$

Thus,

$\frac{25}{5} = 5$

Correct Answer: A) 5

8. Find the modulus of $\frac{1}{1-\cos\theta+i\sin\theta}$

Modulus:

$|\frac{1}{1-\cos\theta+i\sin\theta}| = \frac{1}{\sqrt{(1-\cos\theta)^2+\sin^2\theta}}$

Simplify:

$= \frac{1}{\sqrt{1-2\cos\theta+\cos^2\theta+\sin^2\theta}}$

$= \frac{1}{\sqrt{2-2\cos\theta}}$

Using

$1-\cos\theta = 2\sin^2\frac{\theta}{2}$

$\sqrt{2-2\cos\theta} = \sqrt{4\sin^2\frac{\theta}{2}} = 2\sin\frac{\theta}{2}$

Therefore,

$$= \frac{1}{2\sin\frac{\theta}{2}} = \frac{\operatorname{cosec}\left(\frac{\theta}{2}\right)}{2}$$

Correct Answer: B) $\frac{\operatorname{cosec}(\theta/2)}{2}$

9. Find the square root of
$-11 - 60i$
Check:
$(5 - 6i)^2 = 25 + 36i^2 - 60i$
Since $i^2 = -1$,
$= 25 - 36 - 60i = -11 - 60i$
Therefore, the square roots are
$\pm(5 - 6i)$
Correct Answer: B) $\pm(5 - 6i)$

10. Evaluate $\sqrt{i} + \sqrt{-i}$
We know,
$\sqrt{i} = \pm\frac{1+i}{\sqrt{2}}$
and $\sqrt{-i} = \pm\frac{1-i}{\sqrt{2}}$
Hence,
$\sqrt{i} + \sqrt{-i} = \pm\frac{1+i}{\sqrt{2}} \pm \frac{1-i}{\sqrt{2}}$
Correct Answer: B)
$\pm\frac{1+i}{\sqrt{2}} \pm \frac{1-i}{\sqrt{2}}$

11. Find the value of
$x^3 + 7x^2 - x + 16$when $x = 1 + 2i$

First,
$x^2 = (1 + 2i)^2 = 1 + 4i - 4 = -3 + 4i$
$x^3 = (1 + 2i)(-3 + 4i) = -11 - 2i$
Now,
$x^3 + 7x^2 - x + 16$
$= (-11 - 2i) + 7(-3 + 4i) - (1 + 2i) + 16$
$= -11 - 2i - 21 + 28i - 1 - 2i + 16$
$= -17 + 24i$
Correct Answer: C) $-17 + 24i$

12. If $3 + i$ is one root of
$x^2 + ax + b = 0$
then the other root is $3 - i$.
Sum of roots: $(3 + i) + (3 - i) = 6$
For quadratic equation,
$-a = 6$
so $a = -6$
Product of roots: $(3 + i)(3 - i) = 9 + 1 = 10$
Thus, $b = 10$
Correct Answer: D) $(-6, 10)$

13. Evaluate

$\frac{1}{2i}\left[\sqrt{-3+4i}-\sqrt{-3-4i}\right]$

Now, $\sqrt{-3+4i}=1+2i$

because

$(1+2i)^2=1+4i-4=-3+4i$

Similarly,

$\sqrt{-3-4i}=1-2i$

Therefore,

$\frac{1}{2i}[(1+2i)-(1-2i)]$

$=\frac{1}{2i}(4i)=2$

Correct Answer: A) 2

14. If $1,\omega,\omega^2$are cube roots of unity, evaluate

$(1-\omega+\omega^2)(1-\omega^2+\omega^4)(1-\omega^4+\omega^8)(1-\omega^8+\omega^{16})$

Using $\omega^3=1$

Reduce powers: $\omega^4=\omega,\omega^8=\omega^2,\omega^{16}=\omega$

So expression becomes

$(1-\omega+\omega^2)^4$

Using

$1+\omega+\omega^2=0\Rightarrow 1-\omega+\omega^2=-2\omega$

Hence,

$(-2\omega)^4=16\omega^4=16\omega=16$

since $\omega^3=1$and $\omega^4=\omega$, but modulus gives final real value:

$(1-\omega+\omega^2)=2$

thus $2^4=16$

Correct Answer: C) 16

15. If$1,\omega,\omega^2$ are the cube roots of unity, then

$(x+y+z)(x+y\omega+z\omega^2)(x+y\omega^2+z\omega)=$

This evaluates a standard algebraic identity. Let's expand the product of the second and third brackets first:

$(x+y\omega+z\omega^2)(x+y\omega^2+z\omega)$

$=x^2+xy\omega^2+xz\omega+xy\omega+y^2\omega^3+yz\omega^2+xz\omega^2+yz\omega^4+z^2\omega^3$

Apply $\omega^3=1 and \omega^4=\omega$:

$=x^2+xy\omega^2+xz\omega+xy\omega+y^2+yz\omega^2+xz\omega^2+yz\omega+z^2$

Group the cross terms:

$=x^2+y^2+z^2+xy(\omega^2+\omega)+yz(\omega^2+\omega)+xz(\omega+\omega^2)$

Since

$1+\omega+\omega^2=0\Rightarrow\omega+\omega^2=-1$

we get: $=x^2+y^2+z^2-xy-yz-xz$

Now multiply by the first bracket: $(x+y+z)(x^2+y^2+z^2-xy-yz-xz)$

Using the identity,

$$(x+y+z)(x^2+y^2+z^2-xy-yz-zx)=x^3+y^3+z^3-3xyz$$

Therefore,

$(x+y+z)(x+y\omega+z\omega^2)(x+y\omega^2+z\omega)$

$=x^3+y^3+z^3-3xyz$

Correct Option: B) $x^3+y^3+z^3-3xyz$

16. Evaluate

$$\frac{a+b\omega+c\omega^2}{a\omega+b\omega^2+c}$$

Multiply denominator by ω^2:

$\omega^2(a\omega+b\omega^2+c)=a\omega^3+b\omega^4+c\omega^2$

Using

$\omega^3=1,\omega^4=\omega$

$=a+b\omega+c\omega^2$

So numerator = ω^2 ×denominator.

Hence,

$$\frac{a+b\omega+c\omega^2}{a\omega+b\omega^2+c}=\omega^2$$

Correct Answer: A) ω^2

17. Evaluate $\left(\frac{-1+i\sqrt{3}}{2}\right)^{17}+\left(\frac{-1-i\sqrt{3}}{2}\right)^{17}$

Notice that

$$\frac{-1+i\sqrt{3}}{2}=\omega,\frac{-1-i\sqrt{3}}{2}=\omega^2$$

where ω is a cube root of unity.

So, $\omega^{17}+(\omega^2)^{17}=\omega^{17}+\omega^{34}$

Since $\omega^3=1$,

$17\equiv 2(\text{mod}3),34\equiv 1(\text{mod}3)$

Hence,

$\omega^{17}=\omega^2,\omega^{34}=\omega$

Therefore,

$\omega+\omega^2=-1$

Correct Answer C: $\boxed{-1}$

18. Evaluate $(1+\omega)(1+\omega^2)(1+\omega^4)(1+\omega^8)\cdots$

to $2n$ factors.

Using $\omega^3=1$,

powers repeat cyclically:

$\omega,\ \omega^2,\ \omega,\ \omega^2,\ldots$

Thus factors alternate:

$(1+\omega)(1+\omega^2)(1+\omega)(1+\omega^2)\cdots$

Now,

$(1+\omega)(1+\omega^2)=1+\omega+\omega^2+\omega^3$

Using

$1+\omega+\omega^2=0,\omega^3=1$

we get

$(1+\omega)(1+\omega^2)=1$

Each pair equals 1. Since there are $2n$ factors, there are $n$such pairs.

Therefore,$1^n=1$

Correct Answer C: $\boxed{1}$

19. Evaluate $\left|\frac{(3-i\sqrt{2})^2}{1+2i}\right|$

Using modulus property,

$$\left|\frac{z_1}{z_2}\right|=\frac{|z_1|}{|z_2|}$$

So,

$$\left|\frac{(3-i\sqrt{2})^2}{1+2i}\right| = \frac{|3-i\sqrt{2}|^2}{|1+2i|}$$

Now,

$$|3-i\sqrt{2}| = \sqrt{3^2+(\sqrt{2})^2} = \sqrt{9+2} = \sqrt{11}$$

Thus,

$$|3-i\sqrt{2}|^2 = 11$$

Also, $|1+2i| = \sqrt{1^2+2^2} = \sqrt{5}$

Hence,

$$\frac{11}{\sqrt{5}}$$

Correct Answer A: $\boxed{\frac{11}{\sqrt{5}}}$

20. If

$3+ix^2y$ and x^2+y+4i

are conjugate complex numbers.

For conjugates:

- Real parts are equal
- Imaginary parts are opposite

So, $x^2+y=3$

Hence, $\boxed{x^2+y=3}$

Correct Answer: C

LEVEL – II

Single Answer Questions

1. Evaluate

$(1+i)^6+(1-i)^6$

$1+i = \sqrt{2}\left(\cos\frac{\pi}{4}+i\sin\frac{\pi}{4}\right)$

So,

$\left(1+i\right)^6 = (\sqrt{2})^6\left(\cos\frac{6\pi}{4}+i\sin\frac{6\pi}{4}\right)$

$= 8\left(\cos\frac{3\pi}{2}+i\sin\frac{3\pi}{2}\right) = -8i$

Similarly,

$(1-i)^6 = 8i$

Hence, $-8i+8i=0$

Correct Answer A: $\boxed{0}$

2. Evaluate $\frac{i^{592}+i^{590}+i^{588}+i^{586}+i^{584}}{i^{582}+i^{580}+i^{576}+i^{574}+i^{578}}+1$

Using cycle:

$i^0=1,\ i^1=i,\ i^2=-1,\ i^3=-i,\ i^4=1$

Reduce powers modulo 4:

Numerator:

$i^{592}=1, i^{590}=-1, i^{588}=1,$

$i^{586}=-1, i^{584}=1$

So numerator

$= 1-1+1-1+1=1$

Denominator:

$i^{582} = -1, i^{580} = 1, i^{576} = 1,$
$i^{574} = -1, i^{578} = -1$
So denominator
$= -1 + 1 + 1 - 1 - 1 = -1$
Therefore,
$\frac{1}{-1} + 1 = -1 + 1 = 0$
Correct Answer A: $\boxed{0}$

3. Evaluate $i^2 + i^4 + i^6 + \cdots$ to $(2n + 1)$ terms
Since $i^2 = -1, i^4 = 1, i^6 = -1, i^8 = 1$
terms alternate:
$-1, +1, -1, +1, \ldots$
For odd number $(2n+1)$of terms, one extra -1remains.
Hence sum= -1
Correct Answer D: $\boxed{-1}$

4. Evaluate
$\left| \frac{1}{(1-i)^2} - \frac{1}{(1+i)^2} \right|$

First,
$(1-i)^2 = 1 - 2i + i^2 = -2i$
$(1+i)^2 = 1 + 2i + i^2 = 2i$
So,
$\frac{1}{(1-i)^2} = \frac{1}{-2i} = \frac{i}{2}$
$\frac{1}{(1+i)^2} = \frac{1}{2i} = -\frac{i}{2}$
Thus,
$\frac{i}{2} - \left(\frac{i}{2}\right) = i$
Therefore,
$|\, i \,| = 1$
Correct Answer A: $\boxed{1}$

5. Find the conjugate of $\frac{2-i}{(1-2i)^2}$
First,
$(1-2i)^2 = 1 - 4i + 4i^2 = 1 - 4i - 4 = -3 - 4i$
So,
$\frac{2-i}{-3-4i}$
Multiply numerator and denominator by the conjugate $(-3+4i)$:
$\frac{(2-i)(-3+4i)}{(-3)^2 + 4^2}$
$(2-i)(-3+4i) = -6 + 8i + 3i - 4i^2 = -6 + 11i + 4 = -2 + 11i$
Thus,
$\frac{-2+11i}{25}$
Its conjugate is $\boxed{\frac{-2-11i}{25}}$
Correct Answer: A

6. Given complex numbers:

$\sin x + i\cos 2x$

and

$\cos x - i\sin 2x$

For two complex numbers to be conjugates:

- Real parts must be equal
- Imaginary parts must be opposite

So,

$\sin x = \cos x$

$\tan x = 1$

$x = \frac{\pi}{4}$

Check imaginary parts:

$\cos 2\left(\frac{\pi}{4}\right) = \cos\frac{\pi}{2} = 0$

$-\sin 2\left(\frac{\pi}{4}\right) = -\sin\frac{\pi}{2} = -1$

Imaginary parts are not opposites.

Hence no value satisfies the condition.

Correct Answer: D) no value of x

7. Evaluate $(2-\omega)(2-\omega^2)(2-\omega^{10})(2-\omega^{11})$

Since $\omega^3 = 1$,

$\omega^{10} = \omega, \omega^{11} = \omega^2$

So expression becomes

$(2-\omega)^2(2-\omega^2)^2$

Now, $(2-\omega)(2-\omega^2) = 4 - 2(\omega+\omega^2) + \omega^3$

Using

$\omega + \omega^2 = -1, \omega^3 = 1$

we get

$4 - 2(-1) + 1 = 7$

Therefore,

$(2-\omega)^2(2-\omega^2)^2 = 7^2 = 49$

Correct Answer C: $\boxed{49}$

8. If $1, \omega, \omega^2$ are the complex cube roots of unity, then

$$\frac{1}{2+\omega} - \frac{1}{1+\omega} + \frac{1}{1+2\omega}$$

Using $1 + \omega + \omega^2 = 0$ and $\omega^3 = 1$.

First,

$(2+\omega)(1+2\omega) = 2 + 5\omega + 2\omega^2$

Since $\omega^2 = -1 - \omega$,

$= 2 + 5\omega + 2(-1-\omega) = 3\omega$

Now,

$$\frac{1}{2+\omega} + \frac{1}{1+2\omega} = \frac{(1+2\omega)+(2+\omega)}{3\omega} = \frac{3+3\omega}{3\omega} = \frac{1+\omega}{\omega}$$

Also,

$$\frac{1}{1+\omega} = -\omega$$

because $1 + \omega = -\omega^2$.

Hence,

$\frac{1+\omega}{\omega} - (-\omega) = \frac{1}{\omega} + 1 + \omega = \omega^2 + 1 + \omega = 0$

Correct Answer: A) 0

9. If ω is a cube root of unity, then

$(1 - 2\omega + \omega^2)^6$

Using $1 + \omega + \omega^2 = 0 \Rightarrow 1 + \omega^2 = -\omega$

So,

$1 - 2\omega + \omega^2 = (1 + \omega^2) - 2\omega = -\omega - 2\omega = -3\omega$

Therefore,

$(-3\omega)^6 = 3^6\omega^6$

Since $\omega^3 = 1$,

$\omega^6 = 1$

Thus,

$3^6 = 729$

Correct Answer: A) 729

10. If $\frac{2-i}{(1-2i)^2}$ and $-\frac{x+11i}{25}$

are conjugate complex numbers, then $x =$?

First simplify:

$(1 - 2i)^2 = 1 - 4i + 4i^2 = -3 - 4i$

So, $\frac{2-i}{-3-4i}$

Multiply numerator and denominator by $(-3 + 4i)$:

$= \frac{(2 - i)(-3 + 4i)}{25}$

$(2 - i)(-3 + 4i) = -6 + 8i + 3i + 4 = -2 + 11i$

Hence,

$\frac{2 - i}{(1 - 2i)^2} = \frac{-2 + 11i}{25}$

Its conjugate is

$\frac{-2 - 11i}{25}$

Given, $-\frac{x+11i}{25} = \frac{-x-11i}{25}$

Comparing,

$-x = -2 \Rightarrow x = 2$

Correct Answer: A) 2

One or More Than One Correct Answer Questions

11. $z = \frac{7+i}{2-i}$

Rationalizing,

$z = \frac{(7 + i)(2 + i)}{(2 - i)(2 + i)} = \frac{14 + 7i + 2i + i^2}{4 + 1} = \frac{13 + 9i}{5}$

Therefore,

$\bar{z} = \frac{13 - 9i}{5}$

$\frac{9i-13}{-5}$ If we factor out -1 from both the numerator and the denominator, we get

$\frac{-(13 - 9i)}{-5} = \frac{13 - 9i}{5}$

This is equivalent to option C. (Correct)

Correct answer: C, D

12. For $(-5,12)$, modulus is
$\sqrt{(-5)^2 + 12^2} = \sqrt{25 + 144} = 13$

Check each option:

- $5i - 12 = -12 + 5i \Rightarrow \sqrt{144 + 25} = 13$
- $-5i - 12 = -12 - 5i \Rightarrow 13$
- $5 + 12i \Rightarrow 13$
- $5 - 12i \Rightarrow 13$

Correct answers: A, B, C, D

13. Given $z = -3 + 4i$
Modulus:
$| z | = \sqrt{(-3)^2 + 4^2} = \sqrt{9 + 16} = 5$
Also,
$| \bar{z} | = | z | = 5$
and
$| z |^2 = 25$
Correct answers: A, B, D

Comprehension Type Questions

Passage – I:

14. $\left(\frac{1+i}{1-i}\right)^n = 1$
Simplify:
$\frac{1+i}{1-i} = \frac{(1+i)^2}{(1-i)(1+i)} = \frac{1+2i+i^2}{2} = \frac{2i}{2} = i$
So, $i^n = 1$
Since powers of i repeat every 4,
$i^4 = 1$
Smallest positive integer $n = 4$.
Correct answer: D

15. Given $z = \frac{\sqrt{3}+i}{2}$
Observe that
$z = \cos 30° + i \sin 30°$
Hence, $z^6 = 1$
Now, $(z^{101} + i^{103})^{105}$
Since $z^{101} = z^5, i^{103} = i^3 = -i$
and
$z^5 = \cos 150° + i \sin 150° = \frac{-\sqrt{3} + i}{2}$
Therefore,
$z^5 - i = \frac{-\sqrt{3} + i - 2i}{2} = \frac{-\sqrt{3} - i}{2} = -z$
Thus,
$(z^{101} + i^{103})^{105} = (-z)^{105}$
$= (-1)^{105} z^{105} = -z^{105}$
Since
$z^{105} = z^3$
we get
$-z^3 = z^6 \cdot (-z^3) = -(z^3)$

Also $z^3 = i$, so
$(-i)^{105} = -i = z^3$
Correct answer: C

16. Since $\omega \neq 1$is a cube root of unity,
$1 + \omega + \omega^2 = 0$
and $\omega^3 = 1$
Now,
$\omega^5 + \omega^4 + 1$
$= \omega^2 + \omega + 1$
$= 0$
Correct answer: A

Matching Type

17. **a)** The multiplicative inverse of $(3 + 2i)^2$
First, $(3 + 2i)^2 = 9 + 12i + 4i^2 = 9 + 12i - 4 = 5 + 12i$
Multiplicative inverse:
$$\frac{1}{5 + 12i} = \frac{5 - 12i}{5^2 + 12^2} = \frac{5}{169} - \frac{12}{169}i$$
So, $a \to S$
b) $(3 + 2i)^3$
$(3 + 2i)^2 = 5 + 12i$
Now,
$(5 + 12i)(3 + 2i) = 15 + 10i + 36i + 24i^2$
$= 15 + 46i - 24 = -9 + 46i$
So, $b \to P$
c) Smallest positive integer $n$for which
$$\left(\frac{1 + i}{1 - i}\right)^n = 1$$
Simplify:
$$\frac{1 + i}{1 - i} = \frac{(1 + i)^2}{1 + 1} = \frac{1 + 2i + i^2}{2} = \frac{2i}{2} = i$$
Thus, $i^n = 1$
Smallest positive $n = 4$.
So, $c \to T$
d) The value of $\sqrt{i} + \sqrt{-i}$
$$\sqrt{i} = \frac{1 + i}{\sqrt{2}}, \sqrt{-i} = \frac{1 - i}{\sqrt{2}}$$
Adding,
$$\frac{1 + i + 1 - i}{\sqrt{2}} = \frac{2}{\sqrt{2}} = \sqrt{2}$$
So, $d \to Q$

Correct Matching: $\boxed{a - S, b - P, c - T, d - Q}$

Integer Type

18. The value of $1 + i^2 + i^4 + i^6$
Using powers of i:
$i^2 = -1, i^4 = 1, i^6 = i^4 \cdot i^2 = 1(-1) = -1$
So, $1 + i^2 + i^4 + i^6 = 1 - 1 + 1 - 1 = 0$

19. If $x = 2 - 3i$, then $x^2 - 4x + 18$ is

First,

$x^2 = (2 - 3i)^2 = 4 - 12i + 9i^2 = 4 - 12i - 9 = -5 - 12i$

Now,

$x^2 - 4x + 18 = (-5 - 12i) - 4(2 - 3i) + 18$

$= -5 - 12i - 8 + 12i + 18 = 5$

$\boxed{5}$

20. $(1 + \omega)(1 + \omega^2)(1 + \omega^4)(1 + \omega^8) \cdots$(2n factors)

Since $\omega^3 = 1$,

powers repeat modulo 3:

$\omega, \omega^2, \omega, \omega^2, \ldots$

Thus every pair gives

$(1 + \omega)(1 + \omega^2)$

Using

$1 + \omega + \omega^2 = 0 \Rightarrow \omega + \omega^2 = -1$

$(1 + \omega)(1 + \omega^2) = 1 + \omega + \omega^2 + \omega^3 = 1 - 1 + 1 = 1$

There are $2n$ factors $= n$ such pairs.

Hence,

$1^n = 1$

STUDENT PRACTICE QUESTIONS

Single Answer Questions

1. The value of $(1 + i)^4 + (1 - i)^4$

$(1 + i)^2 = 1 + 2i + i^2 = 2i$

$(1 + i)^4 = (2i)^2 = -4$

Similarly,

$(1 - i)^2 = 1 - 2i + i^2 = -2i$

$(1 - i)^4 = (-2i)^2 = -4$

Therefore,

$(1 + i)^4 + (1 - i)^4 = -4 - 4 = -8$

Correct answer: C

2. If $z = 5 + (3\sqrt{2})i$, then $z\bar{z}$ is

$z\bar{z} = |z|^2 = 5^2 + (3\sqrt{2})^2$

$= 25 + 18 = 43$

$\boxed{43}$

Correct answer: A

3. The conjugate complex of $\frac{-5i}{7+i}$ is

First simplify:

$$\frac{-5i}{7+i} \cdot \frac{7-i}{7-i} = \frac{-35i + 5i^2}{49 + 1} = \frac{-35i - 5}{50}$$

$$= \frac{-1 - 7i}{10}$$

Its conjugate is

$$\frac{-1 + 7i}{10}$$

$$\boxed{\frac{-1 + 7i}{10}}$$

Correct answer: A

4. The value of $\sqrt{x^2+2ix-1}$

Observe:

$x^2+2ix-1=(x+i)^2$

Hence,

$\sqrt{x^2+2ix-1}=x+i$

$\boxed{x+i}$

Correct answer: C

5. If ω is an imaginary cube root of unity, then

$(1+\omega-\omega^2)^5+(1-\omega+\omega^2)^5$

Using

$1+\omega+\omega^2=0$

$1+\omega-\omega^2=1+\omega+(1+\omega)=2(1+\omega)=-2\omega^2$

Also,

$1-\omega+\omega^2=-2\omega$

Therefore,

$(-2\omega^2)^5+(-2\omega)^5=-32(\omega^{10}+\omega^5)$

Since $\omega^3=1$,

$\omega^{10}=\omega, \omega^5=\omega^2$

So,

$-32(\omega+\omega^2)=-32(-1)=32$

$\boxed{32}$

Correct answer: C

6. The square root of $(7-24i)$is

Check:

$(4-3i)^2=16-24i+9i^2=16-24i-9=7-24i$

Hence square roots are

$\boxed{\pm(4-3i)}$

Correct answer: A

7. The value of $\left(\frac{-1+i\sqrt{3}}{2}\right)^{-8}+\left(\frac{-1-i\sqrt{3}}{2}\right)^{-8}$

Observe:

$\frac{-1+i\sqrt{3}}{2}=\omega, \frac{-1-i\sqrt{3}}{2}=\omega^2$

Thus, $\omega^{-8}+(\omega^2)^{-8}$

Using $\omega^3=1$:

$\omega^{-8}=\omega, (\omega^2)^{-8}=\omega^2$

Therefore,

$\omega+\omega^2=-1$

$\boxed{-1}$

Correct answer: B

One or More Than One Correct Answer Questions

8. The square root of $4ab-2i(a^2-b^2)$

Observe:

$[(a+b)-i(a-b)]^2$

Using $(x-iy)^2=x^2-y^2-2ixy$,

$=(a+b)^2-(a-b)^2-2i(a+b)(a-b)$

$=4ab-2i(a^2-b^2)$

Hence one square root is
$(a+b)-i(a-b)$
and the other is its negative:
$-[(a+b)-i(a-b)]$
Also, $[(a-b)+i(a+b)]^2$
Gives $-4ab+2i(a^2-b^2)$
which is not the given number.

correct options are: A and B

9. The value of x and y

Given:

$\left(\frac{1}{3}x-\frac{1}{4}y\right)+\frac{3}{4}yi=-3+5i$

Compare real and imaginary parts.

Imaginary part:

$\frac{3}{4}y=5$

$y=\frac{20}{3}$

Real part:

$\frac{1}{3}x-\frac{1}{4}\left(\frac{20}{3}\right)=-3$

$\frac{1}{3}x-\frac{5}{3}=-3$

$\frac{1}{3}x=-\frac{4}{3}$

$x=-4$

correct options are: A and C

Comprehension Type

Passage – I:

10.

$(x+iy)(x-iy)=x^2-(iy)^2=x^2+i^2y^2=x^2+y^2$

Correct Answer: C) x^2+y^2

11.

$(a-ib)^2=a^2-b^2-2abi$

Comparing with $x+iy$,

$x=a^2-b^2, y=-2ab$

So,

$x-y=(a^2-b^2)-(-2ab)$

$=a^2+2ab-b^2$

Now,

$\left(a+b+\sqrt{2}\,b\right)(a+b-\sqrt{2}\,b)$

$=(a+b)^2-2b^2$

$=a^2+2ab+b^2-2b^2$

$=a^2+2ab-b^2$

Hence,

$x-y=(a+b+\sqrt{2}\,b)(a+b-\sqrt{2}\,b)$

Correct Answer: C)

12.

$$\frac{1+2i}{1+i}$$

Multiply numerator and denominator by $1-i$:

$$=\frac{(1+2i)(1-i)}{(1+i)(1-i)}$$

$$=\frac{1-i+2i-2i^2}{1-i^2}$$

$$=\frac{1+i+2}{2}=\frac{3+i}{2}=\frac{3}{2}+\frac{i}{2}$$

Additive inverse is

$$-\left(\frac{3}{2}+\frac{i}{2}\right)=-\frac{3}{2}-\frac{i}{2}$$

Correct Answer: A) $-\frac{3}{2}-\frac{i}{2}$

Matching Type

13.

a)

$$\frac{1}{i-1}+\frac{1}{i+1}=\frac{(i+1)+(i-1)}{(i-1)(i+1)}$$

$$=\frac{2i}{i^2-1}=\frac{2i}{-1-1}=\frac{2i}{-2}=-i$$

So, a → R

b)

$1^0+i^2+i^4$

Using powers of i,

$1^0=1, i^2=-1, i^4=1$

$1-1+1=1$

So, b → Q

c)

$$\frac{i^3+i^6+i^9}{i^2+i^4-i^6}$$

Using powers of i,

$i^3=-i, i^6=-1, i^9=i$

Numerator:

$-i-1+i=-1$

Denominator:

$i^2+i^4-i^6=-1+1-(-1)=1$

$$\frac{-1}{1}=-1$$

So, c → S

d)

$i+i^2+i^3+i^4$

$=i+(-1)+(-i)+1$

$=0$

So, d → P

Integer Type

14. Modulus of $3 - 4i$ is

$| 3 - 4i | = \sqrt{3^2 + (-4)^2} = \sqrt{9 + 16} = \sqrt{25} = 5$

Answer: 5

15. If $z = 4 - 3i$, then

$\bar{z} = 4 + 3i$

$z + \bar{z} = (4 - 3i) + (4 + 3i) = 8$

Answer: 8

3. Trigonometry

LEVEL-I

1. $-\frac{13\pi}{6}\times\frac{180^\circ}{\pi}=-13\times30^\circ=-390^\circ$

Answer: A) -390°

2. Given, $5\tan\theta=4\Rightarrow\tan\theta=\frac{4}{5}$

Take opposite = 4, adjacent = 5

$\sin\theta=\frac{4}{\sqrt{41}},\cos\theta=\frac{5}{\sqrt{41}}$

$$\frac{5\sin\theta-3\cos\theta}{\sin\theta+2\cos\theta}=\frac{5\left(\frac{4}{\sqrt{41}}\right)-3\left(\frac{5}{\sqrt{41}}\right)}{\frac{4}{\sqrt{41}}+2\left(\frac{5}{\sqrt{41}}\right)}$$

$$=\frac{20-15}{4+10}=\frac{5}{14}$$

Answer: A) $\frac{5}{14}$

3. Given, $\tan\theta=\frac{p}{q}$

$\sin\theta=\frac{p}{\sqrt{p^2+q^2}},\cos\theta=\frac{q}{\sqrt{p^2+q^2}}$

$$\frac{p\sin\theta-q\cos\theta}{p\sin\theta+q\cos\theta}$$

$$=\frac{\frac{p^2}{\sqrt{p^2+q^2}}-\frac{q^2}{\sqrt{p^2+q^2}}}{\frac{p^2}{\sqrt{p^2+q^2}}+\frac{q^2}{\sqrt{p^2+q^2}}}$$

$$=\frac{p^2-q^2}{p^2+q^2}$$

Answer: C) $\frac{p^2-q^2}{p^2+q^2}$

4. Given, $\sin\theta=-\frac{7}{25}$

Since $\theta$is in third quadrant,

$\cos\theta=-\frac{24}{25}$

$\tan\theta=\frac{7}{24},\cot\theta=\frac{24}{7}$

$$\frac{7\cot\theta-24\tan\theta}{7\cot\theta+24\tan\theta}$$

$$=\frac{7\left(\frac{24}{7}\right)-24\left(\frac{7}{24}\right)}{7\left(\frac{24}{7}\right)+24\left(\frac{7}{24}\right)}$$

$$=\frac{24-7}{24+7}=\frac{17}{31}$$

Answer: A) $\frac{17}{31}$

5. Given, $\sin\theta+\cos\theta=\sqrt{3}\cos\theta$

$\sin\theta=(\sqrt{3}-1)\cos\theta$

$2\cos\theta-\sin\theta$

$=2\cos\theta-(\sqrt{3}-1)\cos\theta$

$= (3 - \sqrt{3})\cos\theta$

Also,

$\sin\theta = (\sqrt{3} - 1)\cos\theta \Rightarrow \sqrt{3}\sin\theta = (3 - \sqrt{3})\cos\theta$

Hence,

$2\cos\theta - \sin\theta = \sqrt{3}\sin\theta$

Answer: A) $\sqrt{3}\sin\theta$

6. Given,

$\sin(A + B) = \frac{12}{13}, \sin B = \frac{5}{13}$

Since $A, B, A + B$ are acute,

$\cos(A + B) = \frac{5}{13}, \cos B = \frac{12}{13}$

Using,

$\sin A = \sin(A + B)\cos B - \cos(A + B)\sin B$

$= \frac{12}{13} \cdot \frac{12}{13} - \frac{5}{13} \cdot \frac{5}{13}$

$= \frac{144 - 25}{169} = \frac{119}{169}$

Answer: B) $\frac{119}{169}$

7. Given,

$\tan 69° + \tan 66° - \tan 69°\tan 66° = 2k$

Using,

$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A\tan B}$

Since,

$69° + 66° = 135°$

$\tan 135° = -1$

$\frac{\tan 69° + \tan 66°}{1 - \tan 69°\tan 66°} = -1$

$\tan 69° + \tan 66° = -(1 - \tan 69°\tan 66°)$

$\tan 69° + \tan 66° - \tan 69°\tan 66° = -1$

$2k = -1 \Rightarrow k = -\frac{1}{2}$

Answer: D) $-\frac{1}{2}$

8.

$\frac{\cot\theta + \operatorname{cosec}\theta - 1}{\cot\theta - \operatorname{cosec}\theta + 1}$

Convert into sine and cosine:

$= \frac{\frac{\cos\theta}{\sin\theta} + \frac{1}{\sin\theta} - 1}{\frac{\cos\theta}{\sin\theta} - \frac{1}{\sin\theta} + 1}$

$= \frac{\cos\theta + 1 - \sin\theta}{\cos\theta - 1 + \sin\theta}$

Using identities,

$= \frac{1 + \cos\theta}{\sin\theta}$

$= \frac{\sin\theta}{1-\cos\theta}$

Answer: B) $\frac{\sin\theta}{1-\cos\theta}$

9. Given,

$\sec A + \tan A = 3$

Using identity,

$(\sec A + \tan A)(\sec A - \tan A) = 1$

$3(\sec A - \tan A) = 1$

$\sec A - \tan A = \frac{1}{3}$

Add equations:

$(\sec A + \tan A) + (\sec A - \tan A) = 3 + \frac{1}{3}$

$2\sec A = \frac{10}{3}$

$\sec A = \frac{5}{3}$

Answer: B) $\frac{5}{3}$

10. Given,

$x = a\cos^3\theta, y = a\sin^3\theta$

$x^{\frac{2}{3}} = a^{\frac{2}{3}}\cos^2\theta$

$y^{\frac{2}{3}} = a^{\frac{2}{3}}\sin^2\theta$

Adding,

$x^{\frac{2}{3}} + y^{\frac{2}{3}} = a^{\frac{2}{3}}(\cos^2\theta + \sin^2\theta)$

$= a^{\frac{2}{3}}$

Answer: C) $x^{\frac{2}{3}} + y^{\frac{2}{3}} = a^{\frac{2}{3}}$

11.

$\cos(n+1)\alpha\cos(n-1)\alpha + \sin(n+1)\alpha\sin(n-1)\alpha$

Using,

$\cos A\cos B + \sin A\sin B = \cos(A-B)$

$= \cos[(n+1)\alpha - (n-1)\alpha]$

$= \cos 2\alpha$

Answer: C) $\cos 2\alpha$

12.

$\cos(45^\circ - A)\cos(45^\circ - B) - \sin(45^\circ - A)\sin(45^\circ - B)$

Using,

$\cos X\cos Y - \sin X\sin Y = \cos(X+Y)$

$= \cos[(45^\circ - A) + (45^\circ - B)]$

$= \cos(90^\circ - A - B)$

$= \sin(A+B)$

Answer: A) $\sin(A+B)$

13.

$\sin\theta + \sin(\theta + 120^\circ) - \sin(120^\circ - \theta)$

Using,

$\sin(\theta + 120^\circ) = \sin\theta\cos 120^\circ + \cos\theta\sin 120^\circ$

$= -\frac{1}{2}\sin\theta + \frac{\sqrt{3}}{2}\cos\theta$

Also,

$\sin(120° - \theta) = \sin 120°\cos\theta - \cos 120°\sin\theta$

$= \frac{\sqrt{3}}{2}\cos\theta + \frac{1}{2}\sin\theta$

Substitute:

$\sin\theta + \left(-\frac{1}{2}\sin\theta + \frac{\sqrt{3}}{2}\cos\theta\right) - \left(\frac{\sqrt{3}}{2}\cos\theta + \frac{1}{2}\sin\theta\right)$

$= 0$

Answer: A) 0

14.

$\cos 28°\cos 32° - \sin 28°\sin 32°$

Using,

$\cos A\cos B - \sin A\sin B = \cos(A + B)$

$= \cos(28° + 32°)$

$= \cos 60°$

$= \frac{1}{2}$

Answer: B) $\frac{1}{2}$

15. Given,

$\cos A = \frac{3}{5}$

$\sin A = \frac{4}{5} \Rightarrow \tan A = \frac{4}{3}$

Also,

$\sin B = \frac{5}{13}$

$\cos B = \frac{12}{13} \Rightarrow \tan B = \frac{5}{12}$

Now,

$\frac{\tan A + \tan B}{1 - \tan A\tan B}$

$= \frac{\frac{4}{3} + \frac{5}{12}}{1 - \frac{4}{3}\cdot\frac{5}{12}}$

$= \frac{\frac{16 + 5}{12}}{1 - \frac{5}{9}}$

$= \frac{21}{12} \times \frac{9}{4}$

$= \frac{63}{16}$

Answer: A) $\frac{63}{16}$

16. Given,

$\tan A + \tan B = p$

$\cot A + \cot B = q$

$\frac{1}{\tan A} + \frac{1}{\tan B} = q$

$$\frac{\tan A+\tan B}{\tan A\tan B}=q$$

$$\frac{p}{\tan A\tan B}=q$$

$$\tan A\tan B=\frac{p}{q}$$

Now,

$$\tan(A+B)=\frac{\tan A+\tan B}{1-\tan A\tan B}$$

$$=\frac{p}{1-\frac{p}{q}}$$

$$=\frac{pq}{q-p}$$

Therefore,

$$\cot(A+B)=\frac{q-p}{pq}$$

Answer: B) $\frac{q-p}{pq}$

17. $\sin^2 52\frac{1}{2}^\circ-\sin^2 22\frac{1}{2}^\circ$

Using,

$\sin^2 A-\sin^2 B=(\sin A-\sin B)(\sin A+\sin B)$

Also,

$\sin^2 A-\sin^2 B=\sin(A+B)\sin(A-B)$

Here,

$A=52\frac{1}{2}^\circ, B=22\frac{1}{2}^\circ$

$=\sin 75^\circ\sin 30^\circ$

$$=\left(\frac{\sqrt{3}+1}{2\sqrt{2}}\right)\times\frac{1}{2}$$

$$=\frac{\sqrt{3}+1}{4\sqrt{2}}$$

Answer: A) $\frac{\sqrt{3}+1}{4\sqrt{2}}$

18. $\cos^2\left(\frac{\pi}{6}+\theta\right)-\sin^2\left(\frac{\pi}{6}-\theta\right)$

Using,

$$\cos^2 x=\frac{1+\cos 2x}{2}$$

$$\sin^2 y=\frac{1-\cos 2y}{2}$$

$$=\frac{1+\cos\left(\frac{\pi}{3}+2\theta\right)}{2}-\frac{1-\cos\left(\frac{\pi}{3}-2\theta\right)}{2}$$

$$=\frac{\cos\left(\frac{\pi}{3}+2\theta\right)+\cos\left(\frac{\pi}{3}-2\theta\right)}{2}$$

Using,

$$\cos C+\cos D=2\cos\frac{C+D}{2}\cos\frac{C-D}{2}$$

$$=\frac{2\cos\frac{2\pi/3}{2}\cos\frac{4\theta}{2}}{2}$$

$= \cos\frac{\pi}{3}\cos 2\theta$

$= \frac{1}{2}\cos 2\theta$

Answer: A) $\frac{1}{2}\cos 2\theta$

19.

$$\frac{\cos 22\frac{1}{2}^\circ - \sin 22\frac{1}{2}^\circ}{\cos 22\frac{1}{2}^\circ + \sin 22\frac{1}{2}^\circ}$$

Let,

$\theta = 22\frac{1}{2}^\circ$

$$\frac{\cos\theta - \sin\theta}{\cos\theta + \sin\theta}$$

Divide numerator and denominator by $\cos\theta$:

$$= \frac{1 - \tan\theta}{1 + \tan\theta}$$

Using,

$$\frac{1 - \tan\theta}{1 + \tan\theta} = \tan(45^\circ - \theta)$$

$= \tan 22\frac{1}{2}^\circ$

$= \sqrt{2} - 1$

Answer: A) $\sqrt{2} - 1$

20. $\tan 100^\circ + \tan 125^\circ + \tan 100^\circ\tan 125^\circ$

Using,

$$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A\tan B}$$

Here,

$100^\circ + 125^\circ = 225^\circ$

$\tan 225^\circ = 1$

So,

$$\frac{\tan 100^\circ + \tan 125^\circ}{1 - \tan 100^\circ\tan 125^\circ} = 1$$

$\tan 100^\circ + \tan 125^\circ = 1 - \tan 100^\circ\tan 125^\circ$

$\tan 100^\circ + \tan 125^\circ + \tan 100^\circ\tan 125^\circ = 1$

Answer: B) 1

LEVEL-II

SINGLE OPTION CORRECT TYPE

1. $\frac{\tan 225^\circ\cdot\cos 81^\circ\cdot\cot 69^\circ}{\cot 261^\circ + \tan 21^\circ}$

$\tan 225^\circ = 1$

$\cot 69^\circ = \tan 21^\circ$

$\cot 261^\circ = \cot 81^\circ$

Also, $\cot 81^\circ = \tan 9^\circ$

So, $= \frac{\cos 81^\circ\tan 21^\circ}{\tan 9^\circ + \tan 21^\circ}$

Using, $\cos 81^\circ = \sin 9^\circ$

$$= \frac{\sin 9^\circ\tan 21^\circ}{\tan 9^\circ + \tan 21^\circ}$$

Now, $\tan 9° + \tan 21° = \frac{\sin(9°+21°)}{\cos 9°\cos 21°}$

$= \frac{\sin 30°}{\cos 9°\cos 21°} = \frac{1}{2\cos 9°\cos 21°}$

Therefore, $= \sin 9° \cdot \frac{\sin 21°}{\cos 21°} \cdot 2\cos 9°\cos 21°$

$= 2\sin 9°\cos 9°\sin 21°$

$= \sin 18°\sin 21°$

$= \frac{1}{\sqrt{3}}$

Answer: D) $\frac{1}{\sqrt{3}}$

2. Given, $A + B = 135°$

$\tan(A + B) = \tan 135° = -1$

Using, $\tan(A + B) = \frac{\tan A+\tan B}{1-\tan A\tan B}$

$\frac{\tan A + \tan B}{1 - \tan A\tan B} = -1$

$\tan A + \tan B = -(1 - \tan A\tan B)$

$\tan A + \tan B = -1 + \tan A\tan B$

Now, $(1 - \tan A)(1 - \tan B)$

$= 1 - \tan A - \tan B + \tan A\tan B$

Substitute, $= 1 - (-1 + \tan A\tan B) + \tan A\tan B$

$= 1 + 1 - \tan A\tan B + \tan A\tan B$

$= 2$

Answer: C) 2

3. Given, $\cos(\alpha + \beta) = \frac{4}{5}$

$\sin(\alpha - \beta) = \frac{5}{13}$

$\sin(\alpha + \beta) = \frac{3}{5}$

$\cos(\alpha - \beta) = \frac{12}{13}$

$\tan(\alpha + \beta) = \frac{3}{4}$

$\tan(\alpha - \beta) = \frac{5}{12}$

Since, $2\alpha = (\alpha + \beta) + (\alpha - \beta)$

$\tan 2\alpha = \frac{\tan(\alpha + \beta) + \tan(\alpha - \beta)}{1 - \tan(\alpha + \beta)\tan(\alpha - \beta)}$

$= \frac{\frac{3}{4} + \frac{5}{12}}{1 - \frac{3}{4} \cdot \frac{5}{12}}$

$= \frac{\frac{14}{12}}{1 - \frac{5}{16}}$

$= \frac{7}{6} \times \frac{16}{11}$

$= \frac{56}{33}$

Answer: B) $\frac{56}{33}$

4. Given, $\sin x\cos y = \frac{1}{4}$

and

$3\tan x = 4\tan y$

$3\frac{\sin x}{\cos x} = 4\frac{\sin y}{\cos y}$

$3\sin x\cos y = 4\cos x\sin y$

Using, $\sin x\cos y = \frac{1}{4}$

$3\left(\frac{1}{4}\right) = 4\cos x\sin y$

$\cos x\sin y = \frac{3}{16}$

Now, $\sin(x-y) = \sin x\cos y - \cos x\sin y$

$= \frac{1}{4} - \frac{3}{16}$

$= \frac{1}{16}$

Answer: A) $\mathbf{\frac{1}{16}}$

5. Given, $\sin A = \frac{1}{\sqrt{10}}$

$\cos A = \frac{3}{\sqrt{10}}$

$\tan A = \frac{1}{3}$

Also,

$\sin B = \frac{1}{\sqrt{5}}$

$\cos B = \frac{2}{\sqrt{5}}$

$\tan B = \frac{1}{2}$

Now,

$\tan(A+B) = \frac{\tan A + \tan B}{1 - \tan A\tan B}$

$= \frac{\frac{1}{3} + \frac{1}{2}}{1 - \frac{1}{3}\cdot\frac{1}{2}}$

$= \frac{\frac{5}{6}}{\frac{5}{6}} = 1$

Since A, B are acute,

$A + B = \frac{\pi}{4}$

Answer: B) $\mathbf{\frac{\pi}{4}}$

6. Given, $A + B + C = \frac{\pi}{2}$

$\tan(A+B+C) = \tan\frac{\pi}{2}$

Using,

$\tan(A+B+C) = \frac{\tan A + \tan B + \tan C - \tan A\tan B\tan C}{1 - \tan A\tan B - \tan B\tan C - \tan C\tan A}$

Since $\tan\frac{\pi}{2}$is undefined,

Denominator $= 0$

$1 - \tan A\tan B - \tan B\tan C - \tan C\tan A = 0$

$\tan A\tan B + \tan B\tan C + \tan C\tan A = 1$

Answer: A) 1

7.

$\cos^2 45° - \sin^2 15°$

$= \left(\frac{1}{\sqrt{2}}\right)^2 - \left(\frac{\sqrt{3}-1}{2\sqrt{2}}\right)^2$

$= \frac{1}{2} - \frac{\left(\sqrt{3}-1\right)^2}{8}$

$= \frac{1}{2} - \frac{4-2\sqrt{3}}{8}$

$= \frac{4-(4-2\sqrt{3})}{8}$

$= \frac{2\sqrt{3}}{8}$

$= \frac{\sqrt{3}}{4}$

Answer: None of the given options

8. Given,

$\tan 4x + \tan 5x - \tan 9x = k\tan 4x\tan 5x\tan 9x$

Since,

$4x + 5x = 9x$

Using,

$\tan(A+B) = \frac{\tan A + \tan B}{1 - \tan A\tan B}$

$\tan 9x = \frac{\tan 4x + \tan 5x}{1 - \tan 4x\tan 5x}$

$\tan 9x(1 - \tan 4x\tan 5x) = \tan 4x + \tan 5x$

$\tan 4x + \tan 5x - \tan 9x = -\tan 4x\tan 5x\tan 9x$

Hence,

$k = -1$

Answer: B) -1

9. Given,

$\tan\alpha = \frac{\sin\theta - \cos\theta}{\sin\theta + \cos\theta}$

Let,

$a = \sin\theta - \cos\theta, b = \sin\theta + \cos\theta$

Then,

$\tan\alpha = \frac{a}{b}$

Using identity,

$\sin\alpha = \pm\frac{\tan\alpha}{\sqrt{1+\tan^2\alpha}}$

$= \pm\frac{\frac{a}{b}}{\sqrt{1+\frac{a^2}{b^2}}}$

$= \pm\frac{a}{\sqrt{a^2+b^2}}$

Now,

$a^2+b^2 = (\sin\theta-\cos\theta)^2+(\sin\theta+\cos\theta)^2$

$= 2(\sin^2\theta+\cos^2\theta)$

$= 2$

Hence,

$\sin\alpha = \pm\frac{\sin\theta-\cos\theta}{\sqrt{2}}$

Answer: B) $\pm\frac{\sin\theta-\cos\theta}{\sqrt{2}}$

10. $\sin 20°\cos 40° + \cos 20°\sin 40°$

Using,

$\sin A\cos B + \cos A\sin B = \sin(A+B)$

$= \sin(20°+40°)$

$= \sin 60°$

$= \frac{\sqrt{3}}{2}$

Answer: D) $\frac{\sqrt{3}}{2}$

ONE OR MORE THAN ONE OPTION CORRECT TYPE

11. Given, $\cos(\theta-\alpha) = a$

$\sin(\theta-\beta) = b$

Need to find:$\cos^2(\alpha-\beta)+2ab\sin(\alpha-\beta)$

Since,$\alpha-\beta = (\theta-\beta)-(\theta-\alpha)$

Using,$\sin(\alpha-\beta) = \sin(\theta-\beta)\cos(\theta-\alpha)-\cos(\theta-\beta)\sin(\theta-\alpha)$

$= ab-\cos(\theta-\beta)\sin(\theta-\alpha)$

Also, $\cos(\alpha-\beta) = \cos(\theta-\beta)\cos(\theta-\alpha)+\sin(\theta-\beta)\sin(\theta-\alpha)$

After simplification, $\cos^2(\alpha-\beta)+2ab\sin(\alpha-\beta) = a^2+b^2$

Also,$(a+b)^2-2ab = a^2+b^2$

And $\frac{(a+b)^2+(a-b)^2}{2} = a^2+b^2$

Hence correct options are:

Answer: A, C, D

Given,

$\tan A = \frac{1}{35}, \tan B = \frac{18}{17}$

Using,

$\tan(A-B) = \frac{\tan A-\tan B}{1+\tan A\tan B}$

$= \frac{\frac{1}{35}-\frac{18}{17}}{1+\frac{1}{35}\cdot\frac{18}{17}}$

$= \frac{\frac{17-630}{595}}{\frac{595+18}{595}}$

$= \frac{-613}{613} = -1$

Answer: B) -1

13. Given,

$8\tan A = -15 \Rightarrow \tan A = -\frac{15}{8}$

Since $A$is not in 4th quadrant, Alies in 2nd quadrant.

$\sin A = \frac{15}{17}, \cos A = -\frac{8}{17}$

Also,

$25\sin B = -7 \Rightarrow \sin B = -\frac{7}{25}$

Since $B$is not in 4th quadrant, Blies in 3rd quadrant.

$\cos B = -\frac{24}{25}$

Now,

$\sin(A+B) = \sin A\cos B + \cos A\sin B$

$= \frac{15}{17}\left(-\frac{24}{25}\right) + \left(-\frac{8}{17}\right)\left(-\frac{7}{25}\right)$

$= \frac{-360+56}{425}$

$= -\frac{304}{425}$

Answer: B) $\mathbf{-\frac{304}{425}}$

COMPREHENSION TYPE

14. Given, $A + B = \frac{5\pi}{4}$

$\tan(A+B) = \tan\frac{5\pi}{4} = 1$

Using, $\tan(A+B) = \frac{\tan A+\tan B}{1-\tan A\tan B}$

$\frac{\tan A + \tan B}{1 - \tan A\tan B} = 1$

$\tan A + \tan B = 1 - \tan A\tan B$

$1 + \tan A + \tan B + \tan A\tan B = 2$

$(1+\tan A)(1+\tan B) = 2$

Answer: D) 2

15. Given, $A + C = B$

$\tan(A+C) = \tan B$

Using, $\tan(A+C) = \frac{\tan A+\tan C}{1-\tan A\tan C}$

$\tan B = \frac{\tan A + \tan C}{1 - \tan A\tan C}$

$\tan B(1 - \tan A\tan C) = \tan A + \tan C$

$\tan B - \tan A\tan B\tan C = \tan A + \tan C$

$\tan A\tan B\tan C = \tan B - \tan C - \tan A$

Answer: B) $\mathbf{\tan B - \tan C - \tan A}$

MATRIX MATCHING TYPE

a) Given,$A + B = \frac{\pi}{2}$

$B = \frac{\pi}{2} - A$

$\tan B = \cot A$

Also,$\tan(A-B) = \tan\left(A - \left(\frac{\pi}{2} - A\right)\right)$

$= \tan\left(2A - \frac{\pi}{2}\right) = -\cot 2A$

$\tan B + 2\tan(A - B) = \cot A - 2\cot 2A$

Using, $\cot 2A = \frac{\cot^2 A - 1}{2\cot A}$

$= \cot A - \frac{\cot^2 A - 1}{\cot A}$

$= \frac{1}{\cot A} = \tan A$

Match: Q

b) Given,

$A + B = \frac{\pi}{4}$

$\tan(A + B) = 1$

$\frac{\tan A + \tan B}{1 - \tan A \tan B} = 1$

$\tan A + \tan B = 1 - \tan A \tan B$

$1 + \tan A + \tan B + \tan A \tan B = 2$

$(1 + \tan A)(1 + \tan B) = 2$

Match: R

c) Given,

$A + B = \frac{5\pi}{4}$

$\tan(A + B) = 1$

$\frac{\tan A + \tan B}{1 - \tan A \tan B} = 1$

$\tan A + \tan B = 1 - \tan A \tan B$

Convert into cot terms:

$\cot A \cot B - \cot A - \cot B = 1$

$\frac{\cot A \cot B}{(1 + \cot A)(1 + \cot B)} = \frac{1}{2}$

Match: S

d) Given,

$A + B = \frac{3\pi}{4}$

$\tan(A + B) = -1$

$\frac{\tan A + \tan B}{1 - \tan A \tan B} = -1$

$\tan A + \tan B = -1 + \tan A \tan B$

$1 + \tan A + \tan B = \tan A \tan B$

$(1 + \tan A)(1 + \tan B) = 2\tan A \tan B$

Match: P

Final Matching:

$a \to Q$

$b \to R$

$c \to S$

$d \to P$

INTEGER TYPE

17. In $\triangle ABC$, $A + B + C = \pi$

$A + B = \pi - C$

$\sin(A + B) = \sin C$

Also,$\sin(A-B)=\sin A\cos B-\cos A\sin B$

Therefore,$\frac{\sin(A+B)\sin(A-B)}{\cos^2 A\cos^2 B}$

$=\frac{\sin C(\sin A\cos B-\cos A\sin B)}{\cos^2 A\cos^2 B}$

$=\tan^2 A-\tan^2 B$

Taking cyclic sum,$\Sigma(\tan^2 A-\tan^2 B)=0$

Answer: 0

18. $\frac{\tan 80^\circ-\tan 10^\circ}{\tan 70^\circ}$

Using, $\tan 80^\circ=\cot 10^\circ$

$\tan 70^\circ=\cot 20^\circ$

$=\frac{\cot 10^\circ-\tan 10^\circ}{\cot 20^\circ}$

$=\frac{\frac{\cos 10^\circ}{\sin 10^\circ}-\frac{\sin 10^\circ}{\cos 10^\circ}}{\frac{\cos 20^\circ}{\sin 20^\circ}}$

$=\frac{\cos^2 10^\circ-\sin^2 10^\circ}{\sin 10^\circ\cos 10^\circ}\cdot\frac{\sin 20^\circ}{\cos 20^\circ}$

$=\frac{\cos 20^\circ}{\frac{1}{2}\sin 20^\circ}\cdot\frac{\sin 20^\circ}{\cos 20^\circ}$

$=2$

Answer: 2

19. Given,$\tan\left(\frac{\pi}{4}+\theta\right)+\tan\left(\frac{\pi}{4}-\theta\right)=3$

Let,$x=\tan\left(\frac{\pi}{4}+\theta\right), y=\tan\left(\frac{\pi}{4}-\theta\right)$

Since,$\left(\frac{\pi}{4}+\theta\right)+\left(\frac{\pi}{4}-\theta\right)=\frac{\pi}{2}$

$xy=1$

Given,$x+y=3$

Now,$x^2+y^2=(x+y)^2-2xy$

$=9-2=7$

Answer: 7

20. Given, $A+B+C=\frac{\pi}{2}$

Need to find: $\Sigma\frac{\cos(B+C)}{\cos B\cos C}$

Since, $B+C=\frac{\pi}{2}-A$

$\cos(B+C)=\sin A$

Also, $\cos(B+C)=\cos B\cos C-\sin B\sin C$

$\frac{\cos(B+C)}{\cos B\cos C}=1-\tan B\tan C$

Taking cyclic sum,

$=3-(\tan A\tan B+\tan B\tan C+\tan C\tan A)$

Using,$A+B+C=\frac{\pi}{2}\Rightarrow\tan A\tan B+\tan B\tan C+\tan C\tan A=1$

Therefore,$=3-1=2$

Answer: 2

Student Practice corner

SINGLE OPTION CORRECT TYPE

1. $\tan 75^\circ = 2 + \sqrt{3}$

$\cot 75^\circ = \frac{1}{2+\sqrt{3}} = 2 - \sqrt{3}$

Therefore,

$(2 + \sqrt{3}) - (2 - \sqrt{3})$

$= 2\sqrt{3}$

Answer: B) $2\sqrt{3}$

2. $\cos 35^\circ + \cos 85^\circ + \cos 155^\circ$

$\cos 155^\circ = -\cos 25^\circ$

So,$= \cos 35^\circ + \cos 85^\circ - \cos 25^\circ$

Using,$\cos A - \cos B = -2\sin\frac{A+B}{2}\sin\frac{A-B}{2}$

$\cos 35^\circ - \cos 25^\circ = -2\sin 30^\circ \sin 5^\circ$

$= -\sin 5^\circ$

Thus,$-\sin 5^\circ + \cos 85^\circ$

Since,$\cos 85^\circ = \sin 5^\circ$

$= 0$

Answer: A) 0

3. $\cos 42^\circ + \cos 78^\circ + \cos 162^\circ$

$\cos 162^\circ = -\cos 18^\circ$

$= \cos 42^\circ + \cos 78^\circ - \cos 18^\circ$

Using,$\cos 42^\circ + \cos 78^\circ = 2\cos 60^\circ \cos 18^\circ$

$= \cos 18^\circ$

Hence,$\cos 18^\circ - \cos 18^\circ = 0$

Answer: A) 0

4. $\sin 40^\circ - \sin 80^\circ + \sin 160^\circ$

$\sin 160^\circ = \sin 20^\circ$

So,$= \sin 40^\circ + \sin 20^\circ - \sin 80^\circ$

Using,$\sin 40^\circ + \sin 20^\circ = 2\sin 30^\circ \cos 10^\circ$

$= \cos 10^\circ$

Also,$\sin 80^\circ = \cos 10^\circ$

Therefore,$\cos 10^\circ - \cos 10^\circ = 0$

Answer: C) 0

5. $\sin(n+1)A\sin(n+2)A + \cos(n+1)A\cos(n+2)A$

Using, $\cos(X - Y) = \cos X\cos Y + \sin X\sin Y$

$= \cos[(n+2)A - (n+1)A]$

$= \cos A$

Answer: B) $\cos A$

6. $\tan 25^\circ \tan 31^\circ + \tan 31^\circ \tan 34^\circ + \tan 34^\circ \tan 25^\circ$

Since,$25^\circ + 31^\circ + 34^\circ = 90^\circ$

Using identity, $\tan A\tan B + \tan B\tan C + \tan C\tan A = 1$

when $A + B + C = 90^\circ$

Therefore $= 1$

Answer: A) 1

7. $\frac{\cos 15^\circ + \sin 15^\circ}{\cos 15^\circ - \sin 15^\circ}$

Divide numerator and denominator by $\cos 15^\circ$:

$= \frac{1 + \tan 15^\circ}{1 - \tan 15^\circ}$

Using, $\frac{1+\tan\theta}{1-\tan\theta} = \tan(45^\circ + \theta)$

$= \tan 60^\circ$

$= \sqrt{3}$

Answer: C) $\sqrt{3}$

8. $\sin^2 75^\circ - \sin^2 15^\circ$

Using, $\sin^2 A - \sin^2 B = \sin(A+B)\sin(A-B)$

$= \sin 90^\circ \sin 60^\circ$

$= 1 \cdot \frac{\sqrt{3}}{2}$

$= \frac{\sqrt{3}}{2}$

Answer: A) $\frac{\sqrt{3}}{2}$

ONE OR MORE THAN ONE OPTION CORRECT TYPE

9. For angles of a triangle, $A + B + C = 180^\circ$

Using identity, $\tan A + \tan B + \tan C = \tan A \tan B \tan C$

Therefore, $\tan A + \tan B + \tan C - \tan A \tan B \tan C = 0$

Now, $\cos 90^\circ = 0, \sin 0^\circ = 0$

Correct options:

Answer: B, C

10. Given,

$\cos x + \cos y = 0$

Using,

$\cos x + \cos y = 2\cos\frac{x+y}{2}\cos\frac{x-y}{2}$

$2\cos\frac{x+y}{2}\cos\frac{x-y}{2} = 0$

So,

$\cos\frac{x+y}{2} = 0$

or

$\cos\frac{x-y}{2} = 0$

Using,

$\sin x + \sin y = 2\sin\frac{x+y}{2}\cos\frac{x-y}{2}$

Given condition implies values of

$\cos\left(\frac{x-y}{2}\right)$

can be

$\pm\frac{1}{2}$

Hence possible values are:

Answer: A, B

COMPREHENSION TYPE

Given: $\alpha$and $\beta$are solutions of $a\cos\theta + b\sin\theta = c$

Rewrite as $\frac{a}{\sqrt{a^2+b^2}}\cos\theta + \frac{b}{\sqrt{a^2+b^2}}\sin\theta = \frac{c}{\sqrt{a^2+b^2}}$

Let $a = \sqrt{a^2+b^2}\cos\phi, b = \sqrt{a^2+b^2}\sin\phi$

Then $\cos(\theta-\phi) = \frac{c}{\sqrt{a^2+b^2}}$

So the two solutions are $\alpha = \phi+\gamma, \beta = \phi-\gamma$

Where $\cos\gamma = \frac{c}{\sqrt{a^2+b^2}}$

Hence $\cos(\alpha-\beta) = \cos 2\gamma$

and $\cos 2\gamma = 2\cos^2\gamma - 1 = 2\left(\frac{c^2}{a^2+b^2}\right) - 1 = \frac{2c^2-a^2-b^2}{a^2+b^2}$

Also, $\cos(\alpha-\beta) = \cos\alpha\cos\beta + \sin\alpha\sin\beta$

Now using symmetry, $\sin\alpha\sin\beta = \frac{c^2-a^2}{a^2+b^2}$

and $\cos\alpha\cos\beta = \frac{c^2-b^2}{a^2+b^2}$

Therefore:

11.

$$\sin\alpha\sin\beta = \frac{c^2-a^2}{a^2+b^2}$$

Answer: D

$$\boxed{\frac{c^2-a^2}{a^2+b^2}}$$

12.

$$\cos\alpha\cos\beta = \frac{c^2-b^2}{a^2+b^2}$$

Answer: A

$$\boxed{\frac{c^2-b^2}{a^2+b^2}}$$

MATRIX MATCHING TYPE

a)

$\cos 20^\circ + \cos 140^\circ + \cos 100^\circ$

$\cos 140^\circ = -\cos 40^\circ$

$\cos 100^\circ = -\cos 80^\circ$

$\cos 20^\circ - \cos 80^\circ = -2\sin 50^\circ \sin(-30^\circ) = \sin 50^\circ = \cos 40^\circ$

Therefore,

$\cos 40^\circ - \cos 40^\circ = 0$

a → P

b)

$(1+\tan 20^\circ)(1+\tan 25^\circ)$

$= 1 + \tan 20^\circ + \tan 25^\circ + \tan 20^\circ \tan 25^\circ$

Using

$\tan 20^\circ + \tan 25^\circ + \tan 20^\circ \tan 25^\circ = 1$

$= 1 + 1 = 2$

b → Q

c)

$\tan 20°\tan 32° + \tan 32°\tan 38° + \tan 38°\tan 20°$

Since

$20° + 32° + 38° = 90°$

Using identity

$\tan A\tan B + \tan B\tan C + \tan C\tan A = 1$

$= 1$

c → R

d)

$\sin^2 10° + \sin^2 110° + \sin^2 130°$

$\sin 110° = \sin 70° = \cos 20°$

$\sin 130° = \sin 50° = \cos 40°$

$= \sin^2 10° + \cos^2 20° + \cos^2 40°$

Using standard identity,

$\sin^2 10° + \sin^2 50° + \sin^2 70° = \frac{3}{2}$

$= \frac{3}{2}$

d → S

Final Matching:

a → P

b → Q

c → R

d → S

INTEGER TYPE

14. Given $\alpha + \beta + \gamma = \frac{\pi}{2}$

And $\cot\alpha, \cot\beta, \cot\gamma$

are in A.P.

So, $2\cot\beta = \cot\alpha + \cot\gamma$

Now, $\alpha + \gamma = \frac{\pi}{2} - \beta$

Taking cotangent, $\cot(\alpha + \gamma) = \tan\beta$

Using formula, $\frac{\cot\alpha\cot\gamma - 1}{\cot\alpha + \cot\gamma} = \tan\beta$

Since $\cot\alpha + \cot\gamma = 2\cot\beta$

$\frac{\cot\alpha\cot\gamma - 1}{2\cot\beta} = \frac{1}{\cot\beta}$

$\cot\alpha\cot\gamma - 1 = 2$

$\boxed{\cot\alpha\cot\gamma = 3}$

15. $\cos\theta + \cos(240° + \theta) + \cos(240° - \theta)$

Using $\cos(A + B) + \cos(A - B) = 2\cos A\cos B$

$= \cos\theta + 2\cos 240°\cos\theta$

$= \cos\theta + 2\left(-\frac{1}{2}\right)\cos\theta$

$= \cos\theta - \cos\theta$

$\boxed{0}$

MULTIPLE AND SUBMULTIPLE ANGLES – TRANSFORMATIONS

LEVEL-I

1. $\frac{\sin 2\theta}{1-\cos 2\theta}$

Using $\sin 2\theta = 2\sin\theta\cos\theta$

And $1 - \cos 2\theta = 2\sin^2\theta$

$= \frac{2\sin\theta\cos\theta}{2\sin^2\theta}$

$= \frac{\cos\theta}{\sin\theta}$

$= \cot\theta$

Answer: D

2. $\sqrt{2+\sqrt{2+2\cos 2\theta}}$

Using $1 + \cos 2\theta = 2\cos^2\theta$

$2 + 2\cos 2\theta = 2(1 + \cos 2\theta) = 4\cos^2\theta$

$\sqrt{2 + 2\cos 2\theta} = 2\cos\theta$

So, $\sqrt{2 + 2\cos\theta}$

Using $1 + \cos\theta = 2\cos^2\frac{\theta}{2}$

$2 + 2\cos\theta = 4\cos^2\frac{\theta}{2}$

$= \sqrt{4\cos^2\frac{\theta}{2}}$

$= 2\cos\frac{\theta}{2}$

Answer: D

3. Given $\cos\alpha = \frac{15}{17}$

Using $\cos 2\alpha = 2\cos^2\alpha - 1$

$= 2\left(\frac{15}{17}\right)^2 - 1$

$= 2 \cdot \frac{225}{289} - 1$

$= \frac{450 - 289}{289}$

$= \frac{161}{289}$

Answer: C

4. $\frac{\cos 3\theta}{2\cos 2\theta - 1}$

Using $\cos 3\theta = 4\cos^3\theta - 3\cos\theta$

and $2\cos 2\theta - 1 = 2(2\cos^2\theta - 1) - 1$

$= 4\cos^2\theta - 3$

Thus, $\frac{\cos\theta(4\cos^2\theta - 3)}{4\cos^2\theta - 3}$

$= \cos\theta$

Answer: A

5. Given $\sin\frac{A}{2} = \frac{3}{5}$

$\cos\frac{A}{2} = \frac{4}{5}$

$\sin A = 2\sin\frac{A}{2}\cos\frac{A}{2}$

$= 2 \cdot \frac{3}{5} \cdot \frac{4}{5} = \frac{24}{25}$

$\cos A = 1 - 2\sin^2\frac{A}{2}$

$= 1 - 2\left(\frac{3}{5}\right)^2$

$= 1 - \frac{18}{25} = \frac{7}{25}$

Using

$\sin 3A = 3\sin A - 4\sin^3 A$

$= 3 \cdot \frac{24}{25} - 4\left(\frac{24}{25}\right)^3$

$= \frac{72}{25} - \frac{55296}{15625}$

$= \frac{45000 - 55296}{15625}$

$= -\frac{10296}{15625} = -\frac{117}{125}$

Answer: B

6. $\operatorname{cosec} 2A + \cot 2A$

$= \frac{1}{\sin 2A} + \frac{\cos 2A}{\sin 2A}$

$= \frac{1 + \cos 2A}{\sin 2A}$

Using $1 + \cos 2A = 2\cos^2 A, \sin 2A = 2\sin A\cos A$

$= \frac{2\cos^2 A}{2\sin A\cos A}$

$= \frac{\cos A}{\sin A} = \cot A$

Answer: A

7. $\frac{\sin 3A}{1+2\cos 2A}$

Using $\sin 3A = 3\sin A - 4\sin^3 A$

And $1 + 2\cos 2A = 1 + 2(1 - 2\sin^2 A)$

$= 3 - 4\sin^2 A$

$\frac{\sin A(3 - 4\sin^2 A)}{3 - 4\sin^2 A} = \sin A$

Answer: D

8. $\left(\frac{\sin A}{1-\cos A}\right) - \left(\frac{1-\cos A}{\sin A}\right)$

Taking LCM,

$= \frac{\sin^2 A - (1 - \cos A)^2}{\sin A(1 - \cos A)}$

$= \frac{\sin^2 A - (1 - 2\cos A + \cos^2 A)}{\sin A(1 - \cos A)}$

$= \frac{\sin^2 A - \cos^2 A - 1 + 2\cos A}{\sin A(1 - \cos A)}$

Using

$\sin^2 A + \cos^2 A = 1$

$= \frac{2\cos A(1-\cos A)}{\sin A(1-\cos A)}$

$= \frac{2\cos A}{\sin A}$

$= 2\cot A$

Answer: B

9. Given $\frac{\cos\alpha}{a} = \frac{\sin\alpha}{b}$

$b\cos\alpha = a\sin\alpha$

$\tan\alpha = \frac{b}{a}$

Take triangle sides: $\sin\alpha = \frac{b}{\sqrt{a^2+b^2}}, \cos\alpha = \frac{a}{\sqrt{a^2+b^2}}$

Now, $a\cos 2\alpha + b\sin 2\alpha$

Using $\cos 2\alpha = \frac{a^2-b^2}{a^2+b^2}$

$\sin 2\alpha = \frac{2ab}{a^2+b^2}$

$= \frac{a(a^2-b^2)+2ab^2}{a^2+b^2}$

$= \frac{a^3+ab^2}{a^2+b^2}$

$= a$

Answer: A

10. $(\sin 3A + \sin A)\sin A + (\cos 3A - \cos A)\cos A$

$= \sin 3A\sin A + \sin^2 A + \cos 3A\cos A - \cos^2 A$

Using $\cos(x-y) = \cos x\cos y + \sin x\sin y$

$= \cos(3A - A) + (\sin^2 A - \cos^2 A)$

$= \cos 2A - \cos 2A$

$= 0$

Answer: A

11. $3\sin 6° - 4\cos^3 84°$

Since $\cos 84° = \sin 6°$

Let $x = \sin 6°$

Then, $3x - 4x^3$

Using identity: $\sin 3\theta = 3\sin\theta - 4\sin^3\theta$

So, $3x - 4x^3 = \sin 18°$

And, $\sin 18° = \frac{\sqrt{5}-1}{4}$

Therefore, $3\sin 6° - 4\cos^3 84° = \frac{\sqrt{5}-1}{4}$

Answer: A) $\frac{\sqrt{5}-1}{4}$

12. Find: $\sin 22\frac{1}{2}°$

Using half-angle formula:$\sin\frac{\theta}{2} = \sqrt{\frac{1-\cos\theta}{2}}$

Take $\theta = 45°$:$\sin 22.5° = \sqrt{\frac{1-\cos 45°}{2}}$

Since $\cos 45° = \frac{1}{\sqrt{2}}$

$$= \sqrt{\frac{1 - \frac{1}{\sqrt{2}}}{2}}$$

$$= \sqrt{\frac{\sqrt{2} - 1}{2\sqrt{2}}}$$

Answer: B) $\sqrt{\frac{\sqrt{2}-1}{2\sqrt{2}}}$

13. $180° < \theta < 270°, \cos\theta = -\frac{3}{5}$

θ is in III quadrant, so:

$\sin\theta = -\frac{4}{5}$

Using:

$$\tan\frac{\theta}{2} = \frac{\sin\theta}{1 + \cos\theta}$$

$$\tan\frac{\theta}{2} = \frac{\sin\theta}{1 + \cos\theta}$$

$$\tan\frac{\theta}{2} = \frac{-\frac{4}{5}}{1 - \frac{3}{5}} = \frac{-\frac{4}{5}}{\frac{2}{5}} = -2$$

Answer: B) -2

14. $2\cos\theta = x + \frac{1}{x}$

Using identity:

$$\left(x + \frac{1}{x}\right)^3 = x^3 + \frac{1}{x^3} + 3\left(x + \frac{1}{x}\right)$$

$$\left(2\cos\theta\right)^3 = x^3 + \frac{1}{x^3} + 3(2\cos\theta\Big)$$

$$8\cos^3\theta = x^3 + \frac{1}{x^3} + 6\cos\theta$$

$$x^3 + \frac{1}{x^3} = 8\cos^3\theta - 6\cos\theta$$

Using:

$\cos 3\theta = 4\cos^3\theta - 3\cos\theta$

$2\cos 3\theta = 8\cos^3\theta - 6\cos\theta$

Hence,

$$x^3 + \frac{1}{x^3} = 2\cos 3\theta$$

Answer: B) $2\cos 3\theta$

15. $(\cos\alpha + \cos\beta)^2 + (\sin\alpha + \sin\beta)^2$

Expand:

$= \cos^2\alpha + \cos^2\beta + 2\cos\alpha\cos\beta$

$+\sin^2\alpha + \sin^2\beta + 2\sin\alpha\sin\beta$

$= (\cos^2\alpha + \sin^2\alpha) + (\cos^2\beta + \sin^2\beta)$

$+2(\cos\alpha\cos\beta + \sin\alpha\sin\beta)$

$= 1 + 1 + 2\cos(\alpha - \beta)$

$= 2 + 2\cos(\alpha - \beta)$

Using:

$1 + \cos x = 2\cos^2 \frac{x}{2}$

$= 4\cos^2 \frac{\alpha - \beta}{2}$

Answer: A) $4\cos^2\left(\frac{\alpha-\beta}{2}\right)$

16. $\sin\alpha\sin(60-\alpha)\sin(60+\alpha)$

Using:

$\sin(A-B)\sin(A+B) = \sin^2 A - \sin^2 B$

$= \sin\alpha(\sin^2 60° - \sin^2\alpha)$

$= \sin\alpha\left(\frac{3}{4} - \sin^2\alpha\right)$

$= \frac{3}{4}\sin\alpha - \sin^3\alpha$

Using:

$\sin 3\alpha = 3\sin\alpha - 4\sin^3\alpha$

$\frac{1}{4}\sin 3\alpha = \frac{3}{4}\sin\alpha - \sin^3\alpha$

Therefore,

$\sin\alpha\sin(60-\alpha)\sin(60+\alpha) = \frac{1}{4}\sin 3\alpha$

Answer: C) $\frac{1}{4}\sin 3\alpha$

17. $\sin 20°\sin 40°\sin 60°\sin 80°$

Use identity: $\sin 20°\sin 40°\sin 80° = \frac{\sqrt{3}}{8}$

Also, $\sin 60° = \frac{\sqrt{3}}{2}$

Therefore, $\sin 20°\sin 40°\sin 60°\sin 80° = \frac{\sqrt{3}}{8}\cdot\frac{\sqrt{3}}{2} = \frac{3}{16}$

Answer: B) $\frac{3}{16}$

18. $\cos 20°\cos 40°\cos 60°\cos 80°$

Using identity: $\cos 20°\cos 40°\cos 80° = \frac{1}{8}$

And, $\cos 60° = \frac{1}{2}$

Hence, $\cos 20°\cos 40°\cos 60°\cos 80° = \frac{1}{8}\cdot\frac{1}{2} = \frac{1}{16}$

Answer: D) $\frac{1}{16}$

19. $\frac{\sin x}{\sin\frac{x}{8}}$

Use: $\sin x = 2\sin\frac{x}{2}\cos\frac{x}{2}$

Again, $\sin\frac{x}{2} = 2\sin\frac{x}{4}\cos\frac{x}{4}$

And, $\sin\frac{x}{4} = 2\sin\frac{x}{8}\cos\frac{x}{8}$

Substituting, $\sin x = 2\left(2\sin\frac{x}{4}\cos\frac{x}{4}\right)\cos\frac{x}{2}$

$= 4\sin\frac{x}{4}\cos\frac{x}{4}\cos\frac{x}{2}$

$= 4\left(2\sin\frac{x}{8}\cos\frac{x}{8}\right)\cos\frac{x}{4}\cos\frac{x}{2}$

$= 8\sin\frac{x}{8}\cos\frac{x}{8}\cos\frac{x}{4}\cos\frac{x}{2}$

Therefore, $\frac{\sin x}{\sin\frac{x}{8}} = 8\cos\frac{x}{8}\cos\frac{x}{4}\cos\frac{x}{2}$

Answer: A) $8\cos\frac{x}{8}\cos\frac{x}{4}\cos\frac{x}{2}$

20. Given:

$x \in (45°, 90°), \sin 2x = \frac{1}{2}$

Since $2x \in (90°, 180°)$,

$2x = 150°$

$x = 75°$

Now,

$\sin^3 x - \frac{3}{4}\sin x$

Using identity:

$\sin 3x = 3\sin x - 4\sin^3 x$

$4\sin^3 x - 3\sin x = -\sin 3x$

$\sin^3 x - \frac{3}{4}\sin x = -\frac{1}{4}\sin 3x$

For $x = 75°$,

$3x = 225°$

$\sin 225° = -\frac{1}{\sqrt{2}}$

Hence,

$-\frac{1}{4}\left(-\frac{1}{\sqrt{2}}\right) = \frac{1}{4\sqrt{2}}$

Answer: C) $\frac{1}{4\sqrt{2}}$

LEVEL-II

1. $\cos^4\theta - \sin^4\theta$

$= (\cos^2\theta - \sin^2\theta)(\cos^2\theta + \sin^2\theta)$

$= \cos 2\theta$

Answer: A) $\cos 2\theta$

2. Given: $\tan\theta = \frac{b}{a}$

Then, $\sin\theta = \frac{b}{\sqrt{a^2+b^2}}, \cos\theta = \frac{a}{\sqrt{a^2+b^2}}$

$\cos 2\theta = \frac{a^2 - b^2}{a^2 + b^2}$

$\sin 2\theta = \frac{2ab}{a^2 + b^2}$

Now, $a\cos 2\theta + b\sin 2\theta$

$= a\left(\frac{a^2 - b^2}{a^2 + b^2}\right) + b\left(\frac{2ab}{a^2 + b^2}\right)$

$= \frac{a^3 - ab^2 + 2ab^2}{a^2 + b^2}$

$= \frac{a(a^2 + b^2)}{a^2 + b^2} = a$

Answer: D) a

3. $8\cos^3 20° - 6\cos 20°$

Using identity: $\cos 3\theta = 4\cos^3\theta - 3\cos\theta$

Multiply by 2: $8\cos^3\theta - 6\cos\theta = 2\cos 3\theta$

For $\theta = 20°$,

$= 2\cos 60° = 2 \cdot \frac{1}{2} = 1$

Answer: D) 1

4. $6\sin 15° - 8\sin^3 15°$

Using: $\sin 3\theta = 3\sin\theta - 4\sin^3\theta$

$6\sin\theta - 8\sin^3\theta = 2(3\sin\theta - 4\sin^3\theta)$

$= 2\sin 3\theta$

For $\theta = 15°$, $= 2\sin 45° = 2 \cdot \frac{1}{\sqrt{2}} = \sqrt{2}$

Answer: B) $\sqrt{2}$

5. $\frac{1-\cos 2A}{1+\cos 2A}$

Using identities: $1 - \cos 2A = 2\sin^2 A$

And $1 + \cos 2A = 2\cos^2 A$

$= \frac{2\sin^2 A}{2\cos^2 A}$

$= \tan^2 A$

Answer: B) $\tan^2 A$

6. $4\sin A\cos^3 A - 4\cos A\sin^3 A$

Take $4\sin A\cos A$common $= 4\sin A\cos A(\cos^2 A - \sin^2 A)$

Using: $2\sin A\cos A = \sin 2A$

And $\cos^2 A - \sin^2 A = \cos 2A$

$= 2\sin 2A\cos 2A$

Using:

$2\sin x\cos x = \sin 2x$

$= \sin 4A$

Answer: A) $\sin 4A$

7. $\sin\frac{\pi}{16}\sin\frac{3\pi}{16}\sin\frac{5\pi}{16}\sin\frac{7\pi}{16}$

Use identity:

$\sin x\sin\left(\frac{\pi}{2} - x\right) = \sin x\cos x = \frac{1}{2}\sin 2x$

Pair terms:

$\sin\frac{\pi}{16}\sin\frac{7\pi}{16} = \sin\frac{\pi}{16}\cos\frac{\pi}{16} = \frac{1}{2}\sin\frac{\pi}{8}$

$\sin\frac{3\pi}{16}\sin\frac{5\pi}{16} = \sin\frac{3\pi}{16}\cos\frac{3\pi}{16} = \frac{1}{2}\sin\frac{3\pi}{8}$

Therefore,

$= \frac{1}{4}\sin\frac{\pi}{8}\sin\frac{3\pi}{8}$

Again,

$\sin\frac{3\pi}{8} = \cos\frac{\pi}{8}$

$= \frac{1}{4}\sin\frac{\pi}{8}\cos\frac{\pi}{8} = \frac{1}{8}\sin\frac{\pi}{4}$

$= \frac{1}{8} \cdot \frac{1}{\sqrt{2}} = \frac{\sqrt{2}}{16}$

Answer: B) $\frac{\sqrt{2}}{16}$

8. $\cot A - \tan A$

$= \frac{\cos A}{\sin A} - \frac{\sin A}{\cos A}$

$= \frac{\cos^2 A - \sin^2 A}{\sin A\cos A}$

$$= \frac{\cos 2A}{\frac{1}{2}\sin 2A}$$

$$= 2\cot 2A$$

Answer: C) $2\cot 2A$

9. $\cos\frac{\pi}{7}\cos\frac{2\pi}{7}\cos\frac{3\pi}{7}$

Known identity:

$$\cos\frac{\pi}{7}\cos\frac{2\pi}{7}\cos\frac{3\pi}{7} = \frac{1}{8}$$

Answer: A) $\frac{1}{8}$

10. $\sqrt{2-2\cos\theta}$

Using identity:

$$1-\cos\theta = 2\sin^2\frac{\theta}{2}$$

$$2-2\cos\theta = 2(1-\cos\theta) = 4\sin^2\frac{\theta}{2}$$

$$\sqrt{2-2\cos\theta} = \sqrt{4\sin^2\frac{\theta}{2}} = 2\sin\frac{\theta}{2}$$

Answer: A) $2\sin\frac{\theta}{2}$

ONE OR MORE THAN ONE OPTION CORRECT TYPE

11. $\frac{1+\cos\theta-\sin\theta}{-1+\cos\theta+\sin\theta}$

Rewrite numerator and denominator using half-angle formulas:

$$1+\cos\theta = 2\cos^2\frac{\theta}{2}$$

$$\sin\theta = 2\sin\frac{\theta}{2}\cos\frac{\theta}{2}$$

Numerator:

$$= 2\cos\frac{\theta}{2}\left(\cos\frac{\theta}{2}-\sin\frac{\theta}{2}\right)$$

Denominator:

$$-1+\cos\theta+\sin\theta$$

$$= -(1-\cos\theta)+\sin\theta$$

$$= -2\sin^2\frac{\theta}{2}+2\sin\frac{\theta}{2}\cos\frac{\theta}{2}$$

$$= 2\sin\frac{\theta}{2}\left(\cos\frac{\theta}{2}-\sin\frac{\theta}{2}\right)$$

Therefore,

$$= \frac{2\cos\frac{\theta}{2}(\cos\frac{\theta}{2}-\sin\frac{\theta}{2})}{2\sin\frac{\theta}{2}(\cos\frac{\theta}{2}-\sin\frac{\theta}{2})}$$

$$= \cot\frac{\theta}{2}$$

Also,

$$\frac{\cos\frac{\theta}{2}}{\sin\frac{\theta}{2}} = \cot\frac{\theta}{2}$$

And,

$$\frac{\operatorname{cosec}\frac{\theta}{2}}{\sec\frac{\theta}{2}} = \frac{1/\sin\frac{\theta}{2}}{1/\cos\frac{\theta}{2}} = \cot\frac{\theta}{2}$$

Correct options: A, C and D

12. $\frac{\sin 2\theta}{1+\cos 2\theta}$

Using identities:

$\sin 2\theta = 2\sin\theta\cos\theta$

θ

θ

$\sin(\theta+\theta) \approx 0.866$

$\sin(\theta)\cos(\theta) + \cos(\theta)\sin(\theta) \approx 0.866$

and

$1 + \cos 2\theta = 2\cos^2\theta$

$= \frac{2\sin\theta\cos\theta}{2\cos^2\theta}$

$= \tan\theta$

Also,

$\sin\theta\sec\theta = \frac{\sin\theta}{\cos\theta} = \tan\theta$

Correct options: C and D

13.

$\sqrt{2-2\cos\theta}$

Using identity:

$1 - \cos\theta = 2\sin^2\frac{\theta}{2}$

$2 - 2\cos\theta = 2(1-\cos\theta) = 4\sin^2\frac{\theta}{2}$

$\sqrt{2-2\cos\theta} = \sqrt{4\sin^2\frac{\theta}{2}} = 2\sin\frac{\theta}{2}$

Correct option: B) $2\sin\frac{\theta}{2}$

COMPREHENSION TYPE

Given: $\sin^3\theta + \sin^3(60^\circ-\theta) + \sin^3(240^\circ+\theta) = -\frac{3}{4}\sin 3\theta$

and $\cos^3\theta + \cos^3(120^\circ+\theta) + \cos^3(240^\circ+\theta) = -\frac{3}{4}\cos 3\theta$

14. $\sin^3 10^\circ + \sin^3 50^\circ - \sin^3 70^\circ$

Observe: $\sin 70^\circ = \sin(240^\circ + 10^\circ)$

Taking $\theta = 10^\circ$, $\sin^3 10^\circ + \sin^3 50^\circ + \sin^3 250^\circ = -\frac{3}{4}\sin 30^\circ$

Since

$\sin 250^\circ = -\sin 70^\circ$

$\sin^3 250^\circ = -\sin^3 70^\circ$

Therefore, $\sin^3 10^\circ + \sin^3 50^\circ - \sin^3 70^\circ = -\frac{3}{4}\cdot\frac{1}{2} = -\frac{3}{8}$

Answer: C) $-\frac{3}{8}$

15. $20^\circ + \cos^3 140^\circ + \cos^3 260^\circ$

Take $\theta = 20^\circ$:

$\cos^3 20^\circ + \cos^3(120^\circ+20^\circ) + \cos^3(240^\circ+20^\circ) = -\frac{3}{4}\cos 60^\circ$

$= -\frac{3}{4} \cdot \frac{1}{2} = -\frac{3}{8}$

Answer: C) $-\frac{3}{8}$

16. $\cos^3 10° + \cos^3 130° + \cos^3 250°$

Take $\theta = 10°$:

$\cos^3 10° + \cos^3(120° + 10°) + \cos^3(240° + 10°) = -\frac{3}{4}\cos 30°$

$= -\frac{3}{4} \cdot \frac{\sqrt{3}}{2} = -\frac{3\sqrt{3}}{8}$

Answer: C) $-\frac{3\sqrt{3}}{8}$

17. MATRIX MATCHING TYPE

Given: $\cos\theta + \sin\theta = x$

$\cos\theta - \sin\theta = y$

Add: $2\cos\theta = x + y$

$\cos\theta = \frac{x+y}{2}$

Subtract: $2\sin\theta = x - y$

$\sin\theta = \frac{x-y}{2}$

Also, $x^2 = (\cos\theta + \sin\theta)^2 = 1 + 2\sin\theta\cos\theta$

$x^2 = 1 + \sin 2\theta$

Hence, $\sin 2\theta = x^2 - 1$

So, $b \to T$

Now, $y^2 = (\cos\theta - \sin\theta)^2 = 1 - 2\sin\theta\cos\theta$

$y^2 = 1 - \sin 2\theta$

$\sin 2\theta = 1 - y^2$

So, $b \to P$

Now, $\cos 2\theta = (\cos\theta + \sin\theta)(\cos\theta - \sin\theta) = xy$

So, $a \to Q$

Next,

$\tan\theta = \frac{\sin\theta}{\cos\theta} = \frac{x-y}{x+y}$

$\cot\theta = \frac{x+y}{x-y}$

So, $c \to S$

Finally, $\tan 2\theta = \frac{\sin 2\theta}{\cos 2\theta} = \frac{x^2-1}{xy}$

So, $d \to R$

Matching:

a → Q

b → P

c → S

d → R

18. Given: $\cot\alpha - \tan\alpha = 2\cot 2\alpha$

Expression: $\tan\alpha - \cot\alpha + 2\tan 2\alpha + 4\tan 4\alpha + 8\tan 8\alpha$

Since $\tan\alpha - \cot\alpha = -2\cot 2\alpha$

Expression becomes: $-2\cot 2\alpha + 2\tan 2\alpha + 4\tan 4\alpha + 8\tan 8\alpha$

Using:

$\tan\theta - \cot\theta = -2\cot 2\theta$

$2(\tan 2\alpha - \cot 2\alpha) = 2(-2\cot 4\alpha) = -4\cot 4\alpha$
Now expression: $-4\cot 4\alpha + 4\tan 4\alpha + 8\tan 8\alpha$
Again, $4(\tan 4\alpha - \cot 4\alpha) = -8\cot 8\alpha$
Expression:
$-8\cot 8\alpha + 8\tan 8\alpha$
$= 8(\tan 8\alpha - \cot 8\alpha)$
$= 8(-2\cot 16\alpha)$
$= -16\cot 16\alpha$
Answer: $-16\cot 16\alpha$

19. $\frac{\cos^3\theta - \cos 3\theta}{\cos\theta} + \frac{\sin^3\theta + \sin 3\theta}{\sin\theta}$
Using: $\cos 3\theta = 4\cos^3\theta - 3\cos\theta$
$\cos^3\theta - \cos 3\theta = \cos^3\theta - (4\cos^3\theta - 3\cos\theta)$
$= 3\cos\theta - 3\cos^3\theta$
$= 3\cos\theta(1 - \cos^2\theta)$
$= 3\cos\theta\sin^2\theta$
Dividing by $\cos\theta$: $= 3\sin^2\theta$
Now, $\sin 3\theta = 3\sin\theta - 4\sin^3\theta$
$\sin^3\theta + \sin 3\theta = \sin^3\theta + 3\sin\theta - 4\sin^3\theta$
$= 3\sin\theta - 3\sin^3\theta$
$= 3\sin\theta(1 - \sin^2\theta)$
$= 3\sin\theta\cos^2\theta$
Dividing by $\sin\theta$: $= 3\cos^2\theta$
Therefore, $3\sin^2\theta + 3\cos^2\theta = 3$
Answer: 3

20. $\frac{\sin 3A}{\sin A} - \frac{\cos 3A}{\cos A}$
Using:
$\sin 3A = 3\sin A - 4\sin^3 A$
$\frac{\sin 3A}{\sin A} = 3 - 4\sin^2 A$
Using:
$\cos 3A = 4\cos^3 A - 3\cos A$
$\frac{\cos 3A}{\cos A} = 4\cos^2 A - 3$
Therefore,
$(3 - 4\sin^2 A) - (4\cos^2 A - 3)$
$= 6 - 4(\sin^2 A + \cos^2 A)$
$= 6 - 4 = 2$
Answer: 2

Student practice corner

SINGLE OPTION CORRECT TYPE

1.

$\frac{\sin 2\theta}{1 + \cos 2\theta}$
Using identities:
$\sin 2\theta = 2\sin\theta\cos\theta$
and
$1 + \cos 2\theta = 2\cos^2\theta$

$= \frac{2\sin\theta\cos\theta}{2\cos^2\theta}$

$= \tan\theta$

Answer: C) $\tan\theta$

2.

$\frac{3\sin\theta - \sin 3\theta}{3\cos\theta + \cos 3\theta}$

Using:

$\sin 3\theta = 3\sin\theta - 4\sin^3\theta$

$3\sin\theta - \sin 3\theta = 4\sin^3\theta$

Using:

$\cos 3\theta = 4\cos^3\theta - 3\cos\theta$

$3\cos\theta + \cos 3\theta = 4\cos^3\theta$

Therefore,

$= \frac{4\sin^3\theta}{4\cos^3\theta} = \tan^3\theta$

Answer: C) $\tan^3\theta$

3.

$\cos\alpha\cos(60-\alpha)\cos(60+\alpha)$

Using:

$\cos(A-B)\cos(A+B) = \cos^2 A - \sin^2 B$

$= \cos\alpha(\cos^2 60° - \sin^2\alpha)$

$= \cos\alpha\left(\frac{1}{4} - \sin^2\alpha\right)$

$= \cos\alpha\left(\cos^2\alpha - \frac{3}{4}\right)$

$= \frac{1}{4}(4\cos^3\alpha - 3\cos\alpha)$

Using:

$\cos 3\alpha = 4\cos^3\alpha - 3\cos\alpha$

$= \frac{1}{4}\cos 3\alpha$

Answer: D) $\frac{1}{4}\cos 3\alpha$

4.

$\sin 20°\sin 40°\sin 100°$

Since

$\sin 100° = \sin 80°$

$= \sin 20°\sin 40°\sin 80°$

Known identity:

$\sin 20°\sin 40°\sin 80° = \frac{\sqrt{3}}{8}$

Answer: C) $\frac{\sqrt{3}}{8}$

5.

$\frac{\sin A}{1+\cos A} + \frac{\sin A}{1-\cos A}$

Take LCM:

$= \frac{\sin A(1-\cos A) + \sin A(1+\cos A)}{1-\cos^2 A}$

$= \frac{\sin A - \sin A\cos A + \sin A + \sin A\cos A}{\sin^2 A}$

$= \frac{2\sin A}{\sin^2 A}$

$= \frac{2}{\sin A} = 2\operatorname{cosec} A$

Answer: B) $2\operatorname{cosec} A$

Using identities:

$\sin 2x = 2\sin x\cos x$

and

$1 + \cos 2x = 2\cos^2 x$

Let $x = 2\theta$:

$\frac{\sin 4\theta}{1 + \cos 4\theta} = \frac{2\sin 2\theta\cos 2\theta}{2\cos^2 2\theta}$

$= \tan 2\theta$

Answer: A) $\tan 2\theta$

7.

$\sqrt{2 + \sqrt{2 + 2\cos 4\theta}}$

Using:

$1 + \cos 2x = 2\cos^2 x$

$2 + 2\cos 4\theta = 4\cos^2 2\theta$

$\sqrt{2 + 2\cos 4\theta} = 2\cos 2\theta$

Therefore,

$\sqrt{2 + 2\cos 2\theta}$

Again,

$2 + 2\cos 2\theta = 4\cos^2 \theta$

$\sqrt{4\cos^2 \theta} = 2\cos \theta$

Answer: B) $2\cos \theta$

8. Given:

$\cos \alpha = \frac{8}{17}$

Using identity:

$\cos 2\alpha = 2\cos^2 \alpha - 1$

$\cos 2\alpha = 2\left(\frac{8}{17}\right)^2 - 1$

$= 2 \cdot \frac{64}{289} - 1$

$= \frac{128}{289} - \frac{289}{289}$

$= -\frac{161}{289}$

Answer: A) $-\frac{161}{289}$

ONE OR MORE THAN ONE OPTION CORRECT TYPE

9.

$\frac{\sin 2\theta}{1 - \cos 2\theta}$

Using:

$\sin 2\theta = 2\sin \theta\cos \theta$

and

$1 - \cos 2\theta = 2\sin^2 \theta$

$= \frac{2\sin\theta\cos\theta}{2\sin^2\theta}$

$= \frac{\cos\theta}{\sin\theta} = \cot\theta$

Also,

$\cos\theta\operatorname{cosec}\theta = \frac{\cos\theta}{\sin\theta} = \cot\theta$

Correct options: A and C

10.

$\cot A + \tan A$

$= \frac{\cos A}{\sin A} + \frac{\sin A}{\cos A}$

$= \frac{\cos^2 A + \sin^2 A}{\sin A\cos A}$

$= \frac{1}{\sin A\cos A}$

Since,

$\sin 2A = 2\sin A\cos A$

$= \frac{2}{\sin 2A} = 2\operatorname{cosec} 2A$

Correct option: A) $2\operatorname{cosec} 2A$

MATRIX MATCHING TYPE

a) $\sin 10°\sin 50°\sin 70°$

Known identity:

$\sin 10°\sin 50°\sin 70° = \frac{1}{8}$

$a \to R$

b) $\cos 20°\cos 40°\cos 60°\cos 80°$

$\cos 20°\cos 40°\cos 80° = \frac{1}{8}$

Since

$\cos 60° = \frac{1}{2}$

$= \frac{1}{8} \cdot \frac{1}{2} = \frac{1}{16}$

$b \to Q$

c) $\sin 20°\sin 40°\sin 30°\sin 80°$

Using: $\sin 20°\sin 40°\sin 80° = \frac{\sqrt{3}}{8}$

And $\sin 30° = \frac{1}{2}$

$= \frac{\sqrt{3}}{8} \cdot \frac{1}{2} = \frac{\sqrt{3}}{16}$

$c \to P$

d) $\cos 10°\cos 30°\cos 50°\cos 70°$

Using:

$\cos 50° = \sin 40°, \cos 70° = \sin 20°, \cos 10° = \sin 80°$

$= \sin 20°\sin 40°\sin 80°\cos 30°$

$= \frac{\sqrt{3}}{8} \cdot \frac{\sqrt{3}}{2} = \frac{3}{16}$

$d \to T$

Final Matching:

a → R

b → Q

c → P

d → T

14.

$\sqrt{3}\text{cosec}\,20^\circ - \sec 20^\circ$

$= \dfrac{\sqrt{3}}{\sin 20^\circ} - \dfrac{1}{\cos 20^\circ}$

Take LCM:

$= \dfrac{\sqrt{3}\cos 20^\circ - \sin 20^\circ}{\sin 20^\circ \cos 20^\circ}$

Using:

$\sqrt{3}\cos\theta - \sin\theta = 2\sin(60^\circ - \theta)$

For $\theta = 20^\circ$,

$= \dfrac{2\sin 40^\circ}{\sin 20^\circ \cos 20^\circ}$

Also,

$2\sin 20^\circ \cos 20^\circ = \sin 40^\circ$

$\sin 20^\circ \cos 20^\circ = \dfrac{1}{2}\sin 40^\circ$

Therefore,

$= \dfrac{2\sin 40^\circ}{\frac{1}{2}\sin 40^\circ} = 4$

$\boxed{4}$

15. $\tan 2\alpha - \tan\alpha(1 + \sec 2\alpha)$

Using: $\tan 2\alpha = \frac{2\tan\alpha}{1-\tan^2\alpha}$

And $\sec 2\alpha = \frac{1+\tan^2\alpha}{1-\tan^2\alpha}$

Let $t = \tan\alpha$

Then,

$= \dfrac{2t}{1-t^2} - t\left(1 + \dfrac{1+t^2}{1-t^2}\right)$

$= \dfrac{2t}{1-t^2} - t\left(\dfrac{1-t^2+1+t^2}{1-t^2}\right)$

$= \dfrac{2t}{1-t^2} - \dfrac{2t}{1-t^2} = 0$

$\boxed{0}$

Properties of Triangles

LEVEL-I

1. Given:

$a = \sqrt{3} + 1,\ B = 30^\circ,\ C = 45^\circ$

Using sine rule,

$\dfrac{a}{\sin A} = \dfrac{c}{\sin C}$

$A = 180^\circ - (30^\circ + 45^\circ) = 105^\circ$

$\sin 105° = \sin(60° + 45°) = \frac{\sqrt{6}+\sqrt{2}}{4}$

$\frac{\sqrt{3}+1}{(\sqrt{6}+\sqrt{2})/4} = \frac{c}{1/\sqrt{2}}$

$c = 2$

Answer: A) 2

2. Given:

$b = 20, c = 21, \sin A = \frac{3}{5}$

$\cos A = \frac{4}{5}$

Using cosine rule,

$a^2 = b^2 + c^2 - 2bc\cos A$

$a^2 = 20^2 + 21^2 - 2(20)(21)\left(\frac{4}{5}\right)$

$= 400 + 441 - 672$

$= 169$

$a = 13$

Answer: B) 13

3. Given:

$A = 30°, C = 90°, c = 7\sqrt{3}$

Then

$B = 60°$

In a $30° - 60° - 90°$triangle,

$a : b : c = 1 : \sqrt{3} : 2$

$2k = 7\sqrt{3} \Rightarrow k = \frac{7\sqrt{3}}{2}$

$a = \frac{7\sqrt{3}}{2}$

$b = \frac{21}{2}$

$a + b = \frac{7\sqrt{3}+21}{2}$

Answer: D) $\frac{7\sqrt{3}+21}{2}$

4. Given:

$\sin^2 A + \sin^2 B = \sin^2 C$

Since

$C = 180° - (A + B)$

$\sin C = \sin(A + B)$

Using identities, condition is satisfied when

$C = 90°$

Answer: D) 90°

5. Given:

$A = 45°, B = 75°$

$C = 60°$

By sine rule,

$a : b : c = \sin 45° : \sin 75° : \sin 60°$

Take common factor k,

$a = \frac{k}{\sqrt{2}}$

$c = \frac{\sqrt{3}}{2}k$

Then

$a + c\sqrt{2} = \frac{k}{\sqrt{2}} + \frac{\sqrt{3}}{2}k\sqrt{2}$

$= \frac{k(1+\sqrt{3})}{\sqrt{2}}$

Also,

$b = k\sin 75^\circ = \frac{k(\sqrt{3}+1)}{2\sqrt{2}}$

Hence,

$a + c\sqrt{2} = 2b$

Answer: A) 2b

6. Given:

$B = 30^\circ, c = b$

Equal sides imply equal opposite angles.

$c = b \Rightarrow C = B = 30^\circ$

$A = 180^\circ - 30^\circ - 30^\circ$

$A = 120^\circ$

Answer: D) 120°

7. Angles are $45^\circ$and 60°.

Third angle:

75°

Smallest side is opposite 45°, greatest side opposite 75°.

Ratio:

$\sin 45^\circ : \sin 75^\circ$

$= \frac{1}{\sqrt{2}} : \frac{\sqrt{3}+1}{2\sqrt{2}}$

$= 2 : (\sqrt{3}+1)$

$= \sqrt{3} - 1 : 1$

Answer: A) $\sqrt{3} - 1 : 1$

8. Angles are in ratio $3 : 5 : 10$.

$3x + 5x + 10x = 180^\circ$

$18x = 180^\circ \Rightarrow x = 10^\circ$

Angles:

$30^\circ, 50^\circ, 100^\circ$

Smallest side : greatest side

$= \sin 30^\circ : \sin 100^\circ$

$= \frac{1}{2} : \sin 100^\circ$

$1 : 2\sin 100^\circ$

$\sin 100^\circ = \sin 80^\circ = \cos 10^\circ$

Hence ratio:

$1 : 2\cos 10^\circ$

Answer: D) $1 : 2\cos 10^\circ$

9. In $\triangle ABC$,
$\sum a^3 \sin(B-C)$
$= a^3 \sin(B-C) + b^3 \sin(C-A) + c^3 \sin(A-B)$
Using sine rule:
$a = 2R\sin A, b = 2R\sin B, c = 2R\sin C$
Substituting and simplifying, all terms cancel cyclically.
Hence,
$\sum a^3 \sin(B-C) = 0$
Answer: D) 0

10. Given:
$\sum \frac{b^2-c^2}{a^2} \sin 2A$
Using
$\sin 2A = 2\sin A\cos A$
and cosine rule,
$\cos A = \frac{b^2+c^2-a^2}{2bc}$
After cyclic expansion, terms cancel pairwise.
Therefore,
$\sum \frac{b^2-c^2}{a^2} \sin 2A = 0$
Answer: A) 0

11. In $\triangle ABC$,
$\sum a\cos A$
Using projection formula,
$a = 2R\sin A$
$\sum a\cos A = 2R\sum \sin A\cos A$
$= 2R\left(\frac{\sin 2A + \sin 2B + \sin 2C}{2}\right)$
Using triangle identity,
$\sin 2A + \sin 2B + \sin 2C = 4\sin A\sin B\sin C$
Hence,
$\sum a\cos A = 4R\sin A\sin B\sin C$
Answer: B) $4R\sin A\sin B\sin C$

12. Evaluate:
$\frac{\cos A}{a} + \frac{\cos B}{b} + \frac{\cos C}{c}$
Using cosine rule,
$\cos A = \frac{b^2+c^2-a^2}{2bc}$
So,
$\frac{\cos A}{a} = \frac{b^2+c^2-a^2}{2abc}$
Similarly,
$\frac{\cos B}{b} = \frac{c^2+a^2-b^2}{2abc}$
$\frac{\cos C}{c} = \frac{a^2+b^2-c^2}{2abc}$
Adding,

$= \frac{a^2 + b^2 + c^2}{2abc}$

Answer: A) $\frac{a^2+b^2+c^2}{2abc}$

13. Given:

$a = 26, b = 30, \cos C = \frac{63}{65}$

Using cosine rule,

$c^2 = a^2 + b^2 - 2ab\cos C$

$= 26^2 + 30^2 - 2(26)(30)\left(\frac{63}{65}\right)$

$= 676 + 900 - 1512$

$= 64$

$c = 8$

Answer: D) 8

14. Angles are in A.P.

$2B = A + C$

Since

$A + B + C = 180°$

$3B = 180° \Rightarrow B = 60°$

Using cosine rule,

$b^2 = a^2 + c^2 - 2ac\cos 60°$

$= a^2 + c^2 - ac$

Hence,

$a^2 + c^2 - ac = b^2$

Answer: D) b^2

15. Given:

$2a^2 + 4b^2 + c^2 = 4ab + 2ac$

earrange:

$2a^2 - 4ab + 4b^2 + c^2 - 2ac = 0$

$2(a - b)^2 + (c - a)^2 - a^2 = 0$

Using cosine rule:

$b^2 = a^2 + c^2 - 2ac\cos B$

Substituting and simplifying,

$8\cos B = 7$

$\cos B = \frac{7}{8}$

Answer: C) $\frac{7}{8}$

16. Given**,**

$c^4 - 2(a^2 + b^2)c^2 + a^4 + a^2b^2 + b^4 = 0$

Treat it as a quadratic in c^2.

$c^4 - 2(a^2 + b^2)c^2 + (a^2 + b^2)^2 - a^2b^2 = 0$

$\Rightarrow (c^2 - (a^2 + b^2))^2 = a^2b^2$

$c^2 - (a^2 + b^2) = \pm ab$

Using cosine rule,

$c^2 = a^2 + b^2 - 2ab\cos C$

So,

$-2ab\cos C = \pm ab$

$\cos C = \mp\frac{1}{2}$

Hence,

$C = 60°$

(120°is not among the options)

Answer: C) 60°

17. Evaluate:

$\frac{\cos C + \cos A}{c + a} + \frac{\cos B}{b}$

Using projection formulas,

$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$

$\cos C = \frac{a^2 + b^2 - c^2}{2ab}$

After simplification,

$\frac{\cos C + \cos A}{c + a} = \frac{1}{b} - \frac{\cos B}{b}$

Hence,

$\frac{\cos C + \cos A}{c + a} + \frac{\cos B}{b} = \frac{1}{b}$

Answer: B) $\frac{1}{b}$

18. Given:

$a = 5, b = 4, \cos(A + B) = \frac{31}{32}$

But,

$A + B = 180° - C$

$\cos(A + B) = -\cos C$

Hence,

$\cos C = -\frac{31}{32}$

Using cosine rule,

$c^2 = a^2 + b^2 - 2ab\cos C$

$= 25 + 16 - 2(5)(4)\left(-\frac{31}{32}\right)$

$= 41 + \frac{155}{4}$

$= \frac{319}{4}$

$c = \frac{\sqrt{319}}{2}$

Answer: D) $\frac{\sqrt{319}}{2}$

19. Evaluate:

$\frac{1 + \cos(A - B)\cos C}{1 + \cos(A - C)\cos B}$

Using triangle identities and expansions,

$1 + \cos(A - B)\cos C = \frac{a^2 + b^2}{2ab}$

$1 + \cos(A - C)\cos B = \frac{a^2 + c^2}{2ac}$

Therefore,

$$\frac{1+\cos(A-B)\cos C}{1+\cos(A-C)\cos B}=\frac{a^2+b^2}{a^2+c^2}$$

Answer: B) $\frac{a^2+b^2}{a^2+c^2}$

20. Evaluate:

$c(a\cos B-b\cos A)$

Using projection formulas,

$$a\cos B=\frac{ac^2+ab^2-a^3}{2ac}$$

and

$$b\cos A=\frac{bc^2+ba^2-b^3}{2bc}$$

Simplifying,

$c(a\cos B-b\cos A)=a^2-b^2$

Answer: A) a^2-b^2

LEVEL-II

SINGLE OPTION CORRECT TYPE

1. Given,

$$\frac{b^2+c^2}{b^2-c^2}=\frac{\sin(B+C)}{\sin(B-C)}$$

Since in a triangle,

$B+C=\pi-A$

$\sin(B+C)=\sin A$

Also,

$\sin(B-C)=\sin B\cos C-\cos B\sin C$

Using sine rule:

$b=2R\sin B, c=2R\sin C$

Then,

$$\frac{b^2+c^2}{b^2-c^2}=\frac{\sin^2 B+\sin^2 C}{\sin^2 B-\sin^2 C}$$

After simplification,

$\sin^2 A=\sin^2(B-C)$

This gives either

$A=90°$

or

$B=C$

Hence triangle is either right angled or isosceles.

Answer: C) right angled or isosceles

2. Given,

$$\frac{1}{a+c}+\frac{1}{b+c}=\frac{3}{a+b+c}$$

Cross multiplying,

$(a+b+2c)(a+b+c)=3(a+c)(b+c)$

Expanding,

$a^2+b^2-ab-c^2=0$

Using cosine rule,

$c^2=a^2+b^2-2ab\cos C$

Substitute:

$a^2 + b^2 - ab - (a^2 + b^2 - 2ab\cos C) = 0$

$-ab + 2ab\cos C = 0$

$2\cos C = 1$

$\cos C = \frac{1}{2}$

$C = 60°$

Answer: B) 60°

3. Given:

Perimeter is six times the arithmetic mean of the sine angles and side $a = 1$.

Arithmetic mean of sines:

$\frac{\sin A + \sin B + \sin C}{3}$

So,

$a + b + c = 6 \cdot \frac{\sin A + \sin B + \sin C}{3}$

$a + b + c = 2(\sin A + \sin B + \sin C)$

Using sine rule,

$a = 2R\sin A, b = 2R\sin B, c = 2R\sin C$

Hence,

$2R(\sin A + \sin B + \sin C) = 2(\sin A + \sin B + \sin C)$

$R = 1$

Given $a = 1$,

$1 = 2R\sin A$

$1 = 2\sin A$

$\sin A = \frac{1}{2}$

$A = \frac{\pi}{6}$

Answer: A) $\frac{\pi}{6}$

4. Given,

$\tan A + \tan B + \tan C = 3\sqrt{3}$

In a triangle,

$\tan A + \tan B + \tan C = \tan A\tan B\tan C$

Hence,

$\tan A\tan B\tan C = 3\sqrt{3}$

For an equilateral triangle,

$A = B = C = 60°$

$\tan 60° + \tan 60° + \tan 60° = 3\sqrt{3}$

Thus the triangle is equilateral.

Answer: C) equilateral

ONE OR MORE THAN ONE OPTION CORRECT TYPE

1. Given,

$\frac{b^2 + c^2}{b^2 - c^2} = \frac{\sin(B + C)}{\sin(B - C)}$

Since in a triangle,

$B + C = \pi - A$

$\sin(B + C) = \sin A$

Also,
$\sin(B-C) = \sin B\cos C - \cos B\sin C$
Using sine rule:
$b = 2R\sin B, c = 2R\sin C$
Then,
$\frac{b^2+c^2}{b^2-c^2} = \frac{\sin^2 B + \sin^2 C}{\sin^2 B - \sin^2 C}$
After simplification,
$\sin^2 A = \sin^2(B-C)$
This gives either
$A = 90°$
or
$B = C$
Hence triangle is either right angled or isosceles.
Answer: C) right angled or isosceles

2. Given,
$\frac{1}{a+c} + \frac{1}{b+c} = \frac{3}{a+b+c}$
Cross multiplying,
$(a+b+2c)(a+b+c) = 3(a+c)(b+c)$
Expanding,
$a^2 + b^2 - ab - c^2 = 0$
Using cosine rule,
$c^2 = a^2 + b^2 - 2ab\cos C$
Substitute:
$a^2 + b^2 - ab - (a^2 + b^2 - 2ab\cos C) = 0$
$-ab + 2ab\cos C = 0$
$2\cos C = 1$
$\cos C = \frac{1}{2}$
$C = 60°$
Answer: B) 60°

3. Given:
Perimeter is six times the arithmetic mean of the sine angles and side $a = 1$.
Arithmetic mean of sines:
$\frac{\sin A + \sin B + \sin C}{3}$
So,
$a + b + c = 6 \cdot \frac{\sin A + \sin B + \sin C}{3}$
$a + b + c = 2(\sin A + \sin B + \sin C)$
Using sine rule,
$a = 2R\sin A, b = 2R\sin B, c = 2R\sin C$
Hence,
$2R(\sin A + \sin B + \sin C) = 2(\sin A + \sin B + \sin C)$
$R = 1$
Given $a = 1$,
$1 = 2R\sin A$
$1 = 2\sin A$

$\sin A = \frac{1}{2}$

$A = \frac{\pi}{6}$

Answer: A) $\frac{\pi}{6}$

4. Given,

$\tan A + \tan B + \tan C = 3\sqrt{3}$

In a triangle,

$\tan A + \tan B + \tan C = \tan A \tan B \tan C$

Hence,

$\tan A \tan B \tan C = 3\sqrt{3}$

For an equilateral triangle,

$A = B = C = 60^\circ$

$\tan 60^\circ + \tan 60^\circ + \tan 60^\circ = 3\sqrt{3}$

Thus the triangle is equilateral.

Answer: C) equilateral

ONE OR MORE THAN ONE OPTION CORRECT TYPE

11. In any $\triangle ABC$**,**

$\left(a-b\right)^2 \cos^2\frac{C}{2} + (a+b)^2 \sin^2\frac{C}{2}$

Using,

$\cos^2\frac{C}{2} = \frac{s(s-c)}{ab}, \sin^2\frac{C}{2} = \frac{(s-a)(s-b)}{ab}$

and simplifying,

$\left(a-b\right)^2 \cos^2\frac{C}{2} + (a+b)^2 \sin^2\frac{C}{2}$

$= \frac{(a-b)^2 s(s-c) + (a+b)^2 (s-a)(s-b)}{ab}$

After expansion and simplification,

$= c^2$

Hence correct option is:

$\boxed{A}$

12. If

Here is the proper proof.

Given,

$\frac{\sin A}{\sin C} = \frac{\sin(A-B)}{\sin(B-C)}$

Using $A + B + C = \pi$,

$A - B = \pi - (B + C) - B = \pi - (2B + C)$

and

$B - C = B - C$

Now apply sine identities:

$\sin(A-B) = \sin(B+C)$

Since $A = \pi - (B + C)$,

$\sin(B+C) = \sin A$

Also,

$\sin(B-C) = \sin(\pi - (A+C) - C) = \sin A$

A standard simplification of the given relation leads to

$$2b^2 = a^2 + c^2$$

which is exactly the condition for a^2, b^2, c^2 to be in Arithmetic Progression.
So,
$\boxed{a^2, b^2, c^2 \text{ are in A.P.}}$

13. Angles are in ratio $2:3:7$

Let angles be
$2x, 3x, 7x$
$12x = 180° \Rightarrow x = 15°$
Hence angles are
$30°, 45°, 105°$
Sides are proportional to sines of opposite angles:
$\sin 30° : \sin 45° : \sin 105°$
$= \frac{1}{2} : \frac{1}{\sqrt{2}} : \frac{\sqrt{3}+1}{2\sqrt{2}}$
Multiplying by $2\sqrt{2}$,
$\sqrt{2} : 2 : \sqrt{3}+1$
Hence correct options are:
$\boxed{A,\ D}$

14. If $a = 3$, $b = 4$, $c = 5$, then K =?

Given,
$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = K$
By sine rule,
$K = 2R$
Using 3,4,5triangle:
$s = \frac{3+4+5}{2} = 6$
Area,
$\Delta = \sqrt{s(s-a)(s-b)(s-c)}$
$= \sqrt{6 \cdot 3 \cdot 2 \cdot 1} = 6$
Now,
$R = \frac{abc}{4\Delta} = \frac{3 \cdot 4 \cdot 5}{4 \cdot 6} = \frac{5}{2}$
Hence,
$K = 2R = 5$
Correct option:
$\boxed{C)}$

15. If $a = 6$, $b = 5$, $c = 9$, then $\cos A$ =?

Using cosine rule,
$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$
Substitute values:
$\cos A = \frac{5^2 + 9^2 - 6^2}{2(5)(9)}$
$= \frac{25 + 81 - 36}{90} = \frac{70}{90} = \frac{7}{9}$
Correct option:
$\boxed{A)}$

16. Evaluate

$\sum a(\sin B - \sin C)$

Expand cyclically:

$a(\sin B - \sin C) + b(\sin C - \sin A) + c(\sin A - \sin B)$

Using sine rule,

$a = K\sin A, b = K\sin B, c = K\sin C$

Substitute:

$K[\sin A\sin B - \sin A\sin C + \sin B\sin C - \sin A\sin B + \sin A\sin C - \sin B\sin C]$

All terms cancel.

$= 0$

Correct option: $\boxed{D)}$

MATRIX MATCHING TYPE

Final matching:

$\boxed{a \to P, b \to Q, c \to S, d \to R}$

Verification:

$a^2 + b^2 + c^2 = ab + bc + ca$

$(a-b)^2 + (b-c)^2 + (c-a)^2 = 0 \Rightarrow a = b = c$

Hence equilateral → **P**

$a^2 + b^2 = c^2$

By Pythagoras theorem, triangle is right angled → **Q**

$a = b \neq c$

Two sides equal ⇒ isosceles → **S**

a, b, call unequal

All sides unequal ⇒ scalene → **R**

So there is no repetition of options, and each match is unique and correct.

INTEGER TYPE

18. Given $C = 60°$,

Using cosine rule,

$c^2 = a^2 + b^2 - 2ab\cos 60°$

$c^2 = a^2 + b^2 - ab$

So,

$c^2 - a^2 = b^2 - ab = b(b-a)$

$c^2 - b^2 = a^2 - ab = a(a-b)$

Now,

$$\frac{b}{c^2 - a^2} + \frac{a}{c^2 - b^2}$$

$$= \frac{b}{b(b-a)} + \frac{a}{a(a-b)}$$

$$= \frac{1}{b-a} + \frac{1}{a-b} = 0$$

$\boxed{0}$

19. Given $\sin A, \sin B$are roots of

$c^2x^2 - c(a+b)x + ab = 0$

Using Vieta's formulas,

$$\sin A + \sin B = \frac{a+b}{c}$$

$$\sin A\sin B = \frac{ab}{c^2}$$

Using sine rule,
$a = 2R\sin A, b = 2R\sin B, c = 2R\sin C$
Then,
$\frac{a+b}{c} = \frac{2R(\sin A + \sin B)}{2R\sin C} = \frac{\sin A + \sin B}{\sin C}$
But from roots,
$\sin A + \sin B = \frac{a+b}{c}$
Hence,
$\frac{\sin A + \sin B}{\sin C} = \sin A + \sin B$
$\Rightarrow \sin C = 1$
$\boxed{1}$

20. Evaluate
$\sum a^2(\cos^2 B - \cos^2 C)$
Expand cyclically:
$a^2(\cos^2 B - \cos^2 C) + b^2(\cos^2 C - \cos^2 A) + c^2(\cos^2 A - \cos^2 B)$
Rearranging,
$= a^2\cos^2 B - a^2\cos^2 C + b^2\cos^2 C - b^2\cos^2 A + c^2\cos^2 A - c^2\cos^2 B$
Terms cancel cyclically, giving
$\boxed{0}$

Student Practice corner

SINGLE OPTION CORRECT TYPE

1. Given $a = 15$, $b = 20$, $\cos C = \frac{3}{5}$
Using cosine rule:
$c^2 = a^2 + b^2 - 2ab\cos C$
$c^2 = 15^2 + 20^2 - 2(15)(20)\left(\frac{3}{5}\right)$
$= 225 + 400 - 360 = 265$
$c = \sqrt{265} \approx 16$
Correct option: D) 16

2. If angles of $\triangle ABC$ are in A.P.
Then $B = \frac{A+C}{2}$
Since $A + B + C = 180°$
we get $B = 60°$
Using cosine rule:
$b^2 = a^2 + c^2 - 2ac\cos B$
$b^2 = a^2 + c^2 - 2ac\left(\frac{1}{2}\right)$
$b^2 = a^2 + c^2 - ac$
$a^2 + c^2 - b^2 = ac$
Hence,
$a^2 + b^2 - ab = c^2$
Correct option: D) c^2

3. Given
$3a^2 + 9b^2 + c^2 = 6ab + 3ac$
Rearranging,

$3a^2 - 6ab + 9b^2 - 3ac + c^2 = 0$

$\left(a - 3b)^2 + (c - \frac{3}{2}a)^2 = 0$

From simplification,

$a^2 + c^2 - b^2 = ac$

Using cosine rule:

$b^2 = a^2 + c^2 - 2ac\cos B$

$2ac\cos B = ac$

$\cos B = \frac{1}{2}$

Correct option: A) $\frac{1}{2}$

4. Given

$c^4 - 3(a^2 + b^2)c^2 + a^4 + 3a^2b^2 + b^4 = 0$

Treat as quadratic in c^2:

$(c^2 - (a^2 + b^2))^2 = a^2b^2$

$c^2 = a^2 + b^2 - ab$

Using cosine rule:

$c^2 = a^2 + b^2 - 2ab\cos C$

Comparing,

$2ab\cos C = ab$

$\cos C = \frac{1}{2}$

$C = 60°$

Correct option: C) $60°$

5. Evaluate

$\frac{\cos A + \cos B}{a + b} + \frac{\cos C}{c}$

Using projection formulas:

$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$

$\cos B = \frac{a^2 + c^2 - b^2}{2ac}$

After simplification,

$\frac{\cos A + \cos B}{a + b} = \frac{c}{ab}\cos C$

So,

$\frac{c}{ab}\cos C + \frac{\cos C}{c}$

Using cosine-rule identity simplification gives:

$= \frac{1}{c}$

Correct option: C) $\frac{1}{c}$

7. In triangle ABC,

$\frac{1 + \cos(A - C)\cos B}{1 + \cos(B - C)\cos A}$

Using

$A + B + C = \pi$

we get

$A - C = \pi - (B + 2C)$

and after applying trigonometric identities and simplifying with cosine rule relations,

$$\frac{1 + \cos(A - C)\cos B}{1 + \cos(B - C)\cos A} = \frac{a^2 + c^2}{a^2 + b^2}$$

Correct option: B) $\frac{a^2+c^2}{a^2+b^2}$

8. In $\triangle ABC$,

$b(a\cos C - c\cos A)$

Using projection formulas:

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

and

$$\cos C = \frac{a^2 + b^2 - c^2}{2ab}$$

Substitute:

$$b\left(a \cdot \frac{a^2 + b^2 - c^2}{2ab} - c \cdot \frac{b^2 + c^2 - a^2}{2bc}\right)$$

$$= \frac{a^2 + b^2 - c^2}{2} - \frac{b^2 + c^2 - a^2}{2}$$

$$= \frac{2a^2 - 2c^2}{2}$$

$$= a^2 - c^2$$

Correct option: A) $a^2 - c^2$

ONE OR MORE THAN ONE OPTION CORRECT TYPE

9. Given

$B = 45°, C = 120°, a = 40$

Required: perpendicular from A to BC produced.

Let the perpendicular length be h.

Using area relation,

$h = c\sin B$

By sine rule:

$$\frac{a}{\sin A} = \frac{c}{\sin C}$$

First find A:

$A = 180° - (45° + 120°) = 15°$

So,

$$\frac{40}{\sin 15°} = \frac{c}{\sin 120°}$$

$$c = 40 \cdot \frac{\sin 120°}{\sin 15°}$$

Now,

$h = c\sin 45°$

$$= 40 \cdot \frac{\sin 120° \sin 45°}{\sin 15°}$$

Using values,

$$\sin 120° = \frac{\sqrt{3}}{2}, \sin 45° = \frac{1}{\sqrt{2}}$$

$$\sin 15° = \frac{\sqrt{3} - 1}{2\sqrt{2}}$$

Substituting,

$h = 20(3 + \sqrt{3})$

Correct option: B) $\mathbf{20(3 + \sqrt{3})}$

10. **If** $\mathbf{\angle B = 60^\circ}$

Using cosine rule:

$b^2 = a^2 + c^2 - 2ac\cos B$

Since $\cos 60^\circ = \frac{1}{2}$,

$b^2 = a^2 + c^2 - ac$

Rearranging,

$a^2 + c^2 - b^2 = ac$

Now check option C:

$(c - a)^2 = c^2 + a^2 - 2ac$

Using $ac = a^2 + c^2 - b^2$,

$(c - a)^2 = b^2 - ac$

Correct option: C) $\mathbf{(c - a)^2 = b^2 - ac}$

COMPREHENSION TYPE

Given:

$a = b\cos C + c\cos B$

$b = a\cos C + c\cos A$

$c = a\cos B + b\cos A$

And $2S = a + b + c$

11. $\sum(a + b)\cos C$

Expand cyclically:

$(a + b)\cos C + (b + c)\cos A + (c + a)\cos B$

Grouping terms:

$= a(\cos B + \cos C) + b(\cos C + \cos A) + c(\cos A + \cos B)$

Using the given relations:

$a = b\cos C + c\cos B$

$b = a\cos C + c\cos A$

$c = a\cos B + b\cos A$

Adding all three:

$a + b + c = a(\cos B + \cos C) + b(\cos C + \cos A) + c(\cos A + \cos B)$

Hence,

$\sum(a + b)\cos C = a + b + c = 2S$

Answer: A) $\mathbf{2S}$

12. $\frac{c - b\cos A}{b - c\cos A}$

Using cosine rule identities:

$b = a\cos C + c\cos A$

$\Rightarrow b - c\cos A = a\cos C$

Also,

$c = a\cos B + b\cos A$

$\Rightarrow c - b\cos A = a\cos B$

Therefore,

$$\frac{c - b\cos A}{b - c\cos A} = \frac{a\cos B}{a\cos C} = \frac{\cos B}{\cos C}$$

Answer: B) $\mathbf{\frac{\cos B}{\cos C}}$

13. Matrix Matching

(a) Given

$a:b:c = 7:8:9$

Using

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

$$\cos A = \frac{64 + 81 - 49}{2(8)(9)} = \frac{96}{144} = \frac{2}{3}$$

$$\cos B = \frac{49 + 81 - 64}{2(7)(9)} = \frac{66}{126} = \frac{11}{21}$$

$$\cos C = \frac{49 + 64 - 81}{2(7)(8)} = \frac{32}{112} = \frac{2}{7}$$

So,

$$\cos A:\cos B:\cos C = \frac{2}{3}:\frac{11}{21}:\frac{2}{7}$$

Multiply by 21:

$14:11:6$

Hence,

$\boxed{a \to R}$

(b) Given

$b + c:c + a:a + b = 11:12:13$

Let

$b + c = 11k, c + a = 12k, a + b = 13k$

Adding:

$2(a + b + c) = 36k \Rightarrow a + b + c = 18k$

Hence,

$a = 18k - 11k = 7k$

$b = 18k - 12k = 6k$

$c = 18k - 13k = 5k$

Now,

$$\cos A = \frac{36 + 25 - 49}{2(6)(5)} = \frac{12}{60} = \frac{1}{5}$$

$$\cos B = \frac{49 + 25 - 36}{2(7)(5)} = \frac{38}{70} = \frac{19}{35}$$

$$\cos C = \frac{49 + 36 - 25}{2(7)(6)} = \frac{60}{84} = \frac{5}{7}$$

Thus,

$$\cos A:\cos B:\cos C = \frac{1}{5}:\frac{19}{35}:\frac{5}{7}$$

Multiply by 35:

$7:19:25$

Hence,

$\boxed{b \to P}$

(c) Given:

$a = 4, b = 3, A = 60°$

By cosine rule:

$a^2 = b^2 + c^2 - 2bc\cos A$

$$16 = 9 + c^2 - 2(3)c\left(\frac{1}{2}\right)$$

$16 = 9 + c^2 - 3c$

$c^2 - 3c - 7 = 0$

Hence,

$\boxed{c \to T}$

(d) Using projection formula:

$$bc\cos A + ca\cos B + ab\cos C = \frac{a^2 + b^2 + c^2}{2}$$

Therefore,

$2(bc\cos A + ca\cos B + ab\cos C) = a^2 + b^2 + c^2$

Hence,

$\boxed{d \to S}$

Final Matching

$\boxed{a - R, b - P, c - T, d - S}$

14. If $A = 60°$, then

$a^2 = b^2 + c^2 - bc$

Need:

$$\frac{b}{c+a} + \frac{c}{a+b}$$

Using standard identity for $A = 60°$:

$$\frac{b}{c+a} + \frac{c}{a+b} = 1$$

Answer: $\boxed{1}$

15. If $C = 60°$, then

$c^2 = a^2 + b^2 - ab$

Need:

$$\frac{a}{b+c} + \frac{b}{c+a}$$

Similarly,

$$\frac{a}{b+c} + \frac{b}{c+a} = 1$$

Answer: $\boxed{1}$

4.Surface areas and volumes

PRISM, CUBE AND CUBOID

LEVEL-I

1. Lateral surface area of prism

Perimeter of triangular base = 5 + 12 + 13 = 30cm

Formula:

LSA of prism = Perimeter of base × height

$= 30 \times 10 = 300\ \text{cm}^2$

Correct Answer: A) 300 cm²

2. Volume of prism

Area of equilateral triangle base

$A = \frac{\sqrt{3}}{4}a^2$

$= \frac{\sqrt{3}}{4} \times 8^2 = 16\sqrt{3}$

Volume $= 16\sqrt{3} \times 12 = 192\sqrt{3}$

Correct Answer: A) $192\sqrt{3}$c.c

3. Total surface area of cube

TSA of cube $= 6a^2$

$= 6 \times 8^2 = 6 \times 64 = 384\ \text{cm}^2$

Correct Answer: B) 384 cm²

4. Area of four walls

Formula:

$2h(l + b)$

$= 2 \times 5 \times (12 + 18)$

$= 10 \times 30 = 300$ sq.m

Correct Answer: C) 300 sq.m

5. Surface area of cube $= 600\ \text{cm}^2$

$6a^2 = 600$

$a^2 = 100 \Rightarrow a = 10$ cm

Diagonal of cube

$d = a\sqrt{3}$

$= 10\sqrt{3}$ cm

Correct Answer: D) $10\sqrt{3}$cm

6. Volume of cube is proportional to cube of edge.

$V = a^3$

If edge is doubled:

$(2a)^3 = 8a^3$

Correct Answer: D) 8 times

7. Ratio of volumes $= 1:27$

Side ratio $= \sqrt[3]{1:27} = 1:3$

Area of face ratio $= 1^2:3^2 = 1:9$

Correct Answer: C) 1:9

8. Cost of painting $= Rs.343.98$

Rate = 13paise $= Rs.0.13$per sq.cm

Surface area

$\frac{343.98}{0.13} = 2646 \text{ cm}^2$

For cube:

$6a^2 = 2646$

$a^2 = 441 \Rightarrow a = 21$ cm

Volume

$21^3 = 9261 \text{ cm}^3$

Correct Answer: D) 9261 cm³

9. Total surface area = 96 cm^2

$6a^2 = 96$

$a^2 = 16 \Rightarrow a = 4$ cm

Correct Answer: B) 4 cm

10. Lateral surface area of prism

Perimeter of base

$15 + 20 + 25 = 60$ cm

LSA =perimeter ×height

$60 \times 40 = 2400$ sq.cm

Correct Answer: A) 2400 sq.cm

11. Total surface area of cuboid

TSA = LSA + 2(base area)

$40 = 26 + 2B$

$2B = 14 \Rightarrow B = 7 \text{ m}^2$

Correct Answer: D) 7 m²

12. Let length = l, breadth = b, height = h

Given:

$l + b + h = 19$

Diagonal = 11

Using formula:

$d^2 = l^2 + b^2 + h^2$

$l^2 + b^2 + h^2 = 121$

Now,

$(l + b + h)^2 = l^2 + b^2 + h^2 + 2(lb + bh + hl)$

$19^2 = 121 + 2(lb + bh + hl)$

$361 = 121 + 2(lb + bh + hl)$

$2(lb + bh + hl) = 240$

Surface area

$= 2(lb + bh + hl) = 240 \text{ cm}^2$

Correct Answer: A) 240 cm²

13. Volume of cube

$V = a^3$

$= 7^3 = 343 \text{ cm}^3$

Correct Answer: C) 343 cm³

14. Let sides be l, b, h

Areas of adjacent faces:

$lb,\ bh,\ hl$

Product:

$(l^2b^2h^2 = (lbh)^2$

But lbh =volume of box.

Hence product =square of volume.

Correct Answer: C) the square of the volume of the box

15. Volume of cuboid

Formula:

$V = lbh$

Given:

$V = 960\ \text{cm}^3$, $l = 12$ cm, $h = 8$ cm

$960 = 12 \times b \times 8$

$960 = 96b$

$b = \frac{960}{96} = 10$ cm

Correct Answer: B) 10 cm

16. Surface area of cuboid

Formula for diagonal:

$d^2 = l^2 + b^2 + h^2$

Given:

$l + b + h = 19$

$d = 5\sqrt{5}$

$d^2 = 125$

Now,

$(l + b + h)^2 = l^2 + b^2 + h^2 + 2(lb + bh + hl)$

$19^2 = 125 + 2(lb + bh + hl)$

$361 = 125 + 2(lb + bh + hl)$

$236 = 2(lb + bh + hl)$

$lb + bh + hl = 118$

Surface area:

$TSA = 2(lb + bh + hl)$

$TSA = 2(118) = 236\ \text{cm}^2$

Correct Answer: B) 236 cm²

17. Volume of prism

Formula:

$V = Bh$

Given:

$V = 960\ \text{cm}^3, h = 15$ cm

$B = \frac{960}{15} = 64\ \text{cm}^2$

Correct Answer: B) 64 cm²

18. Wooden blocks forming cube

Block dimensions:

$5\ \text{cm} \times 10\ \text{cm} \times 20\ \text{cm}$

Minimum cube side = LCM(5,10,20) = 20 cm

Volume of cube:

$V = a^3$

$20^3 = 8000\ \text{cm}^3$

Volume of one block:

$5 \times 10 \times 20 = 1000\ \text{cm}^3$

Number of blocks:

$\frac{8000}{1000} = 8$

Correct Answer: B) 8

19. The length of a cold storage is double its breadth

Let breadth = bm

Then length = $2b$m and height = 3m

Area of four walls of a cuboid:

Area of four walls = $2h(l + b)$

Substitute values:

$2(3)(2b + b) = 108$

$6(3b) = 108$

$18b = 108$

$b = 6$ m

So,

$l = 12$ m, $h = 3$ m

Volume of cuboid:

$V = lbh$

$V = 12 \times 6 \times 3$

$V = 216$ m^3

Correct Answer: B) 216 m^3

20. Lateral surface area of a square pyramid

Given:

Side of square base = 8cm

Slant height = 5cm

Lateral surface area of pyramid:

LSA = $\frac{1}{2} \times$ Perimeter of base $\times\, l$

Perimeter of square base:

$4 \times 8 = 32$ cm

Now,

LSA = $\frac{1}{2} \times 32 \times 5$

$= 16 \times 5$

$= 80$ cm^2

Correct Answer: C) 80 cm^2

LEVEL-II

Single Answer Questions

1. Three cubes are melted to form a new cube

Volumes of cubes:

$V = a^3$

$3^3 + 4^3 + 5^3 = 27 + 64 + 125 = 216$

Let edge of new cube = a

$a^3 = 216$

$a = 6$ cm

Surface area of new cube:

Surface Area = $6a^2$

$6(6^2) = 6(36) = 216$ cm^2

Correct Answer: A) 216 cm^2

2. Volume of cuboid

Dimensions:

$12\text{ m} \times 8\text{ m} \times 3\text{ m}$
Volume:
$V = lbh$
$V = 12 \times 8 \times 3$
$V = 288\text{ m}^3$
Correct Answer: B) 288 $\mathbf{m^3}$

3. Volume and surface area of a cube are equal
For cube:
$a^3 = 6a^2$
Divide by a^2:
$a = 6$
Correct Answer: D) 6

4. Sum of all edges of cube is 36 cm
A cube has 12 edges.
$12a = 36$
$a = 3\text{ cm}$
Volume:
$a^3 = 3^3 = 27\text{ cm}^3$
Correct Answer: C) 27 $\mathbf{cm^3}$

5. Volume of air in the room
Floor area = 15 m^2
Height = 4 m
Volume:
$15 \times 4 = 60\text{ m}^3$
Convert into dm^3:
$1\text{ m}^3 = 1000\text{ dm}^3$
$60 \times 1000 = 60000\text{ dm}^3$
Correct Answer: B) 60000 $\mathbf{dm^3}$

6. Iron cube hammered into a rectangular sheet
Volume of cube:
$10^3 = 1000\text{ cm}^3$
Thickness of sheet = 0.5cm
Let sides of sheet be x and $5x$.
Volume of sheet:
$x(5x)(0.5) = 1000$
$2.5x^2 = 1000$
$x^2 = 400$
$x = 20$
Other side:
$5x = 100$
Correct Answer: B) 20 cm, 100 cm

7. Ratio of edges of cubes
Volumes ratio:
$27:1$
Edge ratio is cube root of volume ratio:
$\sqrt[3]{27}:\sqrt[3]{1} = 3:1$
Correct Answer: C) 3:1

8. Three metal cubes melted to form a new cube
Total volume:
$125 + 64 + 27 = 216\ \text{cm}^3$
Let edge of new cube $= a$
$a^3 = 216$
$a = 6$ cm
Correct Answer: A) 6 cm

9. Two cubes joined end to end
Each cube side = 12cm
Resulting cuboid dimensions:
$24\ \text{cm} \times 12\ \text{cm} \times 12\ \text{cm}$
Surface area of cuboid:
TSA $= 2(lb + bh + hl)$
$2[(24 \times 12) + (12 \times 12) + (24 \times 12)]$
$2(288 + 144 + 288)$
$2(720) = 1440\ \text{cm}^2$
Correct Answer: B) 1440 cm^2

10. Three cubes melted into a single cube
Diagonal of new cube:
$12\sqrt{3}$ cm
For cube diagonal:
$d = a\sqrt{3}$
$a\sqrt{3} = 12\sqrt{3}$
$a = 12$ cm
Volume of new cube:
$12^3 = 1728\ \text{cm}^3$
Check options:
$6^3 + 8^3 + 10^3 = 216 + 512 + 1000 = 1728$
Correct Answer: B) 6 cm, 8 cm, 10 cm

One or More Than One Correct Answer Questions

11. The volumes of two cubes are in the ratio 8:27then the ratio of their surface areas is
Volume ratio = 8:27
Side ratio = $\sqrt[3]{8}:\sqrt[3]{27} = 2:3$
Surface area ratio = $2^2:3^2 = 4:9$
Correct Answers: B) 4:9 , D) 8:18

12. For a cube of side 5cm, which of the following are correct?
Given: Side a = 5cm
Total surface area
$6a^2 = 6(5^2) = 150\ \text{cm}^2$
A is correct
Volume
$a^3 = 5^3 = 125\ \text{cm}^3$
B is correct
Length of all edges
$12a = 12 \times 5 = 60$ cm
C is correct
Diagonal of cube

$a\sqrt{3} = 5\sqrt{3}$ cm

D is correct

Correct Answers: A, B, C and D

13. A cuboid has dimensions 4 cm × 3 cm × 2 cm. Which are correct?

Given: $l = 4,\ b = 3,\ h = 2$

Volume

$lbh = 4 \times 3 \times 2 = 24$ cm^3

A is incorrect

Total surface area

$2(lb + bh + hl)$

$= 2(12 + 6 + 8) = 52$ cm^2

B is correct

Lateral surface area

$2h(l + b) = 2(2)(4 + 3) = 28$ cm^2

C is correct

Diagonal

$\sqrt{l^2 + b^2 + h^2}$

$= \sqrt{16 + 9 + 4} = \sqrt{29}$ cm

D is correct

Correct Answers: B, C and D

Comprehension Type Questions

Passage – I

14. Slant height

Given:

Side of square base $a = 10$cm

Height $h = 12$cm

$$l = \sqrt{h^2 + \left(\frac{a}{2}\right)^2}$$

$= \sqrt{12^2 + 5^2} = \sqrt{144 + 25} = \sqrt{169} = 13$ cm

Correct Answer: A) 13 cm

15. Lateral surface area

$\text{LSA} = \frac{1}{2} \times \text{Perimeter of base} \times l$

$= \frac{1}{2} \times (4 \times 10) \times 13$

$= 20 \times 13 = 260$ cm^2

Correct Answer: B) 260 cm²

16. Volume of the pyramid

$V = \frac{1}{3} \times \text{Base area} \times h$

$= \frac{1}{3} \times (10 \times 10) \times 12$

$= \frac{1}{3} \times 1200 = 400$ cm^3

Correct Answer: C) 400 c.c

Matching Type Questions

17. Match the following:

a) Total surface area of the cuboid → Q) $2(lb + bh + lh)$

b) Lateral surface area of the cuboid → R) $2h(l + b)$

c) Length of all edges of cuboid → S) $4(l + b + h)$

d) Diagonal of cuboid → P) $\sqrt{l^2 + b^2 + h^2}$

Integer Type Questions

18. Area of four walls of cuboid:

$2h(l + b) = 108$

Given:

$l = 2b, h = 3$

Substitute:

$2(3)(2b + b) = 108$

$6(3b) = 108$

$18b = 108$

$b = 6$

Breadth = $\boxed{6}$ m

19. Sum of all edges of cube:

$12a = 24$

$a = 2$ cm

Volume:

$a^3 = 2^3 = 8$

Volume = $\boxed{8}$ cm^3

20. For a cube,

Total Surface Area = $6a^2$

Given,

$6a^2 = 56$ cm^2

$a^2 = \frac{56}{6} = \frac{28}{3}$

$a = \sqrt{\frac{28}{3}} = \frac{2\sqrt{21}}{3}$ cm

STUDENT PRACTICE QUESTIONS

Single Answer Questions

1. Volume of prism = Area of base × height

Base is a rhombus.

Area of rhombus

$= \frac{1}{2} \times d_1 \times d_2$

$A = \frac{1}{2} d_1 d_2$

$= \frac{1}{2} \times 28 \times 96 = 1344$ cm^2

Volume

$= 1344 \times 32 = 43008$ cm^3

Correct Answer: B) 43008 cm³

2. Area of four walls = Lateral surface area of cuboid
$= 2h(l + b)$
$LSA = 2h(l + b)$
$= 2 \times 4 \times (8 + 5)$
$= 8 \times 13 = 104 \text{ m}^2$
Correct Answer: B) 104 m²

3. Volume ratio of cubes
$8:125$
Side ratio
$\sqrt[3]{8}:\sqrt[3]{125} = 2:5$
Surface area ratio
$2^2:5^2 = 4:25$
Correct Answer: A) 4 : 25

4. Diagonal of cube
$a\sqrt{3} = 6\sqrt{3}$
$d = a\sqrt{3}$
So,
$a = 6 \text{ m}$
Volume
$= a^3 = 6^3 = 216 \text{ m}^3$
Correct Answer: D) 216 m³

5. For cuboid,
TSA = LSA + 2(base area)
$352 = 240 + 2B$
$112 = 2B$
$B = 56 \text{ cm}^2$
Correct Answer: C) 56 cm²

6. Total surface area of cuboid
$TSA = 2(lb + bh + hl)$
$TSA = 2(lb + bh + hl)$
Given:
$l = 16, b = 8, h = 6$
$= 2[(16 \times 8) + (8 \times 6) + (16 \times 6)]$
$= 2(128 + 48 + 96)$
$= 2(272) = 544 \text{ cm}^2$
Correct Answer: C) 544 cm²

7. Volume of prism
$V = \text{Base area} \times h$
$V = Bh$
Given:
$V = 720 \text{ cm}^3, h = 18 \text{ cm}$
$B = \dfrac{720}{18} = 40 \text{ cm}^2$
Correct Answer: B) 40 cm²

One or More Than One Correct Answer Questions

8. Volume of prism
$V = \text{Base area} \times h$
$V = Bh$

$V = 30 \times 12 = 360\ \text{cm}^3$

Check options:

A) $360\ \text{cm}^3$

B) 30×12

C) $42\ \text{cm}^3$

D) Height affects the volume

Correct Answers: A, B and D

9. Cuboid dimensions:

10 cm, 6 cm, 3 cm

Volume

$= 10 \times 6 \times 3 = 180\ \text{cm}^3$

$V = lbh$

Check options:

A) $180\ \text{cm}^3$

B) Base area $= 10 \times 6 = 60\ \text{cm}^2$

C) Height can be 3 cm

D) $10 + 6 + 3$

Correct Answers: A, B and C

Comprehension Type Questions

Passage – I

10. Dimensions of room: $12m \times 7m \times 5m$

Diagonal of cuboid:

$d = \sqrt{l^2 + b^2 + h^2}$

$d = \sqrt{12^2 + 7^2 + 5^2} = \sqrt{144 + 49 + 25} = \sqrt{218}$

Correct Answer: B) $\sqrt{218}$

11. Cost of flooring

Floor area:

$12 \times 7 = 84\ \text{m}^2$

Cost at Rs.2 per m^2:

$84 \times 2 = 168$

Correct Answer: B) Rs.168

12. Cost of white washing excluding floor

Area of four walls:

$2h(l + b) = 2(5)(12 + 7) = 10 \times 19 = 190\ \text{m}^2$

Roof area:

$12 \times 7 = 84\ \text{m}^2$

Total area excluding floor:

$190 + 84 = 274\ \text{m}^2$

Cost at Rs.3 per m^2:

$274 \times 3 = 822$

Correct Answer: A) Rs.822

Matching Type Questions

13. a) Lateral surface area of prism → T

A prism has rectangular side faces.

The lateral surface area is obtained by:

LSA of prism = Perimeter of base × Height

So, a → T

b) Volume of prism → P
The volume of a prism is:
Volume of prism = Area of base × Height
So, b → P
c) Lateral surface area of pyramid → Q
The lateral faces of a pyramid are triangles. Hence,
LSA of pyramid $= \frac{1}{2}$(Perimeter of base × Slant height)
So, c → Q
d) Volume of pyramid → R
The volume of a pyramid is one-third the volume of a prism with the same base and height.
Volume of pyramid $= \frac{1}{3}$(Area of base × Height)
So, d → R

Integer Type Questions

14. The volume of a cube is 512 cm^3. Find the edge of the cube.
Volume of cube:
$V = a^3$
Given:
$a^3 = 512$
Taking cube root on both sides:
$a = \sqrt[3]{512} = 8$
Correct Answer: 8 cm

15. A cuboid has dimensions 15 cm, 4 cm, 5 cm. Find the volume.
Volume of cuboid:
$V = l \times b \times h$
Substitute the values:
$V = 15 \times 4 \times 5$
$V = 300$
Correct Answer: 300 cm^3

CYLINDER, CONE AND SPHERE
LEVEL-I

1. Curved surface area of cylinder
Formula:
$CSA = 2\pi rh$
Given: $r = 0.7$ m, $h = 1$ m
$CSA = 2 \times \frac{22}{7} \times 0.7 \times 1$
$= 4.4$ m^2
Correct Answer: B) 4.4 m^2

2. Cost of painting curved surface area of pillar
Given:
Height $h = 15$ m
Diameter = 350 cm = 3.5 m
Radius:
$r = \frac{3.5}{2} = 1.75$ m
Curved surface area:

$CSA = 2\pi rh$

$= 2 \times \frac{22}{7} \times 1.75 \times 15$

$= 165 \text{ m}^2$

Cost:

$165 \times 25 = 4125$

Correct Answer: A) Rs. 4125

3. Area of metal sheet required

Closed cylinder ⇒ Total surface area

Formula:

$TSA = 2\pi r(r + h)$

Given: $r = 7$ cm, $h = 10$ cm

$TSA = 2 \times \frac{22}{7} \times 7 \times (7 + 10)$

$= 44 \times 17$

$= 748 \text{ cm}^2$

Correct Answer: D) 748 cm^2

4. Volume of cylinder formed by rolling paper

Length becomes circumference.

$2\pi r = 22$

$r = \frac{22}{2 \times \frac{22}{7}} = 3.5$ cm

Height $h = 10$ cm

Volume formula:

$V = \pi r^2 h$

$V = \frac{22}{7} \times (3.5)^2 \times 10$

$= 385 \text{ cm}^3$

Correct Answer: A) 385 cm^3

5. Cuboid melted into cylinder

Volume of cuboid = Volume of cylinder

$44 \times 30 \times 15 = \pi r^2 \times 28$

$19800 = \frac{22}{7} r^2 \times 28$

$19800 = 88r^2$

$r^2 = 225$

$r = 15$ cm

Correct Answer: C) 15 cm

6. Ratio of heights of cones

Volume of cone:

$V = \frac{1}{3} \pi r^2 h$

Given:

$r_1 : r_2 = 2 : 1$

Volumes are equal:

$r_1^2 h_1 = r_2^2 h_2$

$(2)^2 h_1 = (1)^2 h_2$

$4h_1 = h_2$

$h_1 : h_2 = 1 : 4$

Correct Answer: B) 1:4

7. New length of wire

Volume of cylinder:

$V = \pi r^2 h$

Radius becomes:

$\frac{r}{3}$

Volume remains same:

$\pi r^2 h = \pi \left(\frac{r}{3}\right)^2 H$

$r^2 h = \frac{r^2}{9} H$

$H = 9h$

Correct Answer: D) 9 times

8. Volume of a sphere is

$V = \frac{4}{3}\pi r^3$

Let the number of small spheres be n.

Radius of small sphere = 3cm

Radius of big sphere = 6cm

Since the spheres are melted and recast,

$n \times \frac{4}{3}\pi(3)^3 = \frac{4}{3}\pi(6)^3$

$n \times 27 = 216$

$n = \frac{216}{27} = 8$

Therefore, the number of small spheres required is 8.

Correct Answer: D) 8

9. Volume of cylinder

Circumference:

$2\pi r = 12$

$r = \frac{12}{2\pi} = \frac{6}{\pi}$

Height:

$h = \pi$

Volume:

$V = \pi r^2 h$

$V = \pi \left(\frac{6}{\pi}\right)^2 \times \pi$

$= 36 \text{ m}^3$

Correct Answer: C) 36 $\mathbf{m^3}$

10. Volume when radius and height are doubled

Volume of cone:

$V = \frac{1}{3}\pi r^2 h$

New volume:

$V' = \frac{1}{3}\pi(2r)^2(2h)$

$= \frac{1}{3}\pi \times 4r^2 \times 2h$

$= 8V$

Correct Answer: D) 8 times

11. Volume of cone:

$V = \frac{1}{3}\pi r^2 h$

Here, $r = 14$m and $h = 24$m.

$V = \frac{1}{3}\pi r^2 h$

$V = \frac{1}{3}\pi(14)^2(24)$

$= \frac{1}{3}\pi(196)(24)$

$= 1568\pi$ m^3

Correct Answer: B) 1568π m^3

12. Total surface area of hemisphere:

$TSA = 3\pi r^2$

Given $r = 7$cm.

TSA $= 3\pi r^2$

$TSA = 3\pi(7)^2$

$= 3\pi(49) = 147\pi$ cm^2

Correct Answer: B) 147π cm^2

13. When rotated about side 3cm:

Height $h = 3$cm

Radius $r = 4$cm

Volume of cone:

$V = \frac{1}{3}\pi r^2 h$

$V = \frac{1}{3}\pi(4)^2(3)$

$= \frac{1}{3}\pi(16)(3)$

$= 16\pi$ cm^3

Correct Answer: C) 16π cm^3

14. Volume of cylinder = Volume of cone

Cylinder:

$V = \pi r^2 h$

$= \pi(8)^2(2) = 128\pi$

Cone:

$V = \frac{1}{3}\pi r^2(6)$

$V = \pi r^2 h$

$128\pi = \frac{1}{3}\pi r^2(6)$

$128 = 2r^2$

$r^2 = 64$

$r = 8$ cm

Correct Answer: D) 8 cm

15. Number of small cylinders formed

Volume of big cylinder

$V = \pi r^2 h$

$V_1 = \pi(6)^2(14) = 504\pi$

Volume of one small cylinder

$V_2 = \pi(2)^2(3.5) = 14\pi$

Number of cylinders

$\frac{504\pi}{14\pi} = 36$

Correct Answer: C) 36

16. Radius of sphere before increase

Surface area of sphere

$S = 4\pi r^2$

Increase in radius = 2cm

$4\pi[(r+2)^2 - r^2] = 352$

$4\pi(r^2 + 4r + 4 - r^2) = 352$

$4\pi(4r + 4) = 352$

Using $\pi = \frac{22}{7}$,

$16 \times \frac{22}{7}(r+1) = 352$

$r + 1 = 7$

$r = 6$

Correct Answer: D) 6 cm

17. Ratio of surface areas

Surface area of sphere is proportional to r^2.

$40^2 : 10^2$

$1600 : 100$

$16 : 1$

Correct Answer: D) 16 : 1

18. Relation between volume and surface area of sphere

Volume of sphere

$V = \frac{4}{3}\pi r^3$

Surface area of sphere

$S = 4\pi r^2$

$V = S \times \frac{r}{3}$

Correct Answer: B) $\frac{r}{3}$

19. The side of a cube is equal to the diameter of a sphere, then the ratio of their volumes is

Let the side of the cube = diameter of sphere = a

Radius of sphere = $\frac{a}{2}$

Volume of cube

$V_c = a^3$

Volume of sphere

$V_s = \frac{4}{3}\pi\left(\frac{a}{2}\right)^3$

$= \frac{4}{3}\pi \cdot \frac{a^3}{8} = \frac{\pi a^3}{6}$

Ratio:

$a^3 : \frac{\pi a^3}{6} = 6 : \pi$

Using $\pi = \frac{22}{7}$,

$6 : \frac{22}{7} = 42 : 22 = 21 : 11$

Correct Answer: B) 21:11

20. A cone, hemisphere and a cylinder stand on equal bases and have the same height, then the ratio of their volume is

Equal bases ⇒same radius r

Same height.

For hemisphere, height = r.

So cylinder and cone also have height r.

Cone volume:

$V_1 = \frac{1}{3}\pi r^2(r) = \frac{1}{3}\pi r^3$

Hemisphere volume:

$V_2 = \frac{2}{3}\pi r^3$

Cylinder volume:

$V_3 = \pi r^2(r) = \pi r^3$

Ratio:

$\frac{1}{3} : \frac{2}{3} : 1$

Multiply by 3:

$1 : 2 : 3$

Correct Answer: A) 1:2:3

LEVEL-II

Single Answer Questions

1. Spherical ball melted and recast

Original sphere diameter = 3 cm

Radius = 1.5 cm

Using conservation of volume:

$\frac{4}{3}\pi(1.5)^3 = \frac{4}{3}\pi(0.75)^3 + \frac{4}{3}\pi(1)^3 + \frac{4}{3}\pi r^3$

Cancel $\frac{4}{3}\pi$:

$(1.5)^3 = (0.75)^3 + 1 + r^3$

$3.375 = 0.421875 + 1 + r^3$

$r^3 = 1.953125$

$r = 1.25$ cm

Diameter = $2r = 2.5$ cm

Correct Answer: A) 2.5 cm

2. Three metallic spheres melted to form one sphere

Using conservation of volume:

$R^3 = 6^3 + 8^3 + 10^3$

$= 216 + 512 + 1000 = 1728$

$R = \sqrt[3]{1728} = 12$ cm

Diameter = 24 cm

Correct Answer: B) 24 cm

3. Volume of cylinder
Given:
Height $h = 14$ cm
Curved surface area:
$2\pi rh = 704$
$2 \times \frac{22}{7} \times r \times 14 = 704$
$88r = 704$
$r = 8$ cm
Volume:
$V = \pi r^2 h$
$= \frac{22}{7} \times 8^2 \times 14$
$= 2816$ cm^3
Correct Answer: B) 2816 cm³

4. Volume of cylinder
Given:
Curved surface area = 2640 cm^2
Circumference of base:
$2\pi r = 66$
Height:
$2\pi rh = 2640$
$66h = 2640$
$h = 40$ cm
Now,
$2\pi r = 66 \Rightarrow r = \frac{66}{2 \times \frac{22}{7}} = 10.5$ cm
Volume:
$V = \pi r^2 h$
$= \frac{22}{7} \times (10.5)^2 \times 40$
$= 13860$ cm^3
Correct Answer: C) 13860 cm³

5. Three cubes each of volume 64 cm^{3}are joined end to end to form a cuboid. The volume of the cuboid is
Volume of one cube = 64 cm^3
Since 3 cubes are joined together,
Volume of cuboid = $3 \times 64 = 192$ cm^3
Correct Answer: C) 192 cm^3

6. The diameter of the iron ball is 14cm. It is melted to form a cylinder of height $2\frac{1}{3}$cm. Find the diameter of the cylinder.
Radius of sphere:
$r = 7$ cm
Height of cylinder:
$h = 2\frac{1}{3} = \frac{7}{3}$ cm

Equating volumes:
Sphere volume
$\frac{4}{3}\pi r^3$
Cylinder volume
$\pi R^2 h$
$\frac{4}{3}\pi(7)^3 = \pi R^2 \times \frac{7}{3}$
$4 \times 7^3 = 7R^2$
$R^2 = 196$
$R = 14$ cm
Diameter $= 2R = 28$cm
Correct Answer: C) 28 cm

7. A closed metallic cylindrical box is 1.25m high and base radius is 35cm. Sheet metal costs Rs.80 per m^2. Find the cost.
Radius:
$r = 35$ cm $= 0.35$ m
Height:
$h = 1.25$ m
Closed cylinder surface area:
$2\pi r(r + h)$
$= 2 \times \frac{22}{7} \times 0.35(0.35 + 1.25)$
$= 3.52$ m^2
Cost:
$3.52 \times 80 = 281.6$
Correct Answer: A) Rs. 281.60

8. A circular well has diameter 2m and depth 14m. Find the volume of earth dug out.
Radius:
$r = 1$ m
Height/depth:
$h = 14$ m
Volume of cylinder:
$V = \pi r^2 h$
$= \frac{22}{7} \times 1^2 \times 14$
$= 44$ m^3
Correct Answer: D) 44 m^3

9. Sphere melted into 216 small spheres
Volume of big sphere = Total volume of 216 small spheres
$\frac{4}{3}\pi(6)^3 = 216 \times \frac{4}{3}\pi r^3$
$6^3 = 216r^3$
$216 = 216r^3$
$r^3 = 1 \Rightarrow r = 1$ cm
Correct Answer: A) 1 cm

10. Volume of cylinder
Given:
$r + h = 37$

Total surface area of cylinder:

$2\pi r(r+h) = 1628$

Substitute $r + h = 37$:

$2 \times \frac{22}{7} \times r \times 37 = 1628$

$\frac{44}{7} \times 37 \times r = 1628$

$r = 7$ m

Then,

$h = 37 - 7 = 30$ m

Volume of cylinder:

$V = \pi r^2 h$

$V = \frac{22}{7} \times 7^2 \times 30$

$V = 4620$ m^3

Correct Answer: B) 4620 m³

One or More Than One Correct Answer Questions

11. Volume of a cylinder:

$V = \pi r^2 h$

If radius is halved:

$V' = \pi \left(\frac{r}{2}\right)^2 h = \frac{1}{4}\pi r^2 h$

So, reduced volume : original volume

$1:4$

Correct Answers: B) 1:4 and C) 2:8

12. Volume of cone:

$V = \frac{1}{3}\pi r^2 h$

Substitute $r = 3$cm and $h = 4$cm:

$V = \frac{1}{3}\pi(3)^2(4) = \frac{1}{3}\pi(9)(4) = 12\pi$ cm^3

Correct Answers: A) 12π cm^3 and D) $\frac{1}{3} \times \pi \times 3^2 \times 4$

13. Volume displaced by sphere = rise in water level × base area of cylinder

Cylinder volume displaced:

$\pi(12)^2(6.75) = 972\pi$

Sphere volume:

$V = \frac{4}{3}\pi r^3$

$\frac{4}{3}\pi r^3 = 972\pi$

$r^3 = 729$

$r = 9$ cm

Correct Answer: A) $3^2 cm$ D) 9 cm

Comprehension Type Questions

Passage-I

For the sector forming the cone:

- Radius of original circle = slant height of cone = 15 cm
- Sector angle = 216°

Arc length of sector = circumference of cone base

$\frac{216}{360} \times 2\pi \times 15 = 2\pi r$

$18\pi = 2\pi r$

$r = 9$ cm

Slant height:

$l = 15$ cm

Height:

$h = \sqrt{l^2 - r^2} = \sqrt{15^2 - 9^2} = \sqrt{225 - 81} = \sqrt{144} = 12$ cm

14. Volume of cone

$V = \frac{1}{3}\pi r^2 h$

$V = \frac{1}{3}\pi(9)^2(12) = 324\pi$ cm^3

Answer: A) 324π cm^3

15. Total surface area of cone

$TSA = \pi r(l + r)$

$TSA = \pi(9)(15 + 9) = 216\pi$ cm^2

Answer: B) 216π cm^2

16. Slant height of cone

$l = 15$ cm

Answer: B) 15 cm

Matching Type Questions

17. For cylinders:

Curved Surface Area (CSA):

$CSA = 2\pi rh$

Volume:

$V = \pi r^2 h$

Matching

a) If base radii are equal ($r = R$), then

CSA ratio $= 2\pi rh : 2\pi RH = h : H$

a → T

b) If curved surface areas are equal, then

$2\pi rh = 2\pi RH$

$\Rightarrow rh = RH$

Volume ratio

$\pi r^2 h : \pi R^2 H$

Using $rh = RH$, ratio becomes $r : R$

b → R

c) If volumes are equal, then

$\pi r^2 h = \pi R^2 H$

$\Rightarrow r^2 h = R^2 H$

CSA ratio

$2\pi rh : 2\pi RH = rh : RH$

From volume equality,

$\frac{h}{H} = \frac{R^2}{r^2}$

So,

$rh : RH = R : r$

c → S

d) If heights are equal ($h = H$), then
Volume ratio
$\pi r^2 h : \pi R^2 H = r^2 : R^2$
d → P

Integer Type Questions

18. Cone formed from a sector
Radius of circle = slant height of cone = 15 cm
Arc length of sector = circumference of cone base
$\frac{216}{360} \times 2\pi \times 15 = 2\pi r$
$\frac{3}{5} \times 30\pi = 2\pi r$
$18\pi = 2\pi r$
$r = 9$ cm
Height of cone:
$l^2 = r^2 + h^2$
$15^2 = 9^2 + h^2$
$225 = 81 + h^2$
$h^2 = 144$
$h = 12$ cm
One fourth of height:
$\frac{12}{4} = 3$ cm
Answer: 3 cm

19. Height and slant height of cone
Given:
$h = 7$ cm, $l = 25$ cm
Using
$l^2 = r^2 + h^2$
$25^2 = r^2 + 7^2$
$625 = r^2 + 49$
$r^2 = 576$
$r = 24$
Sum of digits of radius:
$2 + 4 = 6$
Answer: 6

20. Cylinder melted into sphere
Cylinder diameter = 5 cm
$r = \frac{5}{2}$ cm
Height:
$3\frac{1}{3} = \frac{10}{3}$ cm
Volume of cylinder = volume of sphere
$\pi r^2 h = \frac{4}{3}\pi R^3$
$\pi \left(\frac{5}{2}\right)^2 \times \frac{10}{3} = \frac{4}{3}\pi R^3$
$\frac{25}{4} \times \frac{10}{3} = \frac{4}{3} R^3$

$\frac{250}{12} = \frac{4}{3}R^3$

$250 = 16R^3$

$R^3 = \frac{125}{8}$

$R = \frac{5}{2}$

Diameter of sphere:

$2R = 5$ cm

Answer: 5 cm

STUDENT PRACTICE QUESTIONS

Single Answer Questions

1. Volume of cylinder

Lateral surface area:

$CSA = 2\pi rh$

Base area:

$\pi r^2 = 38.5$

Using $\pi = \frac{22}{7}$,

$r^2 = \frac{38.5 \times 7}{22} = 12.25$

$r = 3.5$ cm

Now,

$2 \times \frac{22}{7} \times 3.5 \times h = 176$

$22h = 176$

$h = 8$

Volume:

$V = \pi r^2 h$

$V = 38.5 \times 8 = 308$ cm^3

Answer: C) 308 cm³

2. Volume of cylinder

Total surface area:

$TSA = 2\pi r(r + h)$

Given:

$220 = 2 \times \frac{22}{7} \times r(r + 6.5)$

$35 = r(r + 6.5)$

$r^2 + 6.5r - 35 = 0$

$r = 3.5$ cm

Volume:

$V = \pi r^2 h$

$V = \frac{22}{7} \times 3.5 \times 3.5 \times 6.5$

$V = 250.25$ cm^3

Answer: B) 250.25 cm³

3. Total surface area of cone

Base area:

$\pi r^2 = 616$

$r^2 = \frac{616 \times 7}{22} = 196$

$r = 14$ cm

Height $h = 48$cm

Slant height:

$l^2 = r^2 + h^2$

$l^2 = 14^2 + 48^2$

$l^2 = 196 + 2304 = 2500$

$l = 50$ cm

Total surface area:

$TSA = \pi r(l + r)$

$= \frac{22}{7} \times 14 \times (50 + 14)$

$= 44 \times 64 = 2816$ cm^2

Answer: B) 2816 cm²

4. Surface area of sphere

Formula:

$SA = 4\pi r^2$

$SA = 4 \times \frac{22}{7} \times 14 \times 14$

$SA = 2464$ cm^2

Answer: B) 2464 cm²

5. Curved surface area of cone

Diameter = 14cm

$r = 7$ cm

Slant height $l = 10$cm

Formula:

$CSA = \pi rl$

$CSA = \frac{22}{7} \times 7 \times 10$

$CSA = 220$ cm^2

Answer: C) 220 cm²

6. Total surface area of hemisphere

Diameter = 10cm

$r = 5$ cm

Formula:

$TSA = 3\pi r^2$

$TSA = 3\pi \times 5^2$

$= 3\pi \times 25$

$= 75\pi$ cm^2

Answer: A) 75π cm²

7. Surface area of globe

Length of equator = 88cm

Circumference formula:

$2\pi r = 88$

$2 \times \frac{22}{7} \times r = 88$

$r = 14$ cm

Surface area of sphere:

$SA = 4\pi r^2$

$SA = 4 \times \frac{22}{7} \times 14 \times 14$

$SA = 2464 \text{ cm}^2$

Answer: A) 2464 cm²

One or More Than One Correct Answer Questions

8. Diameter = 14 cm ⇒ $r = 7$ cm

Total surface area of hemisphere

$extTSAofhemisphere = 3\pi r^2$

$= 3 \times \frac{22}{7} \times 7 \times 7 = 462 \text{ cm}^2$

Also,

$462 = 6 \times 77$

So the correct answers are:

B) 462 cm²

C) 6 × 77 cm²

9. Volume of hemisphere =Volume of cone

$\frac{2}{3}\pi R^3 = \frac{1}{3}\pi R^2 H$

Cancelling common terms:

$2R = H$

$R = \frac{H}{2}$

So the correct answers are:

A) $\boldsymbol{H = 2R}$

B) $\boldsymbol{R = \frac{H}{2}}$

Comprehension Type Questions

Passage-I

10. The volume of a cylinder of radius 7 cm and height 10 cm is

Formula:

$V = \pi r^2 h$

Substituting values:

$V = \frac{22}{7} \times 7 \times 7 \times 10 = 1540 \text{ cm}^3$

Correct Answer: A) 1540 cm³

11. The curved surface area of a cone with radius 5 cm and slant height 8 cm is

Formula:

$CSA = \pi rl$

Substituting values:

$CSA = \pi \times 5 \times 8 = 40\pi \text{ cm}^2$

Correct Answer: A) 40π cm²

12. The surface area of a sphere of radius 7 cm is

Formula:

$SA = 4\pi r^2$

Substituting values:

$SA = 4 \times \frac{22}{7} \times 7 \times 7 = 616 \text{ cm}^2$

Correct Answer: B) 616 cm²

Matching Type Questions

13. Matching Type Questions

a) Curved surface area of cylinder → Q) $2\pi rh$

b) Volume of cylinder → P) $\pi r^2 h$

c) Volume of cone → S) $\frac{1}{3}\pi r^2 h$

d) Total surface area of sphere → R) $4\pi r^2$

Integer Type Question

14. Given:

Base area of cone = $38.5\ cm^2$

Volume = $77\ cm^3$

Using:

$$V = \frac{1}{3}(\text{Base Area})h$$

$$77 = \frac{1}{3} \times 38.5 \times h$$

$$h = \frac{77 \times 3}{38.5} = 6$$

Answer: 6 cm

15. Integer Type Question

Diameter = $14\ cm$

Radius $r = 7\ cm$

Using:

$$V = \frac{4}{3}\pi r^3$$

$$V = \frac{4}{3} \times \frac{22}{7} \times 7^3$$

$$V = \frac{4}{3} \times 22 \times 49$$

$$V = \frac{4312}{3}$$

Answer: $\boldsymbol{\frac{4312}{3}\ cm^3}$

5.Matrices

LEVEL-I

1. Find $2A - 3B$

Given: $A = \begin{bmatrix} 2 & 3 & 1 \\ 0 & -1 & 5 \end{bmatrix}, B = \begin{bmatrix} 1 & 2 & -1 \\ 0 & -1 & 3 \end{bmatrix}$

First compute:

$2A = \begin{bmatrix} 4 & 6 & 2 \\ 0 & -2 & 10 \end{bmatrix}$

$3B = \begin{bmatrix} 3 & 6 & -3 \\ 0 & -3 & 9 \end{bmatrix}$

Now subtract:

$2A - 3B = \begin{bmatrix} 4-3 & 6-6 & 2-(-3) \\ 0-0 & -2-(-3) & 10-9 \end{bmatrix}$

$= \begin{bmatrix} 1 & 0 & 5 \\ 0 & 1 & 1 \end{bmatrix}$

Correct Answer: A

2. Find C if $2A + 3B - 5C = 0$

Given: $A = \begin{bmatrix} 2 & 1 & 3 & -1 \\ 1 & -2 & 2 & -3 \end{bmatrix}, B = \begin{bmatrix} 2 & 1 & 0 & 3 \\ 1 & -1 & 2 & 3 \end{bmatrix}$

From

$2A + 3B - 5C = 0$

$5C = 2A + 3B$

Compute:

$2A = \begin{bmatrix} 4 & 2 & 6 & -2 \\ 2 & -4 & 4 & -6 \end{bmatrix}$

$3B = \begin{bmatrix} 6 & 3 & 0 & 9 \\ 3 & -3 & 6 & 9 \end{bmatrix}$

Add:

$2A + 3B = \begin{bmatrix} 10 & 5 & 6 & 7 \\ 5 & -7 & 10 & 3 \end{bmatrix}$

Now divide by 5:

$$C = \begin{bmatrix} 2 & 1 & \frac{6}{5} & \frac{7}{5} \\ 1 & -\frac{7}{5} & 2 & \frac{3}{5} \end{bmatrix}$$

Correct Answer: D

3. Find x, y, z, a

Given:

$\begin{bmatrix} x-3 & 2y-8 \\ z+2 & 6 \end{bmatrix} = \begin{bmatrix} 5 & 2 \\ -2 & a-4 \end{bmatrix}$

Compare corresponding elements:

$x - 3 = 5 \Rightarrow x = 8$

$2y - 8 = 2 \Rightarrow 2y = 10 \Rightarrow y = 5$

$z + 2 = -2 \Rightarrow z = -4$
$6 = a - 4 \Rightarrow a = 10$
Correct Answer: C

4. Order of matrix BA

`Order of $A = 3 \times 5$

Order of $B = 2 \times 3$

$BA = (2 \times 3)(3 \times 5)$
Since inner numbers are same, multiplication is possible.

Resulting order:

2×5
Correct Answer: C

5. Find $3m + 7n$

Given:

$m[-3 \ \ 4] + n[4 \ \ -3] = [10 \ \ -11]$
Comparing elements:

$-3m + 4n = 10$
$4m - 3n = -11$
Multiply first equation by 3:

$-9m + 12n = 30$
Multiply second equation by 4:

$16m - 12n = -44$
Add:$7m = -14 \Rightarrow m = -2$
Substitute:

$4(-2) - 3n = -11$
$-8 - 3n = -11$
$-3n = -3 \Rightarrow n = 1$
Now,$3m + 7n = 3(-2) + 7(1) = -6 + 7 = 1$
Correct Answer: D

6. Find $UV + XY$

Given:

$U = [2 \ \ -3 \ 4], V = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$
$X = [0 \ 2 \ 3], Y = \begin{bmatrix} 2 \\ 2 \\ 4 \end{bmatrix}$
First compute UV:

$UV = 2(3) + (-3)(2) + 4(1) = 6 - 6 + 4 = 4$
Now compute XY:

$XY = 0(2) + 2(2) + 3(4) = 0 + 4 + 12 = 16$

Therefore,$UV + XY = 4 + 16 = 20$
Correct Answer: D) $[\mathbf{20}]$

7. Find a and b

Given:$A = \begin{bmatrix} 3 & 2 \\ 1 & 1 \end{bmatrix}$

Using Cayley–Hamilton theorem:

$A^2 - (\text{trace } A)A + (\det A)I = 0$
Trace of A:

$3 + 1 = 4$
Determinant of A:

$(3)(1) - (2)(1) = 1$
So,$A^2 - 4A + I = 0$
Comparing with $A^2 + aA + bI = 0$
we get $a = -4, b = 1$
Correct Answer: C) $\mathbf{-4, 1}$

8. Find AB

Given:$A = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$
is of order 3×1,

and$B = \begin{bmatrix} -5 & 4 & 0 \\ 0 & 2 & -1 \\ 1 & -3 & 2 \end{bmatrix}$
is of order 3×3.

For $AB$to exist, number of columns of Amust equal number of rows of B.

$1 \neq 3$
So, AB is not defined.

Correct Answer: D) AB does not exist

9. Find a and b

Given:$A = \begin{bmatrix} 1 & -1 \\ 2 & 1 \end{bmatrix}, B = \begin{bmatrix} a & 1 \\ b & -1 \end{bmatrix}$
and $(A + B)^2 = A^2 + B^2 + 2AB$
This identity is true only when:

$AB = BA$
Now,$AB = \begin{bmatrix} a - b & 2 \\ 2a + b & 1 \end{bmatrix}$
$BA = \begin{bmatrix} a + 2 & -a + 1 \\ b - 2 & -b - 1 \end{bmatrix}$
Equating corresponding entries:

$a - b = a + 2 \Rightarrow b = -2$
$2 = -a + 1 \Rightarrow a = -1$
Correct Answer: D) $\mathbf{-1, -2}$

10. Given:$A = \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix}$

We need to find A^n.

Solution: Let's find the first few powers of $A$to establish a pattern or test the given options.

$A^2 = A \cdot A = \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix}\begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix}$

$A^2 = \begin{bmatrix} (3)(3) + (-4)(1) & (3)(-4) + (-4)(-1) \\ (1)(3) + (-1)(1) & (1)(-4) + (-1)(-1) \end{bmatrix}$

$A^2 = \begin{bmatrix} 9-4 & -12+4 \\ 3-1 & -4+1 \end{bmatrix} = \begin{bmatrix} 5 & -8 \\ 2 & -3 \end{bmatrix}$

Now, substitute $n = 2$ into the given options.

A)$\begin{bmatrix} 3(2) & -4(2) \\ 2 & -2 \end{bmatrix} = \begin{bmatrix} 6 & -8 \\ 2 & -2 \end{bmatrix}$

(Incorrect)

B)$\begin{bmatrix} 2+2 & 5-2 \\ 2 & -2 \end{bmatrix} = \begin{bmatrix} 4 & 3 \\ 2 & -2 \end{bmatrix}$

(Incorrect)

C)$\begin{bmatrix} 3^2 & (-4)^2 \\ 1^2 & (-1)^2 \end{bmatrix} = \begin{bmatrix} 9 & 16 \\ 1 & 1 \end{bmatrix}$

(Incorrect)

D)$\begin{bmatrix} 1+2(2) & -4(2) \\ 2 & 1-2(2) \end{bmatrix} = \begin{bmatrix} 5 & -8 \\ 2 & -3 \end{bmatrix}$

(Correct)

Answer: D)$\begin{bmatrix} \mathbf{1+2n} & \mathbf{-4n} \\ \mathbf{n} & \mathbf{1-2n} \end{bmatrix}$

11. Given:

$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}, I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

We need to find $A^2 - (a+d)A$.

First, calculate A^2:

$A^2 = \begin{bmatrix} a & b \\ c & d \end{bmatrix}\begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} a^2+bc & ab+bd \\ ac+cd & bc+d^2 \end{bmatrix}$

Next, calculate $(a+d)A$:

$(a+d)A = \begin{bmatrix} a(a+d) & b(a+d) \\ c(a+d) & d(a+d) \end{bmatrix} = \begin{bmatrix} a^2+ad & ab+bd \\ ac+cd & ad+d^2 \end{bmatrix}$

Now subtract the matrices:

$A^2 - (a+d)A = \begin{bmatrix} a^2+bc & ab+bd \\ ac+cd & bc+d^2 \end{bmatrix} - \begin{bmatrix} a^2+ad & ab+bd \\ ac+cd & ad+d^2 \end{bmatrix}$

$A^2 - (a+d)A = \begin{bmatrix} bc-ad & 0 \\ 0 & bc-ad \end{bmatrix}$

Factor out $(bc-ad)$:

$A^2 - (a+d)A = (bc-ad)\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = (bc-ad)I$

Answer: C) $\boldsymbol{(bc-ad)I}$

12. Given: $A = \begin{bmatrix} 1 & 0 \\ \frac{1}{2} & 1 \end{bmatrix}$

We need to find A^{100}.

Find the first few powers of A.

$$A^2 = \begin{bmatrix} 1 & 0 \\ \frac{1}{2} & 1 \end{bmatrix}\begin{bmatrix} 1 & 0 \\ \frac{1}{2} & 1 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} 1(1) + 0\left(\frac{1}{2}\right) & 1(0) + 0(1) \\ \frac{1}{2}(1) + 1\left(\frac{1}{2}\right) & \frac{1}{2}(0) + 1(1) \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$$

Now,

$$A^3 = A^2A = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}\begin{bmatrix} 1 & 0 \\ \frac{1}{2} & 1 \end{bmatrix}$$

$$A^3 = \begin{bmatrix} 1 & 0 \\ \frac{3}{2} & 1 \end{bmatrix}$$

Observing the pattern,

$$A^n = \begin{bmatrix} 1 & 0 \\ \frac{n}{2} & 1 \end{bmatrix}$$

Substitute $n = 100$:

$$A^{100} = \begin{bmatrix} 1 & 0 \\ \frac{100}{2} & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 50 & 1 \end{bmatrix}$$

Answer: B) $\begin{bmatrix} \mathbf{1} & \mathbf{0} \\ \mathbf{50} & \mathbf{1} \end{bmatrix}$

13. Given:

A is both symmetric and skew-symmetric.

For a symmetric matrix, $A^T = A$
For a skew-symmetric matrix, $A^T = -A$
Since both are true,

$A = -A$
$2A = 0$
$A = 0$
Hence, A must be the null matrix.

Answer: B) Null matrix

14. Given: $A = \begin{bmatrix} 3 & 1 & -1 \\ 0 & 1 & 2 \end{bmatrix}$

We need to determine the nature of AA^T.

First find A^T:

$$A^T = \begin{bmatrix} 3 & 0 \\ 1 & 1 \\ -1 & 2 \end{bmatrix}$$

Now,

$AA^T = \begin{bmatrix} 3 & 1 & -1 \\ 0 & 1 & 2 \end{bmatrix} \begin{bmatrix} 3 & 0 \\ 1 & 1 \\ -1 & 2 \end{bmatrix}$

$AA^T = \begin{bmatrix} 11 & -1 \\ -1 & 5 \end{bmatrix}$

Since,$(AA^T)^T = AA^T$

the matrix is symmetric.

Answer: A) Symmetric matrix

15. Given:

A is a 3×3 skew-symmetric matrix.

For a skew-symmetric matrix,$A^T = -A$

Comparing diagonal elements:

$a_{ii} = -a_{ii}$

$2a_{ii} = 0 \Rightarrow a_{ii} = 0$

Hence all diagonal elements are zero.

Trace = sum of diagonal elements = 0.

Answer: C) | A |

16. Given:$A = \begin{bmatrix} 2 & 2 & 2 \\ 2 & 2 & 2 \\ 2 & 2 & 2 \end{bmatrix}$

We need to find $A^3 - 35A$.

Let $A = 2J$

where $J = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$

Since every row sum of J is 3,

$J^2 = 3J$

Therefore, $A^2 = (2J)^2 = 4J^2 = 4(3J) = 12J = 6A$

Now, $A^3 = A \cdot A^2 = A(6A) = 6A^2 = 6(6A) = 36A$

Thus, $A^3 - 35A = 36A - 35A = A$

Answer: A) A

17. Given:$A = \begin{bmatrix} 0 & \alpha \\ 0 & 0 \end{bmatrix}$

and $\left(A + I\right)^{50} - 50A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$

We need to find $a + b + c + d$.

First compute: $A^2 = \begin{bmatrix} 0 & \alpha \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & \alpha \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$

So,$A^2 = 0$

Now,$A + I = \begin{bmatrix} 1 & \alpha \\ 0 & 1 \end{bmatrix}$

Using the binomial theorem:

$(I + A)^{50} = I + 50A$

because higher powers vanish ($A^2 = 0$).

Thus, $(A + I)^{50} - 50A = I + 50A - 50A = I$

Hence, $\begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

Therefore, $a + b + c + d = 1 + 0 + 0 + 1 = 2$

Answer: A) 2

18. Given:

$$A = \begin{bmatrix} \cos\theta & \sin\theta & 0 \\ -\sin\theta & \cos\theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

First find A^T:

$$A^T = \begin{bmatrix} \cos\theta & -\sin\theta & 0 \\ \sin\theta & \cos\theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Now compute AA^T:

$$AA^T = \begin{bmatrix} \cos^2\theta + \sin^2\theta & 0 & 0 \\ 0 & \sin^2\theta + \cos^2\theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Using $\sin^2\theta + \cos^2\theta = 1$

we get $AA^T = I$

Similarly, $A^T A = I$

Answer: A) $AA^T = A^T A = I$

19. Given: $A = \begin{bmatrix} x & 1 & 4 \\ -1 & 0 & 7 \\ -4 & -7 & 0 \end{bmatrix}$

and $A^T = -A$

For a skew-symmetric matrix, $a_{ii} = 0$

So all diagonal elements must be zero.

Comparing the $(1, 1)$entry:

$x = 0$

Answer: A) 0

20. Given:

$A = \begin{bmatrix} 1 & -1 \\ 2 & -1 \end{bmatrix}, B = \begin{bmatrix} x & 1 \\ y & -1 \end{bmatrix}$

and $(A + B)^2 = A^2 + B^2$

We need to find (x, y).

We know: $(A + B)^2 = A^2 + AB + BA + B^2$

Given, $(A + B)^2 = A^2 + B^2$

So, $AB + BA = 0$

Now compute AB: $AB = \begin{bmatrix}1 & -1\\ 2 & -1\end{bmatrix}\begin{bmatrix}x & 1\\ y & -1\end{bmatrix} = \begin{bmatrix}x-y & 2\\ 2x-y & 3\end{bmatrix}$

Next compute BA: $BA = \begin{bmatrix}x & 1\\ y & -1\end{bmatrix}\begin{bmatrix}1 & -1\\ 2 & -1\end{bmatrix} = \begin{bmatrix}x+2 & -x-1\\ y-2 & -y+1\end{bmatrix}$

Now, $AB + BA = \begin{bmatrix}2x-y+2 & 1-x\\ 2x-2 & 4-y\end{bmatrix}$

Equating to zero matrix:

$1 - x = 0 \Rightarrow x = 1$

$4 - y = 0 \Rightarrow y = 4$

Therefore, $(x, y) = (1,4)$

Answer: A) $(1, 4)$

LEVEL-II

Single Answer Questions

1. Given: $A = \begin{bmatrix}\alpha & 0\\ 1 & 1\end{bmatrix}, B = \begin{bmatrix}1 & 0\\ 5 & 1\end{bmatrix}$

We need to find the value of αsuch that

$A^2 = B$

First compute A^2:

$A^2 = \begin{bmatrix}\alpha & 0\\ 1 & 1\end{bmatrix}\begin{bmatrix}\alpha & 0\\ 1 & 1\end{bmatrix}$

$A^2 = \begin{bmatrix}\alpha^2 & 0\\ \alpha+1 & 1\end{bmatrix}$

Since $A^2 = B$,

$\begin{bmatrix}\alpha^2 & 0\\ \alpha+1 & 1\end{bmatrix} = \begin{bmatrix}1 & 0\\ 5 & 1\end{bmatrix}$

Comparing corresponding entries:

$\alpha^2 = 1$

$\alpha + 1 = 5 \Rightarrow \alpha = 4$

Checking: $4^2 \neq 1$

Hence no real value satisfies both conditions.

Answer: D) No real values

2. Given:$A = \begin{bmatrix}0 & c & -b\\ -c & 0 & a\\ b & -a & 0\end{bmatrix}$

and $B = \begin{bmatrix}a^2 & ab & ac\\ ab & b^2 & bc\\ ac & bc & c^2\end{bmatrix}$

We need to find AB.

Multiply A and B.

First row of AB:

$[0, c, -b]\begin{bmatrix}a^2\\ ab\\ ac\end{bmatrix} = 0 + c(ab) - b(ac) = abc - abc = 0$

Similarly, all remaining entries become 0.

Hence, $AB = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$

Therefore, $AB = O$

Answer: C) O

3. Given: $[1 \ x \ 1]\begin{bmatrix} 1 & 3 & 2 \\ 0 & 5 & 1 \\ 0 & 3 & 2 \end{bmatrix}\begin{bmatrix} 1 \\ 1 \\ x \end{bmatrix} = 0$

First multiply the matrix and column vector:

$$\begin{bmatrix} 1 & 3 & 2 \\ 0 & 5 & 1 \\ 0 & 3 & 2 \end{bmatrix}\begin{bmatrix} 1 \\ 1 \\ x \end{bmatrix} = \begin{bmatrix} 1+3+2x \\ 5+x \\ 3+2x \end{bmatrix} = \begin{bmatrix} 4+2x \\ 5+x \\ 3+2x \end{bmatrix}$$

Now multiply by $[1 \ x \ 1]$:

$$[1 \ x \ 1]\begin{bmatrix} 4+2x \\ 5+x \\ 3+2x \end{bmatrix} = 0$$

$$(4+2x) + x(5+x) + (3+2x) = 0$$

$$4 + 2x + 5x + x^2 + 3 + 2x = 0$$

$$x^2 + 9x + 7 = 0$$

Using quadratic formula:

$$x = \frac{-9 \pm \sqrt{81 - 28}}{2}$$

$$x = \frac{-9 \pm \sqrt{53}}{2}$$

Answer: D) $\frac{-9 \pm \sqrt{53}}{2}$

4. Given: $A = \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix}, B = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, C = \begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix}$

We need to find A^2, B^2, C^2.

First compute A^2:

$$A^2 = \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix}\begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix} = \begin{bmatrix} i^2 & 0 \\ 0 & (-i)^2 \end{bmatrix}$$

Since $i^2 = -1$,

$$A^2 = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} = -I$$

Now compute B^2:

$$B^2 = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} = -I$$

Now compute C^2:

$$C^2 = \begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix}\begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix} = \begin{bmatrix} i^2 & 0 \\ 0 & i^2 \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} = -I$$

Therefore, $A^2 = B^2 = C^2 = -I$

Answer: A) $A^2 = B^2 = C^2 = -I$

5. Given:

$$A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$$

We need to find: $A^3 - 4A^2 - 6A$

Observe that

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}$$

Using direct multiplication,

$$A^2 = \begin{bmatrix} 9 & 8 & 8 \\ 8 & 9 & 8 \\ 8 & 8 & 9 \end{bmatrix}$$

and

$$A^3 = \begin{bmatrix} 41 & 40 & 40 \\ 40 & 41 & 40 \\ 40 & 40 & 41 \end{bmatrix}$$

Now compute:

$$A^3 - 4A^2 - 6A$$
$$= \begin{bmatrix} 41 & 40 & 40 \\ 40 & 41 & 40 \\ 40 & 40 & 41 \end{bmatrix} - \begin{bmatrix} 36 & 32 & 32 \\ 32 & 36 & 32 \\ 32 & 32 & 36 \end{bmatrix} - \begin{bmatrix} 6 & 12 & 12 \\ 12 & 6 & 12 \\ 12 & 12 & 6 \end{bmatrix}$$
$$= \begin{bmatrix} -1 & -4 & -4 \\ -4 & -1 & -4 \\ -4 & -4 & -1 \end{bmatrix} = -A$$

Answer: C) $-A$

6. Given: $A = \begin{bmatrix} a^2 & ab & ac \\ ab & b^2 & bc \\ ac & bc & c^2 \end{bmatrix}$

and $a^2 + b^2 + c^2 = 1$

We need to find A^2.

Multiply A by itself.

First element:

$$(a^2)(a^2) + (ab)(ab) + (ac)(ac)$$
$$= a^2(a^2 + b^2 + c^2)$$

Since $a^2 + b^2 + c^2 = 1$

we get $= a^2$

Similarly, every entry remains unchanged.

Hence, $A^2 = A$

Answer: B) A

7. Given: $3A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \\ x & 2 & y \end{bmatrix}$

and $AA' = I$

Since $AA' = I$, rows of A are orthonormal.

So rows of $3A$ are mutually orthogonal and each row has length 3.

Let the rows of $3A$ be:

$R_1 = (1,2,2), R_2 = (2,1,-2), R_3 = (x,2,y)$
Using orthogonality with R_1:

$R_1 \cdot R_3 = 0$
$1(x) + 2(2) + 2(y) = 0$
$x + 4 + 2y = 0$
$x + 2y = -4 \quad (1)$
Using orthogonality with R_2:

$R_2 \cdot R_3 = 0$
$2x + 1(2) - 2y = 0$
$2x - 2y = -2$
$x - y = -1 \quad (2)$
From (2): $x = y - 1$
Substitute into (1):

$(y - 1) + 2y = -4$
$3y = -3$
$y = -1$
$x = -2$
Therefore, $x + y = -2 + (-1) = -3$
Correct Answer: C) -3

8. Given:$A = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$
We need to find A^{n+1}.

First compute A^2:

$A^2 = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}\begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$
$A^2 = \begin{bmatrix} 2 & -2 \\ -2 & 2 \end{bmatrix} = 2A$
Now, $A^3 = A^2A = 2A^2 = 2(2A) = 2^2A$
Similarly, $A^4 = 2^3A$
Continuing the pattern,

$A^{n+1} = 2^nA$
Therefore,

$A^{n+1} = 2^n \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$
Answer:

C)$2^n \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$

9. Given:

$A = \begin{bmatrix} 4 & 1 & 0 \\ 1 & -2 & 2 \end{bmatrix}, B = \begin{bmatrix} 2 & 0 & -1 \\ 3 & 1 & 4 \end{bmatrix}, C = \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}$
and $(3B - 2A)C + 2X = 0$
First find:

$3B = \begin{bmatrix} 6 & 0 & -3 \\ 9 & 3 & 12 \end{bmatrix}$

$2A = \begin{bmatrix} 8 & 2 & 0 \\ 2 & -4 & 4 \end{bmatrix}$

$3B - 2A = \begin{bmatrix} -2 & -2 & -3 \\ 7 & 7 & 8 \end{bmatrix}$

Now multiply by C:

$$\begin{bmatrix} -2 & -2 & -3 \\ 7 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix} = \begin{bmatrix} -2-4+3 \\ 7+14-8 \end{bmatrix} = \begin{bmatrix} -3 \\ 13 \end{bmatrix}$$

So, $\begin{bmatrix} -3 \\ 13 \end{bmatrix} + 2X = 0$

$2X = \begin{bmatrix} 3 \\ -13 \end{bmatrix}$

$$X = \begin{bmatrix} \frac{3}{2} \\ -\frac{13}{2} \end{bmatrix}$$

Correct Answer: C) $\boxed{\begin{bmatrix} \frac{3}{2} \\ -\frac{13}{2} \end{bmatrix}}$

10. We are given

$A = \begin{bmatrix} \cos\alpha & \sin\alpha \\ -\sin\alpha & \cos\alpha \end{bmatrix}$

and need to find

$(2A)\left(\frac{1}{4}A^T\right)$

Step 1: Simplify the expression

Using scalar multiplication:

$$\left(2A\right)\left(\frac{1}{4}A^T\right) = \left(2 \times \frac{1}{4}\right)(AA^T) = \frac{1}{2}(AA^T)$$

Step 2: Find $A^T$and compute AA^T

$A^T = \begin{bmatrix} \cos\alpha & -\sin\alpha \\ \sin\alpha & \cos\alpha \end{bmatrix}$

Now,

$$AA^T = \begin{bmatrix} \cos\alpha & \sin\alpha \\ -\sin\alpha & \cos\alpha \end{bmatrix} \begin{bmatrix} \cos\alpha & -\sin\alpha \\ \sin\alpha & \cos\alpha \end{bmatrix}$$

$$= \begin{bmatrix} \cos^2\alpha + \sin^2\alpha & -\cos\alpha\sin\alpha + \sin\alpha\cos\alpha \\ -\sin\alpha\cos\alpha + \cos\alpha\sin\alpha & \sin^2\alpha + \cos^2\alpha \end{bmatrix}$$

Using $\sin^2\alpha + \cos^2\alpha = 1$

we get

$AA^T = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$

Step 3: Final Answer

$$\frac{1}{2}(AA^T) = \frac{1}{2}I = \begin{bmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{2} \end{bmatrix}$$

Correct Answer: B

One or More Than One Correct Answer Questions

11. Given $A = \begin{bmatrix} 0 & 2\beta & \gamma \\ \alpha & \beta & -\gamma \\ \alpha & -\beta & \gamma \end{bmatrix}$ is orthogonal.

For an orthogonal matrix, columns are orthonormal.

Columns are:

$C_1 = \begin{bmatrix} 0 \\ \alpha \\ \alpha \end{bmatrix}, C_2 = \begin{bmatrix} 2\beta \\ \beta \\ -\beta \end{bmatrix}, C_3 = \begin{bmatrix} \gamma \\ -\gamma \\ \gamma \end{bmatrix}$

Step 1: Use unit length condition

$| C_1 |^2 = 0^2 + \alpha^2 + \alpha^2 = 2\alpha^2 = 1$

$\alpha = \pm\frac{1}{\sqrt{2}}$

$| C_2 |^2 = (2\beta)^2 + \beta^2 + (-\beta)^2 = 6\beta^2 = 1$

$\beta = \pm\frac{1}{\sqrt{6}}$

Also, $\alpha^2 = \frac{1}{2}, 3\beta^2 = 3 \cdot \frac{1}{6} = \frac{1}{2}$

Hence $\alpha = \pm\sqrt{3}\,\beta$

Correct Answers: A, C, D

12. Given $A = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix}$

Write $A = I + N, N = \begin{bmatrix} 0 & 0 \\ 2 & 0 \end{bmatrix}$

and $N^2 = 0$

Using

$(I + N)^n = I + nN$

$A^{2012} = \begin{bmatrix} 1 & 0 \\ 4024 & 1 \end{bmatrix}$

Thus $a = 1, b = 0, c = 4024, d = 1$

Check options:

A)$a = d$

True

B) $a + b + c + d = 1 + 0 + 4024 + 1 = 4026$

True

C)$a^2 + b^2 + d^2 = 1 + 0 + 1 = 2$

True

D)$d = 2012$

False

Correct Answers: A, B, C

13. Given

$$[x \;\; 4 \;\; -1]\begin{bmatrix}2 & 1 & 0\\ 1 & 0 & 2\\ 0 & 2 & 4\end{bmatrix}\begin{bmatrix}x\\ 4\\ -1\end{bmatrix} = 0$$

First compute:

$$\begin{bmatrix}2 & 1 & 0\\ 1 & 0 & 2\\ 0 & 2 & 4\end{bmatrix}\begin{bmatrix}x\\ 4\\ -1\end{bmatrix} = \begin{bmatrix}2x+4\\ x-2\\ 4\end{bmatrix}$$

Now multiply:

$$[x \;\; 4 \;\; -1]\begin{bmatrix}2x+4\\ x-2\\ 4\end{bmatrix} = 0$$

$x(2x+4) + 4(x-2) - 4 = 0$

$2x^2 + 8x - 12 = 0$

$x^2 + 4x - 6 = 0$

$x = \frac{-4 \pm \sqrt{16+24}}{2} = \frac{-4 \pm \sqrt{40}}{2} = -2 \pm \sqrt{10}$

Correct Answers: B, C

Comprehension Type Questions

Passage – I:

14. Given $A = \begin{bmatrix}1\\ 0\\ -1\end{bmatrix}, B = [2 \;\; 3 \;\; 2]$

First find

$$B^T = \begin{bmatrix}2\\ 3\\ 2\end{bmatrix}$$

Then $A + B^T = \begin{bmatrix}1\\ 0\\ -1\end{bmatrix} + \begin{bmatrix}2\\ 3\\ 2\end{bmatrix} = \begin{bmatrix}3\\ 3\\ 1\end{bmatrix}$

Now transpose:

$(A + B^T)^T = [3 \;\; 3 \;\; 1]$

Correct Answer: B

15. Given $A + 2B = \begin{bmatrix}2 & -4\\ 1 & 6\end{bmatrix}$

and $A^T + B^T = \begin{bmatrix}1 & 2\\ 0 & -1\end{bmatrix}$

Using $(A+B)^T = A^T + B^T$

we get

$$A + B = \begin{bmatrix}1 & 0\\ 2 & -1\end{bmatrix}$$

Now,

$(A + 2B) - (A + B) = B$

$B = \begin{bmatrix}2 & -4\\ 1 & 6\end{bmatrix} - \begin{bmatrix}1 & 0\\ 2 & -1\end{bmatrix} = \begin{bmatrix}1 & -4\\ -1 & 7\end{bmatrix}$

Then $A = (A + B) - B$

$A = \begin{bmatrix} 1 & 0 \\ 2 & -1 \end{bmatrix} - \begin{bmatrix} 1 & -4 \\ -1 & 7 \end{bmatrix} = \begin{bmatrix} 0 & 4 \\ 3 & -8 \end{bmatrix}$

Correct Answer: A

16. Given $A = \begin{bmatrix} 1 & 5 & 3 \\ 2 & 4 & 0 \\ 3 & -1 & -5 \end{bmatrix}, B = \begin{bmatrix} 2 & -1 & 0 \\ 0 & -2 & 5 \\ 1 & 2 & 0 \end{bmatrix}$

First find

$B^T = \begin{bmatrix} 2 & 0 & 1 \\ -1 & -2 & 2 \\ 0 & 5 & 0 \end{bmatrix}$

Now,

$3A = \begin{bmatrix} 3 & 15 & 9 \\ 6 & 12 & 0 \\ 9 & -3 & -15 \end{bmatrix}$

and

$4B^T = \begin{bmatrix} 8 & 0 & 4 \\ -4 & -8 & 8 \\ 0 & 20 & 0 \end{bmatrix}$

Thus,

$3A - 4B^T = \begin{bmatrix} 3-8 & 15-0 & 9-4 \\ 6-(-4) & 12-(-8) & 0-8 \\ 9-0 & -3-20 & -15-0 \end{bmatrix}$

$= \begin{bmatrix} -5 & 15 & 5 \\ 10 & 20 & -8 \\ 9 & -23 & -15 \end{bmatrix}$

Correct Answer: A

Matching Type Questions

a) $\begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix}\begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$

So, $a \to S$

b) $\begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$

So, $b \to P$

c) $\begin{bmatrix} 1 & 4 \\ 2 & 9 \end{bmatrix}\begin{bmatrix} 9 & -4 \\ -2 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

So, $c \to T$

d) $\begin{bmatrix} 1 & 0 & 1 \end{bmatrix}\begin{bmatrix} 2 & 1 \\ 1 & 0 \\ 3 & 2 \end{bmatrix} = \begin{bmatrix} 5 & 3 \end{bmatrix}$

So, $d \to Q$

Correct Matching: $\boxed{a \to S, b \to P, c \to T, d \to Q}$

Integer Type Questions

18. $\begin{bmatrix} 3 & -2 \\ -1 & 2 \end{bmatrix}\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 11 \\ -5 \end{bmatrix}$

Equations:

$3x - 2y = 11$
$-x + 2y = -5$
Adding,

$2x = 6 \Rightarrow x = 3$
Substitute in second equation:

$-3 + 2y = -5$
$2y = -2 \Rightarrow y = -1$
Now,

$3x + 7y = 3(3) + 7(-1) = 9 - 7 = 2$
$\boxed{2}$

19.

$5A = \begin{bmatrix} 3 & -4 \\ 4 & x \end{bmatrix}$
$A = \frac{1}{5}\begin{bmatrix} 3 & -4 \\ 4 & x \end{bmatrix}$
Since $AA^T = I$
rows are orthonormal.

Using second row norm:

$\left(\frac{4}{5}\right)^2 + \left(\frac{x}{5}\right)^2 = 1$
$16 + x^2 = 25$
$x^2 = 9$
Using orthogonality:

$\frac{3}{5} \cdot \frac{4}{5} + \frac{-4}{5} \cdot \frac{x}{5} = 0$
$12 - 4x = 0$
$x = 3$
$\boxed{3}$

20. $\text{trace}(A) = 3$
Property:

$\text{trace}(kA) = k\,\text{trace}(A)$
$\text{trace}(3A) = 3 \times 3 = 9$
$\boxed{9}$

STUDENT PRACTICE QUESTIONS

Single Answer Questions

1. Given:

$A + B = \begin{bmatrix} 2 & 3 & 1 \\ 6 & -1 & 5 \end{bmatrix}, A - B = \begin{bmatrix} 2 & 1 & -1 \\ 0 & -1 & 3 \end{bmatrix}$
Using,

$B = \frac{(A + B) - (A - B)}{2}$

$B = \frac{1}{2}\begin{bmatrix}0 & 2 & 2\\ 6 & 0 & 2\end{bmatrix} = \begin{bmatrix}0 & 1 & 1\\ 3 & 0 & 1\end{bmatrix}$

$\boxed{\begin{bmatrix}0 & 1 & 1\\ 3 & 0 & 1\end{bmatrix}}$

Correct Answer: A

2. $A = \begin{bmatrix}ab & b^2\\ -a^2 & -ab\end{bmatrix}$

Now,

$A^2 = A \cdot A$

$= \begin{bmatrix}ab & b^2\\ -a^2 & -ab\end{bmatrix}\begin{bmatrix}ab & b^2\\ -a^2 & -ab\end{bmatrix}$

First element:

$(ab)(ab) + b^2(-a^2) = a^2b^2 - a^2b^2 = 0$

Similarly all entries become 0.

Hence,

$A^2 = \begin{bmatrix}0 & 0\\ 0 & 0\end{bmatrix}$

$\boxed{\begin{bmatrix}0 & 0\\ 0 & 0\end{bmatrix}}$

Correct Answer: C

3. Given:

The matrix is

$A = \begin{bmatrix}-1 & 0\\ 0 & 2\end{bmatrix}$

We need to find the value of $A^3 - A^2$.

Since $A$is a diagonal matrix, we can find its powers by simply raising its diagonal elements to the corresponding power.

Step 1: Calculate A^2

$A^2 = \begin{bmatrix}(-1)^2 & 0\\ 0 & 2^2\end{bmatrix} = \begin{bmatrix}1 & 0\\ 0 & 4\end{bmatrix}$

Step 2: Calculate A^3

$A^3 = \begin{bmatrix}(-1)^3 & 0\\ 0 & 2^3\end{bmatrix} = \begin{bmatrix}-1 & 0\\ 0 & 8\end{bmatrix}$

Step 3: Calculate $A^3 - A^2$

$A^3 - A^2 = \begin{bmatrix}-1 & 0\\ 0 & 8\end{bmatrix} - \begin{bmatrix}1 & 0\\ 0 & 4\end{bmatrix}$

$A^3 - A^2 = \begin{bmatrix}-1-1 & 0-0\\ 0-0 & 8-4\end{bmatrix} = \begin{bmatrix}-2 & 0\\ 0 & 4\end{bmatrix}$

Step 4: Check the options

Option A:

$2A = 2\begin{bmatrix}-1 & 0\\ 0 & 2\end{bmatrix} = \begin{bmatrix}-2 & 0\\ 0 & 4\end{bmatrix}$

This matches perfectly.

Correct Answer: A) $2A$

4. Identify the given matrix

The given matrix is:

$$A = \begin{bmatrix} -5 & -8 & 0 \\ 3 & 5 & 0 \\ 1 & 2 & -1 \end{bmatrix}$$

Recall the definitions of the given options

- Idempotent matrix: A matrix $A$is idempotent if $A^2 = A$.
- Involutory matrix: A matrix $A$is involutory if $A^2 = I$, where $I$is the identity matrix.
- Nilpotent matrix: A matrix $A$is nilpotent if $A^k = 0$for some positive integer k.
- Identity matrix: A square matrix having 1s on the principal diagonal and 0s elsewhere.

Calculate A^2

$$A^2 = \begin{bmatrix} -5 & -8 & 0 \\ 3 & 5 & 0 \\ 1 & 2 & -1 \end{bmatrix}\begin{bmatrix} -5 & -8 & 0 \\ 3 & 5 & 0 \\ 1 & 2 & -1 \end{bmatrix}$$

Now multiply rows by columns.

First row:

$(-5)(-5) + (-8)(3) + (0)(1) = 25 - 24 = 1$
$(-5)(-8) + (-8)(5) + (0)(2) = 40 - 40 = 0$
$(-5)(0) + (-8)(0) + (0)(-1) = 0$
Second row:

$(3)(-5) + (5)(3) + (0)(1) = -15 + 15 = 0$
$(3)(-8) + (5)(5) + (0)(2) = -24 + 25 = 1$
$(3)(0) + (5)(0) + (0)(-1) = 0$
Third row:

$(1)(-5) + (2)(3) + (-1)(1) = -5 + 6 - 1 = 0$
$(1)(-8) + (2)(5) + (-1)(2) = -8 + 10 - 2 = 0$
$(1)(0) + (2)(0) + (-1)(-1) = 1$
Therefore,

$$A^2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I$$

Conclusion

Since $A^2 = I$
the matrix A is an Involutory matrix.

Correct Answer: B) Involutory matrix

5. Find AB

Given

$$A = \begin{bmatrix} 1 & -1 \\ 3 & 4 \\ -1 & 5 \end{bmatrix}, B = \begin{bmatrix} -1 & 2 \\ 3 & 1 \end{bmatrix}$$

Multiply A and B:

$$AB = \begin{bmatrix} 1 & -1 \\ 3 & 4 \\ -1 & 5 \end{bmatrix} \begin{bmatrix} -1 & 2 \\ 3 & 1 \end{bmatrix}$$

Row 1:

$(1)(-1) + (-1)(3) = -1 - 3 = -4$
$(1)(2) + (-1)(1) = 2 - 1 = 1$
Row 2:

$(3)(-1) + (4)(3) = -3 + 12 = 9$
$(3)(2) + (4)(1) = 6 + 4 = 10$
Row 3:

$(-1)(-1) + (5)(3) = 1 + 15 = 16$
$(-1)(2) + (5)(1) = -2 + 5 = 3$
Therefore,

$$AB = \begin{bmatrix} -4 & 1 \\ 9 & 10 \\ 16 & 3 \end{bmatrix}$$

Correct Answer: A)

6. Find A^2

Given

$$A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$

Now calculate $A^2 = A \times A$:

$$A^2 = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$

First row:

$1(1) + 0(0) + 1(1) = 2$
$1(0) + 0(0) + 1(0) = 0$
$1(1) + 0(0) + 1(1) = 2$

Second row gives all zeros.

Third row:

$1(1) + 0(0) + 1(1) = 2$
$1(0) + 0(0) + 1(0) = 0$
$1(1) + 0(0) + 1(1) = 2$
Thus,

$$A^2 = \begin{bmatrix} 2 & 0 & 2 \\ 0 & 0 & 0 \\ 2 & 0 & 2 \end{bmatrix} = 2A$$

Correct Answer: C) $2A$

7. Given

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$$

We need to find $A^3 - 9A$.

First calculate A^2.

Since every row and column contains all 1s,

$$A^2 = \begin{bmatrix} 3 & 3 & 3 \\ 3 & 3 & 3 \\ 3 & 3 & 3 \end{bmatrix} = 3A$$

Now calculate A^3:

$A^3 = A^2A = (3A)A = 3A^2$
But $A^2 = 3A$, so

$A^3 = 3(3A) = 9A$
Therefore,

$A^3 - 9A = 9A - 9A = 0$

Correct Answer: C) 0

One or More Than One Correct Answer Questions

8. Given

$$A = \begin{bmatrix} 0 & 0 & -1 \\ 0 & -1 & 0 \\ -1 & 0 & 0 \end{bmatrix}$$

Check each statement.

A) Trace of A is -1

Trace$(A) = 0 + (-1) + 0 = -1$
So, A is correct.

B) A is an involutory matrix

Calculate A^2:

$$A^2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I$$

Since $A^2 = I$, A is involutory.

So, B is correct.

C) Trace of $A^{2013}$is 1

Since $A^2 = I$,

$A^{2013} = A^{2(1006)+1} = (A^2)^{1006}A = IA = A$
Thus,

$\text{Trace}(A^{2013}) = \text{Trace}(A) = -1$
So, C is false.

D) Trace of $A^{2012}$is 3

$A^{2012} = (A^2)^{1006} = I$
$\text{Trace}(I) = 1 + 1 + 1 = 3$
So, D is correct.

Correct Answers: A, B and D

9. Given $A = \begin{bmatrix} 5 & 7 & 3 \\ 1 & 5 & 2 \\ 3 & 2 & 1 \end{bmatrix}$
and $A^3 - kA^2 + \ell A = I$
Using Cayley–Hamilton theorem:

$A^3 - (\text{trace}A)A^2 + (\text{sum of principal minors})A - (\det A)I = 0$
Rearranging,

$A^3 - (\text{trace}A)A^2 + (\text{sum of principal minors})A = (\det A)I$
Comparing with

$A^3 - kA^2 + \ell A = I$
we need

$\det(A) = 1$
Now calculate values.

Trace of A

$5 + 5 + 1 = 11$
Hence,

$k = 11$
Sum of principal minors

$\left|\begin{matrix} 5 & 7 \\ 1 & 5 \end{matrix}\right| = 25 - 7 = 18$
$\left|\begin{matrix} 5 & 3 \\ 3 & 1 \end{matrix}\right| = 5 - 9 = -4$
$\left|\begin{matrix} 5 & 2 \\ 2 & 1 \end{matrix}\right| = 5 - 4 = 1$
Sum:

$18 + (-4) + 1 = 15$
Thus,

$\ell = 15$
Correct Answers: A and C

Comprehension Type Questions

Passage – I:

10. Find $A + B$

Given

$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}, B = \begin{bmatrix} 5 & 6 \\ 7 & 8 \end{bmatrix}$

Add corresponding elements:

$A + B = \begin{bmatrix} 1+5 & 2+6 \\ 3+7 & 4+8 \end{bmatrix} = \begin{bmatrix} 6 & 8 \\ 10 & 12 \end{bmatrix}$

Correct Answer: A)

11. Find the transpose

Given

$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$

Transpose is obtained by interchanging rows and columns.

$A^T = \begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$

Correct Answer: A)

12. Find $2A$

Given

$A = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$

Multiply each element by 2:

$2A = \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix}$

Correct Answer: B)

Matching Type Questions

13. (a) Let

$A = \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix}$

$A^2 = \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix}\begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix} = \begin{bmatrix} (4)(4)+(0)(0) & (4)(0)+(0)(4) \\ (0)(4)+(4)(0) & (0)(0)+(4)(4) \end{bmatrix} = \begin{bmatrix} 16 & 0 \\ 0 & 16 \end{bmatrix}$

This matches (iv).

(b) Let

$A = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$

$A^2 = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}\begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} = \begin{bmatrix} (2)(2)+(0)(0) & (2)(0)+(0)(2) \\ (0)(2)+(2)(0) & (0)(0)+(2)(2) \end{bmatrix} = \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix}$

This matches (i).

(c) Let

$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$

$A^2 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} (0)(0)+(1)(1) & (0)(1)+(1)(0) \\ (1)(0)+(0)(1) & (1)(1)+(0)(0) \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

This matches (ii).

(d) Let

$A = \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix}$

$A^2 = \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix}\begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix} = \begin{bmatrix} (3)(3)+(0)(0) & (3)(0)+(0)(3) \\ (0)(3)+(3)(0) & (0)(0)+(3)(3) \end{bmatrix} = \begin{bmatrix} 9 & 0 \\ 0 & 9 \end{bmatrix}$

This matches (iii).

Integer Type Questions

14. Given

$A = \begin{bmatrix} x & 1 \\ 1 & 0 \end{bmatrix}$

and $A^2 = I$

Compute A^2:

$A^2 = \begin{bmatrix} x & 1 \\ 1 & 0 \end{bmatrix}\begin{bmatrix} x & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} x^2+1 & x \\ x & 1 \end{bmatrix}$

Since

$A^2 = I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

Comparing corresponding entries:

$x = 0$

Hence, $\boxed{x = 0}$

15. Given

$A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 5 \\ x & 5 & 6 \end{bmatrix}$

And $A^T = A$

Since A is symmetric,

$a_{13} = a_{31}$

So, $3 = x$

Hence,

$\boxed{x = 3}$

6. Binomial Theorem Solutions

LEVEL-I

1. In the expansion of $(x+a)^n$, powers of $x$go from $x^n$to x^0.
Hence total number of terms $= n+1$.
Answer: B) $n+1$

2. The formula for binomial coefficient is
$^nC_r = \frac{n!}{(n-r)!r!}$
Answer: B) $\frac{n!}{(n-r)!r!}$

3. General term in $(x+a)^n$is
$T_{r+1} = \ ^nC_r x^{n-r}a^r$
Answer: C) $^nC_r x^{n-r}a^r$

4. If $n$is even, then there is one middle term in the expansion of $(a+b)^n$.
Middle term $= \left(\frac{n}{2}+1\right)^{th}$ term.
Answer: A) $\left(\frac{n}{2}+1\right)^{th}$ **term**

5. Expansion of $(x+1)^3$:
$(x+1)^3 = x^3+3x^2+3x+1$
There are 4 terms, so middle terms are 2nd and 3rd terms:
$3x^2$and $3x$.
Answer: D) Both (A) and (B)

6. Expansion of
$\left(x+\frac{1}{x}\right)^2 = x^2+2+\frac{1}{x^2}$
Term independent of $x$is 2.
Answer: B) 2

7. Expansion of $(1+x)^4$:
$1+4x+6x^2+4x^3+x^4$
Largest binomial coefficient is 6, which is $^4C_2$.
Answer: B) $^4C_2$

8. $(x+2y)^{12}(x-2y)^{12}$
Using identity:
$(a+b)^n(a-b)^n = (a^2-b^2)^n$
So,
$(x^2-4y^2)^{12}$
This expansion has powers from 0to 12, hence number of terms $= 12+1 = 13$.
Answer: A) 13

9. $(3x+5y)^{16}+(3x-5y)^{16}$
In the sum, odd-powered terms cancel and only even-powered terms remain.
Even powers from 0to 16:
0,2,4, …,16
Number of even powers:
$\frac{16}{2}+1 = 9$
Answer: C) 9

10. $\left(x^2+\frac{1}{2y}\right)^{49}-\left(x^2-\frac{1}{2y}\right)^{49}$

In subtraction, all even-powered terms cancel and only odd-powered terms remain.

Odd powers from 1to 49:

1,3,5, … ,49

Number of odd integers:

$\frac{49+1}{2}=25$

Answer: C) 25

11. The $(r+1)^{th}$term in $(a+b)^n$is:

$T_{r+1}= \ {}^nC_r a^{n-r}b^r$

For $(2+x)^{10}$, the 6th term means $r=5$:

$T_6= \ {}^{10}C_5(2)^{10-5}x^5$

$= \ {}^{10}C_5 2^5 x^5$

Answer: B) ${}^{10}C_5\ 2^5x^5$

12. $\left(x-\frac{1}{2x}\right)^{10}$

Since $n=10$is even, middle term is

$\left(\frac{10}{2}+1\right)=6^{th}$ term

General term:

$T_{r+1}= \ {}^{10}C_r x^{10-r}\left(\ -\frac{1}{2x}\right)^r$

For 6th term, $r=5$:

$T_6= \ {}^{10}C_5 x^5\left(\ -\frac{1}{2x}\right)^5$

$=252\left(\ -\frac{1}{32}\right)$

$=-\frac{63}{8}$

Answer: D) $-\frac{63}{8}$

13. Find term independent of $x$in

$\left(x^2-\frac{1}{x}\right)^9$

General term:

$T_{r+1}= \ {}^9C_r(x^2)^{9-r}\left(\ -\frac{1}{x}\right)^r$

Power of x:

$x^{18-2r-r}=x^{18-3r}$

For constant term:

$18-3r=0$

$r=6$

Constant term:

${}^9C_6=84$

Answer: A) 84

14. 7th term in

$\left(\frac{4}{x^3}+\frac{x^2}{2}\right)^{14}$

Using general term:

$T_{r+1}= \ {}^{14}C_r\left(\frac{4}{x^3}\right)^{14-r}\left(\frac{x^2}{2}\right)^r$

For 7th term, $r=6$:

$T_7 = \quad {}^{14}C_6 \left(\frac{4}{x^3}\right)^8 \left(\frac{x^2}{2}\right)^6$

$= \quad {}^{14}C_6 \frac{4^8 x^{12}}{x^{24} 2^6}$

$= \quad {}^{14}C_6 \frac{4^5}{x^{12}}$

Answer: B) ${}^{14}C_6 \frac{4^5}{x^{12}}$

15. The 4th term from the end in $(2a + 5b)^8$

4th term from end = $(8 - 4 + 2) = 6^{th}$term from beginning.

General term:

$T_{r+1} = \quad {}^nC_r a^{n-r} b^r$

For 6th term, $r = 5$:

$T_6 = \quad {}^8C_5 (2a)^3 (5b)^5$

$= \quad {}^8C_5 \cdot 2^3 \cdot 5^5 a^3 b^5$

Answer: B) ${}^8C_5\, 2^3 5^5 a^3 b^5$

16. $\left(\sqrt{x^5} + \frac{3}{\sqrt{x^3}}\right)^6$

Rewrite:

$\left(x^{5/2} + 3x^{-3/2}\right)^6$

General term:

$T_{r+1} = \quad {}^6C_r (x^{5/2})^{6-r} (3x^{-3/2})^r$

Power of x:

$\frac{5}{2}(6 - r) - \frac{3r}{2} = 15 - 4r$

For coefficient of x^3:

$15 - 4r = 3$

$r = 3$

Coefficient:

${}^6C_3 \cdot 3^3 = 20 \cdot 27 = 540$

Answer: D) 540

17. Coefficient of $x$in

$\left(x^2 + \frac{k}{x}\right)^5$

General term:

$T_{r+1} = \quad {}^5C_r (x^2)^{5-r} \left(\frac{k}{x}\right)^r$

Power of x:

$x^{10-2r-r} = x^{10-3r}$

For coefficient of x:

$10 - 3r = 1$

$r = 3$

Coefficient:

${}^5C_3 k^3 = 10k^3$

Given:

$10k^3 = 270$

$k^3 = 27$

$k = 3$

Answer: C) 3

18. In $(1+x)^{18}$, coefficients of $(2r+4)^{th}$and $(r-2)^{th}$terms are equal.
Coefficient of k^{th}term is: ${}^{18}C_{k-1}$
So, ${}^{18}C_{2r+3} = {}^{18}C_{r-3}$
Using property: ${}^{n}C_a = {}^{n}C_b \Rightarrow a+b=n$
$(2r+3)+(r-3)=18$
$3r=18$
$r=6$
Answer: C) 6

19. Fourth term in $\left(px+\frac{1}{x}\right)^n$
is $\frac{5}{2}$.
Fourth term means $r=3$: $T_4 = {}^{n}C_3(px)^{n-3}\left(\frac{1}{x}\right)^3$
For constant term: $n-3-3=0$
$n=6$
Then
$T_4 = {}^{6}C_3p^3$
$20p^3 = \frac{5}{2}$
$p^3 = \frac{1}{16}$
$p = \frac{1}{2}$
Hence,
$(n,p) = \left(6,\frac{1}{2}\right)$
Answer: B) $\left(6,\frac{1}{2}\right)$

20. Coefficients of 5th, 6th, and 7th terms in $(1+x)^n$are in A.P.
So,
$2\cdot {}^{n}C_5 = {}^{n}C_4 + {}^{n}C_6$
Using properties of binomial coefficients and simplifying:
$n^2-21n+98=0$
$(n-7)(n-14)=0$
$n=7$ or 14
Answer: C) 7 or 14

LEVEL-II

SINGLE OPTION CORRECT TYPE

1. In $(1+x)^{20}$,
Coefficient of r^{th}term:
${}^{20}C_{r-1}$
Coefficient of $(r+1)^{th}$term:
${}^{20}C_r$
Given ratio:
$\frac{{}^{20}C_{r-1}}{{}^{20}C_r} = \frac{1}{2}$
Using,
$\frac{{}^{n}C_{r-1}}{{}^{n}C_r} = \frac{r}{n-r+1}$
So,

$$\frac{r}{21-r}=\frac{1}{2}$$
$$2r=21-r$$
$$3r=21$$
$$r=7$$
Answer: C) 7

2. General term in
$$\left(x+\frac{1}{y}\right)^{23}$$
is
$$T_{r+1}= \ {}^{23}C_r x^{23-r}y^{-r}$$
Now,
$$T_{12}= \ {}^{23}C_{11}x^{12}y^{-11}$$
$$T_{13}= \ {}^{23}C_{12}x^{11}y^{-12}$$
Since,
$${}^{23}C_{11}= \ {}^{23}C_{12}$$
we get
$$T_{12}=xy\ T_{13}$$
Answer: C) $T_{12}=xyT_{13}$

3. Constant term in
$$\left(\sqrt{x}-\frac{k}{x^2}\right)^{10}$$
General term:
$$T_{r+1}= \ {}^{10}C_r(x^{1/2})^{10-r}(\ \ -kx^{-2})^r$$
Power of x:
$$\frac{10-r}{2}-2r=5-\frac{5r}{2}$$
For constant term:
$$5-\frac{5r}{2}=0$$
$$r=2$$
Constant term:
$${}^{10}C_2k^2=45k^2$$
Given:
$$45k^2=405$$
$$k^2=9$$
$$k=\pm 3$$
Answer: C) ± 3

4. Put $x=i$in
$$(1+x)^{10}=a_0+a_1x+\cdots+a_{10}x^{10}$$
Then
$$(1+i)^{10}=(a_0-a_2+a_4-a_6+a_8-a_{10})+i(a_1-a_3+a_5-a_7+a_9)$$
Taking modulus squared:
$$(a_0-a_2+a_4-a_6+a_8-a_{10})^2+(a_1-a_3+a_5-a_7+a_9)^2=|\ 1+i\ |^{20}$$
Since
$$|\ 1+i\ |=\sqrt{2}$$
we get
$$\left(\sqrt{2}\right)^{20}=2^{10}$$
Answer: B) 2^{10}

5. Let $C_r = \ {}^nC_r$.
We need:
$C_0 + 2C_1 + 3C_2 + \cdots + (n+1)C_n$
This equals
$\sum_{r=0}^{n} (r+1) \ {}^nC_r$
Using identities:
$\sum \ {}^nC_r = 2^n$
and
$\sum r \ {}^nC_r = n2^{n-1}$
Hence,
$2^n + n2^{n-1} = (n+2)2^{n-1}$
Answer: D) $(n+2)2^{n-1}$

6. Evaluate
$3C_1 - 4C_2 + 5C_3 - \cdots + (-1)^{n-1}(n+2)C_n$
where $C_r = \ {}^nC_r$.
Write as
$\sum_{r=1}^{n} (-1)^{r-1}(r+2) \ {}^nC_r$
Using standard binomial identities:
$\sum(-1)^r \ {}^nC_r = 0$
and
$\sum(-1)^r r \ {}^nC_r = 0$
After simplification, value becomes
2
Answer: B) 2

7. Evaluate
$\frac{C_1}{C_0} + 2\frac{C_2}{C_1} + 3\frac{C_3}{C_2} + \cdots + n\frac{C_n}{C_{n-1}}$
Using
$\frac{{}^nC_r}{{}^nC_{r-1}} = \frac{n-r+1}{r}$
Therefore,
$r \cdot \frac{C_r}{C_{r-1}} = n - r + 1$
So sum becomes
$n + (n-1) + \cdots + 1$
$= \frac{n(n+1)}{2}$
Answer: B) $\frac{n(n+1)}{2}$

8. Evaluate
$C_0^2 + 3C_1^2 + 5C_2^2 + \cdots + (2n+1)C_n^2$
where $C_r = \ {}^nC_r$.
Using standard identity:
$\sum_{r=0}^{n} (2r+1)\,(\ {}^nC_r)^2 = (n+1) \ {}^{2n}C_n$
Answer: C) $(n+1) \ {}^{2n}C_n$

9. Given

$$\left(x-2\right)^{100}=\sum_{r=0}^{100}a_r\,x^r$$

We need:

$a_1+2a_2+3a_3+\cdots+100a_{100}$

Let

$$f(x)=\sum_{r=0}^{100}a_r\,x^r$$

Then

$$f'(x)=\sum_{r=1}^{100}r\,a_r x^{r-1}$$

Putting $x=1$:

$f'(1)=a_1+2a_2+\cdots+100a_{100}$

Now,

$f(x)=(x-2)^{100}$

$f'(x)=100(x-2)^{99}$

Hence,

$f'(1)=100(1-2)^{99}$

$=100(-1)^{99}=-100$

Answer: B) -100

10. We use the standard identity:

$\sum_{r=0}^{n}(\ {}^nC_r)^2=\ {}^{2n}C_n$

Therefore,

$$\sum_{r=0}^{n}(\ {}^nC_r)^2=\ {}^{2n}C_n$$

$$=\ {}^{2n}C_n=\frac{(2n)!}{(n!)^2}$$

Answer: D) $\frac{(2n)!}{(n!)^2}$

ONE OR MORE THAN ONE OPTION CORRECT TYPE

11. Coefficient of x^2 in $(1-x)^2$

$(1-x)^2=1-2x+x^2$

Coefficient of $x^2=1$.

Also,

$$\frac{n^2+3n+2}{(n+2)(n+1)}=\frac{(n+1)(n+2)}{(n+2)(n+1)}=1$$

Hence both B and C are correct.

Answers: B, C

12. Expression:

$$\left(x^3+3\cdot 2^{-\log_{\sqrt{2}}\sqrt{x^3}}\right)^{11}$$

Simplify the second term.

Since

$\log_{\sqrt{2}}\sqrt{x^3}=\log_{2^{1/2}}x^{3/2}=3\log_2 x$

Thus,

$2^{-3\log_2 x} = x^{-3}$

So expression becomes

$(x^3 + 3x^{-3})^{11}$

General term:

$T_{r+1} = {}^{11}C_r (x^3)^{11-r} (3x^{-3})^r$

Power of x:

$33 - 3r - 3r = 33 - 6r$

Possible powers:

$33,27,21,15,9,3,-3,-9,\ldots$

So x^2never appears, but x^{-3}appears when $r = 6$.

Coefficient of x^3:

For $33 - 6r = 3$,

$r = 5$

Coefficient:

${}^{11}C_5 3^5$

Coefficient of x^{-3}:

For $33 - 6r = -3$,

$r = 6$

Coefficient:

${}^{11}C_6 3^6$

Their ratio:

$$\frac{{}^{11}C_5 3^5}{{}^{11}C_6 3^6} = \frac{1}{3}$$

since ${}^{11}C_5 = {}^{11}C_6$.

Hence B, C, and D are correct.

Answers: B, C, D

13. Sum of coefficients in

$(1 + 3x - 5x^2)^{2n}$

is obtained by putting $x = 1$:

$(1 + 3 - 5)^{2n} = (-1)^{2n} = 1$

Answer: A) 1

COMPREHENSION TYPE

Passage – I:

14. Second term

General term:

$T_{r+1} = {}^{7}C_r m^{7-r} (-n)^r$

For 2nd term, $r = 1$:

$T_2 = {}^{7}C_1 m^6 (-n)$

$= -7m^6 n$

Answer: B) $-7m^6 n$

15. Third term

For 3rd term, $r = 2$:

$T_3 = {}^{7}C_2 m^5 (-n)^2$

$= 21m^5 n^2$

Answer: A) $21m^5 n^2$

16. Fourth term

For 4th term, $r = 3$:

$T_4 = {}^7C_3 m^4(-n)^3$

$= -35m^4n^3$

Answer: B) $-35m^4n^3$

17. Matrix Matching

Given

$(1 + x + x^2)^n = a_0 + a_1x + a_2x^2 + \cdots + a_{2n}x^{2n}$

(a) $\boldsymbol{a_0 + a_1 + a_2 + \cdots + a_{2n}}$

Put $x = 1$:

$(1 + 1 + 1)^n = 3^n$

So,

$a_0 + a_1 + \cdots + a_{2n} = 3^n$

Matches with **P**

(b) $\boldsymbol{a_0 + a_2 + a_4 + \cdots + a_{2n}}$

Even-power sum:

$\frac{f(1) + f(-1)}{2}$

where

$f(x) = (1 + x + x^2)^n$

Now,

$f(1) = 3^n, f(-1) = 1$

Thus,

$\frac{3^n + 1}{2}$

Matches with **R**

(c) $\boldsymbol{a_1 + a_3 + a_5 + \cdots + a_{2n-1}}$

Odd-power sum:

$\frac{f(1) - f(-1)}{2}$

$= \frac{3^n - 1}{2}$

Matches with **Q**

(d) $\boldsymbol{a_0 + a_3 + a_6 + \cdots}$

Using cube roots of unity property:

$a_0 + a_3 + a_6 + \cdots = 3^{n-1}$

Matches with **S**

Final Matching

- **a → P**
- **b → R**
- **c → Q**
- **d → S**

INTEGER TYPE

18. Given:

T_{r+1} is independent of x in $\left(3x - \frac{5}{x^3}\right)^8$

Using general term:

$T_{r+1} = \binom{8}{r}(3x)^{8-r}\left(-\frac{5}{x^3}\right)^r$

Power of x:

$x^{8-r} \cdot x^{-3r} = x^{8-4r}$

For term independent of x,

$8 - 4r = 0$

$r = 2$

Answer: 2

19. Expansion of:

$(3 - 2x)^5$

At $x = 1$:

$(3 - 2)^5 = 1$

General term:

$T_{r+1} = \binom{5}{r} 3^{5-r}(-2)^r$

Numerical values:

$T_1 = 243$

$T_2 = -810$

$T_3 = 1080$

$T_4 = -720$

$T_5 = 240$

$T_6 = -32$

Greatest numerically means greatest absolute value.

Largest magnitude is 1080, which is the **3rd term**.

Answer: 3

20. Given:

$$\sum_{i=1}^{n} \sum_{j=1}^{n} \sum_{k=1}^{n} 1$$

Each summation contributes $n$terms:

$n \times n \times n = n^3$

So volume $= k = n^3$.

For $n \neq 0$, taking $n = 1$gives:

$k = 1$

Sum of digits of k:

1

Answer: 1

Student practice corner

Single option correct type

1. Given:

$\left(3 + 2x\right)^{15}, x = \frac{5}{2}$

General term:

$T_{r+1} = \binom{15}{r} 3^{15-r}(2x)^r$

Substitute $x = \frac{5}{2}$:

$2x = 5$

So,

$T_{r+1} = \binom{15}{r} 3^{15-r} 5^r$

For numerically greatest term:

$\frac{T_{r+2}}{T_{r+1}} = \frac{15-r}{r+1} \cdot \frac{5}{3}$

Set

$\frac{T_{r+2}}{T_{r+1}} \geq 1$

$\frac{15-r}{r+1} \cdot \frac{5}{3} \geq 1$

$5(15-r) \geq 3(r+1)$

$75 - 5r \geq 3r + 3$

$72 \geq 8r$

$r \leq 9$

Greatest term corresponds to $r = 9$.

Hence term is:

T_{10}

Answer: 10th term (Option C)

2. Using binomial expansion:

$\left(x+\frac{2}{x}\right)^2 = x^2 + 2(x)\left(\frac{2}{x}\right) + \left(\frac{2}{x}\right)^2$

$= x^2 + 4 + \frac{4}{x^2}$

The term independent of $x$is 4.

Answer: A) 4

3. The binomial coefficients are:

$^6C_0, \quad ^6C_1, \quad ^6C_2, \quad ^6C_3, \quad ^6C_4, \quad ^6C_5, \quad ^6C_6$

Their values are:

1, 6, 15, 20, 15, 6, 1

Largest coefficient = 20

$20 = \quad ^6C_3$

Answer: D) 20

4. For $(a+b)^n$, the number of terms is:

Number of terms = $n + 1$

Here $n = 8$,

$8 + 1 = 9$

Answer: C) 9

5. Sum of coefficients in:

$(1-x)^{10}$

Put $x = 1$:

$(1-1)^{10} = 0$

Answer: 0(Option D)

6. Using binomial theorem:

$\sum_{r=0}^{n} \quad ^nC_r = (1+1)^n = 2^n$

So,

$^nC_0 + \quad ^nC_1 + \quad ^nC_2 + \cdots + \quad ^nC_n = 2^n$

Answer: 2^n(Option C)

7. We know:

$^nC_0 + \quad ^nC_1 + \quad ^nC_2 + \cdots + \quad ^nC_n = 2^n$

Also,

$^nC_0 + \ ^nC_2 + \ ^nC_4 + \cdots = \ ^nC_1 + \ ^nC_3 + \ ^nC_5 + \cdots = 2^{n-1}$

Given:

$^nC_2 + \ ^nC_4 + \ ^nC_6 + \cdots$

Subtract $^nC_0 = 1$:

$2^{n-1} - 1$

Answer: $2^{n-1} - 1$(Option A)

8. For the expansion of $(a+b)^n$:

Total number of terms $= n + 1$

Here $n = 8$,

$8 + 1 = 9$ terms

Since the number of terms is odd, there is one middle term.

Middle term position:

$\frac{9+1}{2} = 5$

Hence, the middle term is the **5th term**.

Answer: B) 5th term

ONE OR MORE THAN ONE OPTION CORRECT TYPE

9. In the expansion of $(1+x)^8$, the greatest binomial coefficient(s) is/are

The binomial coefficients are:

$^8C_0, \ ^8C_1, \ ^8C_2, \ ^8C_3, \ ^8C_4, \ ^8C_5, \ ^8C_6, \ ^8C_7, \ ^8C_8$

Their values are:

1, 8, 28, 56, 70, 56, 28, 8, 1

The greatest coefficient is:

$^8C_4 = 70$

So correct options are:

- A) $^8C_4$✓
- B) 70✓

Correct Answers: A, B

10. In the expansion of $(a+b)^7$

For $(a+b)^n$:

Number of terms $= n + 1$

Here $n = 7$,

$7 + 1 = 8$

So, A is correct.

Since the number of terms is even, there are two middle terms:

$\frac{8}{2} = 4^{th}, \frac{8}{2} + 1 = 5^{th}$

So, B is correct.

For odd n, the two greatest coefficients are equal:

$^7C_3 = \ ^7C_4 = 35$

So, C is correct.

Last term of $(a+b)^7$is:

b^7

So, D is correct.

Correct Answers: A, B, C, D

COMPREHENSION TYPE

Passage – I:

11. Given:

Sum of odd terms = P

Sum of even terms = Q

For expansion of: $(x+a)^n$

We know: $P+Q=(x+a)^n$

And $P-Q=(x-a)^n$

Therefore, $P^2-Q^2=(P+Q)(P-Q)$

$=(x+a)^n(x-a)^n$

$=(x^2-a^2)^n$

Answer: $\boxed{(x^2-a^2)^n}$

(Option B)

12. Using: $P+Q=(x+a)^n$

$P-Q=(x-a)^n$

Now, $(P+Q)^2-(P-Q)^2=4PQ$

So, $4PQ=(x+a)^{2n}-(x-a)^{2n}$

Answer: $\boxed{(x+a)^{2n}-(x-a)^{2n}}$

(Option A)

13. Match the Following

Solution

Greatest coefficient in $(1+x)^8$

$^8C_4 = 70$

So, a → Q

Number of terms in $(a+b)^8$

$8+1=9$

So, b → P

Middle term in $(x+y)^6$

Since total terms = 7, middle term is 4th term:

$T_4 = {}^6C_3x^3y^3 = 20x^3y^3$

So, c → R

Coefficient of x^3 **in** $(1+x)^6$

$^6C_3 = 20$

So, d → S

Correct Answer:

a–Q, b–P, c–R, d–S

INTEGER TYPE

14.

Given: $\left(x^2+\frac{k}{x}\right)^5$

General term: $T_{r+1} = {}^5C_r(x^2)^{5-r}\left(\frac{k}{x}\right)^r$

$= {}^5C_rk^rx^{10-3r}$

Coefficient of $x$is 270.

So power of $x$must be 1: $10-3r=1$

$3r=9$

$r=3$

Thus required term: $T_4 = \quad {}^5C_3k^3x$

Coefficient: $\quad {}^5C_3k^3 = 270$

$10k^3 = 270$

$k^3 = 27$

$k = 3$

Answer: $\boxed{3}$

15.

Given successive coefficients: 28, 56, 70

These are: $\quad {}^nC_r = 28, \quad {}^nC_{r+1} = 56, \quad {}^nC_{r+2} = 70$

Using:

$$\frac{{}^nC_{r+1}}{{}^nC_r} = \frac{n-r}{r+1}$$

$$\frac{56}{28} = 2$$

$$\frac{n-r}{r+1} = 2$$

$n - r = 2r + 2$

$n = 3r + 2$

Again, $\frac{{}^nC_{r+2}}{{}^nC_{r+1}} = \frac{n-r-1}{r+2}$

$$\frac{70}{56} = \frac{5}{4}$$

$$\frac{n-r-1}{r+2} = \frac{5}{4}$$

$4n - 4r - 4 = 5r + 10$

$4n = 9r + 14$

Substitute $n = 3r + 2$: $12r + 8 = 9r + 14$

$3r = 6$

$r = 2$

Hence, $n = 3(2) + 2 = 8$

Answer: $\boxed{8}$

7. Sets and Relations

LEVEL – I

Single option correct type

1. Let A and B be two sets in the same universal set. Then, $A - B =$
$A - B = \{x: x \in A \text{ and } x \notin B\} = A \cap B'$
Correct Answer: C) $A \cap B'$

2. The number of subsets of a set containing $n$elements is
A set with $n$elements has:
2^n
subsets.
Correct Answer: D) 2^n

3. For any two sets A and B,
$A \cap (A \cup B)$
Using the absorption law:
$A \cap (A \cup B) = A$
Correct Answer: A) A

4. For any two sets $A$and B,
$(A - B) \cup (B - A)$
This represents elements belonging to exactly one of the sets, i.e. symmetric difference:
$(A \cup B) - (A \cap B)$
Correct Answer: C) $(A \cup B) - (A \cap B)$

5. Which of the following statement is false?
Check option C:
$A - B = A - B'$
But, $A - B = A \cap B'$
and $A - B' = A \cap B$
These are generally not equal.
`Hence, option C is false.
Correct Answer: C) $A - B = A - B'$

6. Let $A = \{x: x \in \mathbb{R}, x > 4\}, B = \{x: x \in \mathbb{R}: x < 5\}$
Then, $A \cap B = \{x: 4 < x < 5\} = (4,5)$
Correct Answer: B) $(4, 5)$

7. Let $n(A) = 16$, $n(B) = 14$, $n(A \cup B) = 25$
Using the formula:
$n(A \cup B) = n(A) + n(B) - n(A \cap B)$
$25 = 16 + 14 - n(A \cap B)$
$25 = 30 - n(A \cap B)$
$n(A \cap B) = 5$
Correct Answer: C) 5

8. If $A = \{1,2,3,4,5\}$
Number of subsets: $2^5 = 32$
Proper subsets exclude the set itself:
$32 - 1 = 31$
Correct Answer: C) 31

9. For two sets, $A \cup B = A$
This happens when every element of B is already in A.

$B \subseteq A$

Correct Answer: A) $B \subseteq A$

10. If A and B are two given sets, then

$A \cap (A \cap B)^c$

Using De Morgan's law:

$(A \cap B)^c = A^c \cup B^c$

So,

$A \cap (A^c \cup B^c)$

$= (A \cap A^c) \cup (A \cap B^c)$

$= \phi \cup (A \cap B^c)$

$= A \cap B^c$

Correct Answer: D) $A \cap B^c$

11. If $A = \{x: x$ is a multiple of $3\}, B = \{x: x$ is a multiple of $5\}$

$A - B$ means elements in A but not in B.

$A - B = A \cap \bar{B}$

Correct Answer: B) $A \cap \bar{B}$

12. In a city, 20%travel by car, 50%by bus and 10%by both.

Using:

$n(A \cup B) = n(A) + n(B) - n(A \cap B)$

$20 + 50 - 10 = 60$

So, persons travelling by car or bus:

60%

Correct Answer: C) 60%

13. Two finite sets have $m$and $n$elements.

Number of elements in power sets:

$2^m, 2^n$

Given:

$2^m = 2^n + 48$

Check options:

For $m = 6$, $n = 4$,

$2^6 - 2^4 = 64 - 16 = 48$

Hence correct.

Correct Answer: C) 6, 4

14. $A = \{1,2,3\}, B = \{2,3,4\}$

Given: $C \subset B$

and C is disjoint with A.

Only element in B not in A is 4.

So, $C = \{4\}$

Correct Answer: D) $\{4\}$

15. If $n(A \cup B) = n(A) + n(B)$

Using the formula: $n(A \cup B) = n(A) + n(B) - n(A \cap B)$

Comparing, $n(A \cap B) = 0$

So, A and B are disjoint.

Correct Answer: B) disjoint sets

16. $A = \{1,2,3\}$

Number of elements in the power set:

$2^n = 2^3 = 8$

Correct Answer: A) 8

17. $U = \{1,2,3,4\}, A = \{1,2\}, B = \{2,3\}$
First find: $A \cup B = \{1,2,3\}$
Complement in U: $(A \cup B)' = \{4\}$
Hence, $n((A \cup B)') = 1$
Correct Answer: A) 1

18. Let S contain n elements.
A binary operation on S is a function:
$S \times S \to S$
Number of ordered pairs in $S \times S$:
n^2
Each pair can map to any of $n$elements.
So total binary operations:
n^{n^2}
Correct Answer: C) n^{n^2}

19. $A = \{x: x$ is a multiple of 3$\}$
$B = \{x: x$ is a multiple of 5$\}$
$A \cap B$ means numbers divisible by both 3 and 5, i.e. multiples of 15.
$A \cap B = \{15,30,45, \dots\}$
Correct Answer: C) $\{\mathbf{15}, \mathbf{30}, \mathbf{45}, \dots\}$

20. Let $U = A \cup B \cup C$
We need to find:$\{(A - B) \cup (B - C) \cup (C - A)\}'$
Write differences as intersections:
$A - B = A \cap B'$
$B - C = B \cap C'$
$C - A = C \cap A'$
The union contains elements belonging to one set but not the next.
Its complement contains elements common to all three sets.
Hence,
$\{(A - B) \cup (B - C) \cup (C - A)\}' = A \cap B \cap C$
Correct Answer: C) $A \cap B \cap C$

LEVEL-2

1. For any three sets A, B, C, which statement is false?
Using: $B - C = B \cap C'$
Then,$A \cup (B - C) = A \cup (B \cap C')$
By distributive law: $= (A \cup B) \cap (A \cup C')$
So option C is true.
Now check option D:
$(A \cup B) - (A \cup C)$
$= (A \cup B) \cap (A \cup C)'$
This is not equal to $A \cup (B - C)$.
Hence option D is false.
Correct Answer: D)

2. Let $n(U) = 700$, $n(A) = 200$, $n(B) = 300$, $n(A \cap B) = 100$
First find:
$n(A \cup B) = n(A) + n(B) - n(A \cap B)$
$= 200 + 300 - 100 = 400$

Now, $A' \cap B' = (A \cup B)'$
So, $n(A' \cap B') = 700 - 400 = 300$
Correct Answer: C) 300

3. Total students = 100
Given:
All three = 10
Milk & coffee = 20
Milk & tea = 25
Tea & coffee = 10
Milk only = 12
Coffee only = 5
Tea only = 8
Students taking exactly two drinks:
$(20 - 10) + (25 - 10) + (10 - 10)$
$= 10 + 15 + 0 = 25$
Students taking at least one drink:
$12 + 5 + 8 + 25 + 10$
$= 60$
Therefore, students taking none:
$100 - 60 = 40$
Correct Answer: C) 40

4. Let
$n(M) = 100, n(P) = 70, n(C) = 40$
$n(M \cap P) = 30, n(M \cap C) = 28, n(P \cap C) = 23$
$n(M \cap P \cap C) = 18$
Mathematics only:
$= n(M) - n(M \cap P) - n(M \cap C) + n(M \cap P \cap C)$
$= 100 - 30 - 28 + 18$
$= 60$
Correct Answer: C) 60

5. Which of the following is the empty set?
Check each option:
A)$x^2 - 1 = 0 \Rightarrow x = \pm 1$
Not empty.
B)$x^2 + 1 = 0$
No real solution.
So this set is empty.
Correct Answer: B)

6. Let $R = \{(1,1), (2,2), (3,3), (1,2), (2,3), (1,3)\}$on $A = \{1,2,3\}$. Then $R$is
Reflexive: Since $(1,1), (2,2), (3,3) \in R$, relation is reflexive.
Symmetric: $(1,2) \in R$but $(2,1) \notin R$, so not symmetric.
Transitive:
$(1, 2)$and $(2,3) \Rightarrow (1,3) \in R$.
Other required pairs also satisfy transitivity.
Therefore, $R$is transitive.
Correct Answer: B) Transitive

7. The shaded region in the figure is
The shaded portion represents only the part of set Aexcluding the common regions with $B$and C.
Hence the region is:
$A - (B \cup C)$
Correct Answer: D) $\boldsymbol{A - (B \cup C)}$

8. If A and B are two sets, then $(A - B) \cup (B - A) \cup (A \cap B)$ is equal to

- $A - B$: elements only in A
- $B - A$: elements only in B
- $A \cap B$: common elements

Their union gives all elements belonging to either Aor B.
$(A - B) \cup (B - A) \cup (A \cap B) = A \cup B$
Correct Answer: A) $\boldsymbol{A \cup B}$

9. Since 65% own neither,
$P(\text{car} \cup \text{phone}) = 100\% - 65\% = 35\%$
So statement 2 is correct.
Using inclusion-exclusion:
$35\% = 25\% + 15\% - P(\text{both})$
$P(\text{both}) = 40\% - 35\% = 5\%$
So statement 1 is false.
Given 5% corresponds to 2000 families,
$0.05 \times \text{Total families} = 2000$
$\text{Total families} = \frac{2000}{0.05} = 40000$
So statement 3 is correct.
Correct Answer: C) 2 and 3

10. Let
$A = \{(x, y): y = e^x,\ x \in \mathbb{R}\}$
$B = \{(x, y): y = e^{-x},\ x \in \mathbb{R}\}$
To find intersection, equate:
$e^x = e^{-x}$
$e^{2x} = 1$
$2x = 0 \Rightarrow x = 0$
Then $y = e^0 = 1$
So the common point is $(0, 1)$.
Hence,$A \cap B \neq \phi$
Correct Answer: B) $\boldsymbol{A \cap B \neq \phi}$

ONE OR MORE THAN ONE CORRECT ANSWER TYPE

11. Singleton sets:
A singleton set has exactly one element.
A) $\{0\}$→ one element
B) $\{x: x$ is an even prime number$\}$→ only 2
C) $\{1, 2\}$→ two elements
D) $\{a\}$→ one element
Correct Answers: A, B, D

12. True statements regarding complements:
A) $A \cup A' = U$
B) $A \cap A' = \emptyset$
C) Complement of universal set is universal set
($U' = \emptyset$)
D) Complement of $\emptyset$is U
Correct Answers: A, B, D

13. Given $A = \{1,2,3\}$, $B = \{3,4,5\}$:
A) $A \cap B = \{3\}$
B) $A \cup B = \{1,2,3,4,5\}$
C) $A - B = \{1,2\}$
Correctly, $B - A = \{4,5\}$
Correct Answers: A, B, C

14. Given $n(A) = 20$, $n(B) = 15$, $n(A \cap B) = 5$:
Using
$n(A \cup B) = n(A) + n(B) - n(A \cap B)$
$n(A \cup B) = 20 + 15 - 5 = 30$
A) $n(A \cup B) = 30$
B) $n(A - B) = 20 - 5 = 15$
C) $n(B - A) = 15 - 5 = 10$
(since intersection $= 5 \neq 0$)
Correct Answers: A, B, C

COMPREHENSION TYPE

PASSAGE

15. If $A = \{2,4,6,8,10\}$and $B = \{1,2,3,4,5\}$, then $A \cap B$is
Common elements in both sets are 2and 4.
$A \cap B = \{2,4\}$
Correct Answer: A) $\{2, 4\}$

16. Let $U = \{1,2,3,4,5,6,7,8,9\}$and $A = \{2,4,6,8\}$. Then $A'$is
Complement means elements in Uthat are not in A.
$A' = \{1,3,5,7,9\}$
Correct Answer: A) $\{1, 3, 5, 7, 9\}$

17. If $n(A) = 20$, $n(B) = 18$, and $n(A \cap B) = 7$, then $n(A \cup B)$equals
Using the formula:
$n(A \cup B) = n(A) + n(B) - n(A \cap B)$
$n(A \cup B) = 20 + 18 - 7 = 31$
Correct Answer: A) 31

MATRIX MATCHING TYPE

18. Given:
$A = \{1,2,3,4,5,6\}$
$B = \{2,4,6,8\}$
$C = \{1,3,5,7\}$
Evaluate Column A

1. $A \cap B = \{2,4,6\} \rightarrow$ C
2. $A \cup B = \{1,2,3,4,5,6,8\} \rightarrow$ A
3. $A - B = \{1,3,5\} \rightarrow$ B
4. $B \cap C = \emptyset \rightarrow$ D

Correct Matching

$1 \to C, 2 \to A, 3 \to B, 4 \to D$

INTEGER TYPE

19. $A = \{2,4,6,8,10\}$(multiples of 2 less than 12)
$B = \{2,3,5,7,11\}$(prime numbers less than 12)
$A \cap B = \{2\}$
$n(A \cap B) = 1$
Answer: $\boxed{1}$

20. Using the formula:
$n(A \cup B) = n(A) + n(B) - n(A \cap B)$
Substitute the values:
$28 = 15 + 20 - n(A \cap B)$
$28 = 35 - n(A \cap B)$
$n(A \cap B) = 35 - 28 = 7$
Answer: $\boxed{7}$

STUDENT PRACTISE CORNER

SINGLE ANSWER CORRECT TYPE

1. $A = \{2,3,5,7,11,13\}$(prime numbers less than 15)
$B = \{1,3,5,7,9,11,13\}$(odd numbers less than 15)
$A \cap B = \{3,5,7,11,13\}$
Answer: B) $\{\mathbf{3}, \mathbf{5}, \mathbf{7}, \mathbf{11}, \mathbf{13}\}$

2. $A \cap B$= numbers that are multiples of both 2 and 3
$\Rightarrow$ multiples of 6 in $U = \{1,2, \ldots, 20\}$
$\{6, 12, 18\}$
$n(A \cap B) = 3$
Answer: B) 3

3. $A - B = \{1,2,3,4,5\} - \{2,4,6,8\}$
Remove common elements 2,4:
$A - B = \{1,3,5\}$
Answer: B) $\{\mathbf{1}, \mathbf{3}, \mathbf{5}\}$

4. Using formula:
$n(M \cup S) = 35 + 25 - 10 = 50$
Students who like neither:
$60 - 50 = 10$
Answer: B) 10

5. Using the formula:
$n(A \cup B) = n(A) + n(B) - n(A \cap B)$
Substitute the values:
$35 = 25 + 20 - n(A \cap B)$
$35 = 45 - n(A \cap B)$
$n(A \cap B) = 45 - 35 = 10$
Answer: B) 10

6. $A = \{1,2,3\}, B = \{3,4,5\}, C = \{5,6,7\}$
First,
$A \cap B = \{3\}$
Now,
$\{3\} \cap C = \emptyset$
$A \cap B \cap C = \emptyset$

Answer: C) $\emptyset$

ONE OR MORE THAN ONE CORRECT ANSWER TYPE

7. Equal sets have the same elements.

A) $\{1,2,3\} = \{3,2,1\}$

C) $\{2,4,6\} = \{6,4,2\}$

D) Even prime number = 2, so sets are equal

Correct Answers: A, C, D

8. Given:

$U = \{1,2,3,4,5,6,7,8\}$

$A = \{2,4,6,8\}, B = \{1,2,3,4\}$

A) $A' = \{1,3,5,7\}$

B) $B' = \{5,6,7,8\}$

C) $A \cap B = \{2,4\}$

Correct Answers: A, B, C

9. $A = \{2,4,6,8\}, B = \{2,4,6,8\}$

A) $A = B$

B) $n(A) = 4$

D) Number of elements in power set:

$| P(A) | = 2^4 = 16$

Correct Answers: A, B, D

COMPREHENSION TYPE

PASSAGE

10. $R = \{1,2,3,4,5\}, C = \{4,5,6,7\}$

Common elements are 4,5.

$R \cap C = \{4,5\}$

Answer: A) $\{4, 5\}$

11.

$R = \{a, b, c, d\}, C = \{c, d, e, f\}$

$R \cup C = \{a, b, c, d, e, f\}$

Number of elements = 6.

Answer: A) 6 elements

12. $R = \{2,4,6,8\}, C = \{1,2,3,4\}$

Remove common elements 2,4from R:

$R - C = \{6,8\}$

Answer: A) $\{6, 8\}$

MATRIX MATCHING TYPE

13. Given:

P = {prime numbers less than 10} = {2,3,5,7}

Q = {odd numbers less than 10} = {1,3,5,7,9}

Match the Following

1. $P = \{2,3,5,7\} \rightarrow$ B
2. $Q = \{1,3,5,7,9\} \rightarrow$ A
3. $P \cap Q = \{3,5,7\} \rightarrow$ C
4. $Q - P = \{1,9\} \rightarrow$ D

Correct Matching $\mathbf{1 \rightarrow B, 2 \rightarrow A, 3 \rightarrow C, 4 \rightarrow D}$

INTEGER TYPE

14. Using the formula:
$n(A \cup B) = n(A) + n(B) - n(A \cap B)$
Substitute the values:
$n(A \cup B) = 12 + 8 - 3 = 17$
Answer: $\boxed{17}$

15. $A = \{1,2,3,4\}, B = \{3,4,5,6\}$
$A \cap B = \{3,4\}$
$n(A \cap B) = 2$
Answer: $\boxed{2}$

RELATIONS

LEVEL-1

1. $A = \{1,2,4\}, B = \{2,4,5\}, C = \{2,5\}$
$A - B = \{1\}, B - C = \{4\}$
$(A - B) \times (B - C) = \{(1,4)\}$
Answer: B) $\{(\mathbf{1},\mathbf{4})\}$

2. Relation defined by:
$xRy \iff y = 3x$
Possible ordered pairs:
$(1,3),\ (2,6),\ (3,9)$
Hence,
$R = \{(1,3),(2,6),(3,9)\}$
Answer: D) $\{(\mathbf{1},\mathbf{3}),(\mathbf{2},\mathbf{6}),(\mathbf{3},\mathbf{9})\}$

3. Given:
$R = \{(1,3),(2,5),(3,3)\}$
Inverse relation is obtained by interchanging coordinates:
$R^{-1} = \{(3,1),(5,2),(3,3)\}$
Answer: A) $\{(3,3),(3,1),(5,2)\}$

4. $A = \{1,2,3\}, B = \{1,4,6,9\}$
Relation defined by $x > y$.
Possible related pairs:
$(2,1),\ (3,1)$
Range of R is:
$\{1\}$
Answer: C) $\{\mathbf{1}\}$

5. Given:
$R = \{(x,y): x,y \in \mathbb{Z},\ x^2 + y^2 \leq 4\}$
For domain, find all possible $x$values satisfying:
$x^2 + y^2 \leq 4$
Possible integer values of x:
$-2,-1,0,1,2$
Answer: C) $\{-\mathbf{2},-\mathbf{1},\mathbf{0},\mathbf{1},\mathbf{2}\}$

6. Relation on Ndefined by:

$x + 2y = 8$

Check natural number solutions:

$y = 1 \Rightarrow x = 6$

$y = 2 \Rightarrow x = 4$

$y = 3 \Rightarrow x = 2$

Thus domain = {2,4,6}.

Answer: C) $\{\mathbf{2}, \mathbf{4}, \mathbf{6}\}$

7. Given:

$y = x - 3$

From $A = \{11,12,13\}$to $B = \{8,10,12\}$:

$11 \to 8$

$13 \to 10$

So,

$R = \{(11,8), (13,10)\}$

Inverse relation:

$R^{-1} = \{(8,11), (10,13)\}$

Answer: A) $\{(\mathbf{8}, \mathbf{11}), (\mathbf{10}, \mathbf{13})\}$

8. Number of elements in Cartesian product:

$| A \times B | = pq$

Answer: C) $\boldsymbol{pq}$

9. A relation from set $A$to set $B$is a subset of $A \times B$.

$R \subseteq A \times B$

Answer: C) $\boldsymbol{R \subseteq A \times B}$

10. If set A has m elements and set B has n elements, then

$| A \times B | = mn$

A relation from A to B is any subset of $A \times B$.

Number of subsets of a set having mn elements is:

2^{mn}

Answer: A) $\boldsymbol{2^{mn}}$

11. For a set A with $n$elements:

$| A \times A | = n^2$

A relation on A is any subset of $A \times A$.

Hence number of relations on $A$is:

2^{n^2}

Answer: B) $\boldsymbol{2^{n^2}}$

12. Given:

$A = \{1,2,3\}, R = \{(2,3)\}$

Not symmetric since $(3,2) \notin R$

Transitive is true vacuously (no pairs to violate transitivity)

Hence, relation is transitive only.

Answer: C) transitive only

13.

$R = \{(1,2), (2,2), (1,1), (4,4), (1,3), (3,3), (3,2)\}$
Reflexive: $(1,1), (2,2), (3,3), (4,4) \in R$

Transitive: conditions satisfy transitivity

Hence, $R$is reflexive and transitive but not symmetric.

Answer: B) Reflexive and transitive but not symmetric

14.

Since $(1, 2)$belongs to an equivalence relation, symmetry and transitivity imply:
$(2,1), (1,1), (2,2) \in R$
Possible equivalence relations on $\{1, 2, 3\}$:

1. $\{\{1,2\}, \{3\}\}$
2. $\{\{1,2,3\}\}$

Hence, there are 2such equivalence relations.

Answer: B) 2

15.

Relation on integers defined by:
$(x, y) \in R \iff |\, x - y \,| \leq 1$
Reflexive: $|\, x - x \,| = 0 \leq 1$✓

Symmetric: $|\, x - y \,| = |\, y - x \,|$✓

Not transitive:

$(1,2) \in R,\ (2,3) \in R$
but

$(1,3) \notin R$
Hence, relation is reflexive and symmetric only.

Answer: B) reflexive and symmetric

16. Given

$(a, b)R(c, d) \iff a + d = b + c$
Check properties:

Reflexive:

$a + b = b + a$
True.

Symmetric:
If $a + d = b + c$, then $c + b = d + a$, true.

Transitive:
If

$a + d = b + c$ and$c + f = d + e$
then adding,

$a + f = b + e$
So transitive.

Hence $R$is an equivalence relation.

Correct Answer: C) an equivalence relation

17. The largest equivalence relation on a set $A$is

$A \times A$

Every relation Son $A$is contained in $A \times A$.

So,

$S \subset R$

Correct Answer: B) $S \subset R$

18. Check which is not an equivalence relation on $\mathbb{Z}$.

$a - b$even ⇒ equivalence relation.

$a = b$⇒ equivalence relation.

$a < b$is not reflexive and not symmetric.

Correct Answer: C) $aRb \iff a < b$

19. "Is congruent to" between triangles is:

Reflexive

Symmetric

Transitive

Therefore it is an equivalence relation.

Correct Answer: C) equivalence

20. Reflexive relation from $A = \{1,2\}$to $B = \{1,2,3\}$:

Total possible ordered pairs:

$| A \times B | = 2 \times 3 = 6$

For reflexive relation, $(1,1)$and $(2,2)$must be included.

Remaining pairs:

$6 - 2 = 4$

Each can be chosen or not:

$2^4 = 16$

Correct Answer: A) 16

LEVEL – II

1. Given

$A = \{a, b, c\}$

$R_1 = \{(c,a), (b,b), (a,c), (b,c), (a,a)\}$

$R_2 = \{(a,b), (b,a), (c,c), (c,a), (a,a), (b,b), (a,c)\}$

Check symmetry and transitivity.

For R_1

$(b,c) \in R_1$, but $(c,b) \notin R_1$

Hence, $R_1$is not symmetric.

Now check transitivity:

$(a,c)$and $(c,a) \Rightarrow (a,a) \in R_1$

$(c,a)$and $(a,c) \Rightarrow (c,c)$should be in R_1, but it is not.

Hence, $R_1$is not transitive.

For R_2

$(a,b) \in R_2$and $(b,a) \in R_2$

$(a,c) \in R_2$and $(c,a) \in R_2$

Hence, $R_2$is symmetric.

Now check transitivity:

$(b, a)$and $(a, c) \in R_2$
then (b, c)should belong to R_2, but it does not.
Hence, $R_2$is not transitive.
Correct Answer: A

2. Given
$R_1 = \{(x, y) \in \mathbb{N} \times \mathbb{N}: 2x + y = 0\}$
$R_2 = \{(x, y) \in \mathbb{N} \times \mathbb{N}: x + 2y = 10\}$
For R_1
Since $x, y \in \mathbb{N}$,
$2x + y = 0$
has no natural number solution.
So,
$R_1 = \emptyset$
Range of $R_1$is empty, so option C is false.
An empty relation is vacuously symmetric and transitive.
For R_2
$x + 2y = 10$
Possible natural number solutions:
$(8,1), (6,2), (4,3), (2,4)$
Hence,
Range$(R_2) = \{1,2,3,4\}$
So option B is true.
Now check symmetry:
$(8,1) \in R_2$, but $(1,8) \notin R_2$
Hence, $R_2$is not symmetric.
Also not transitive.
Correct Answer: B

3. Given
$R = \{(x, y): x, y \in \mathbb{Z},\ x^2 + 3y^2 \leq 8\}$
We need the domain of R^{-1}.
Domain of R^{-1}= Range of R.
So find all possible $y$such that
$x^2 + 3y^2 \leq 8$
for some integer x.
$3y^2 \leq 8 \Rightarrow y^2 \leq \frac{8}{3}$
Hence,
$y = -1,0,1$
Check each:
$y = 0 \Rightarrow x^2 \leq 8$possible
$y = \pm 1 \Rightarrow x^2 + 3 \leq 8 \Rightarrow x^2 \leq 5$possible
Thus all are valid.
Therefore,
Domain$(R^{-1}) = \{-1,0,1\}$
Correct Answer: A

4. Problem: Let $R_1$and R_2be two relations defined as follows:
$R_1 = \{(a, b) \in \mathbb{R}^2: a^2 + b^2 \in \mathbb{Q}\}$
and $R_2 = \{(a, b) \in \mathbb{R}^2: a^2 + b^2 \notin \mathbb{Q}\}$
where $\mathbb{Q}$is the set of all rational numbers. Determine their transitivity.

A relation $R$is transitive if
$(a, b) \in R$ and $(b, c) \in R \Rightarrow (a, c) \in R$
We test both relations using counterexamples.
Check Transitivity of R_1
Condition: $(x, y) \in R_1 \iff x^2 + y^2 \in \mathbb{Q}$
Let $a^2 = 1 + \sqrt{2}, b^2 = 1 - \sqrt{2}$
Then,
$a^2 + b^2 = (1 + \sqrt{2}) + (1 - \sqrt{2}) = 2 \in \mathbb{Q}$
So,$(a, b) \in R_1$
Now let $c^2 = 1 + \sqrt{2}$
Then,$b^2 + c^2 = (1 - \sqrt{2}) + (1 + \sqrt{2}) = 2 \in \mathbb{Q}$
So,$(b, c) \in R_1$
Now check (a, c):
$a^2 + c^2 = (1 + \sqrt{2}) + (1 + \sqrt{2}) = 2 + 2\sqrt{2}$
Since $2 + 2\sqrt{2} \notin \mathbb{Q}$
we get $(a, c) \notin R_1$
Result: R_1 is not transitive
Check Transitivity of R_2
Condition: $(x, y) \in R_2 \iff x^2 + y^2 \notin \mathbb{Q}$
Let $a^2 = \sqrt{2}, b^2 = 1, c^2 = 2 - \sqrt{2}$
Check (a, b):
$a^2 + b^2 = \sqrt{2} + 1 \notin \mathbb{Q}$
So, $(a, b) \in R_2$
Check (b, c):
$b^2 + c^2 = 1 + (2 - \sqrt{2}) = 3 - \sqrt{2} \notin \mathbb{Q}$
So, $(b, c) \in R_2$
Now check (a, c):
$a^2 + c^2 = \sqrt{2} + (2 - \sqrt{2}) = 2$
Since $2 \in \mathbb{Q}$
we get $(a, c) \notin R_2$
Result: R_2 is not transitive
Correct Option: C) Neither R_1 nor R_2 is transitive

5. Given $A = \{1,2,3\}$
and $R = \{(1,1), (2,2), (3,3)\}$
Check the properties of the relation.
Reflexive
Since $(1,1), (2,2), (3,3) \in R$
R is reflexive.
Symmetric
Every ordered pair is of the form (a, a), so symmetry holds automatically.
Hence, $R$is symmetric.
Transitive
If $(a, a) \in R$ and $(a, a) \in R$
then $(a, a) \in R$
So, $R$is transitive.
Antisymmetric
If $(a, b) \in R$ and $(b, a) \in R$

then necessarily $a = b$.
Hence, $R$is antisymmetric.
Therefore, $R$is both:
an equivalence relation
a partial order relation

Correct Option: C) both equivalence and partial ordered

6. Given $A = \{1,2,3,4\}, B = \{1,3,5\}$
Relation R from A to B is defined by
$(a, b) \in R \iff a < b$
First find R.
Possible pairs:
$R = \{(1,3), (1,5), (2,3), (2,5), (3,5), (4,5)\}$
Now find inverse relation R^{-1}:
Reverse each ordered pair.
$R^{-1} = \{(3,1), (5,1), (3,2), (5,2), (5,3), (5,4)\}$
Now compute $R \circ R^{-1}$
For composition: $(a, c) \in R \circ R^{-1}$
if there exists b such that
$(a, b) \in R^{-1}$ and $(b, c) \in R$
Checking all possibilities gives:
$R \circ R^{-1} = \{(3,3), (3,5), (5,3), (5,5)\}$

Correct Option: C

7. Given $(a, b) \in S \iff ab \geq 0$
where S is a relation on $\mathbb{R}$.
Reflexive
For every real number a, $a \cdot a = a^2 \geq 0$
Hence, S is reflexive.
Symmetric
If $ab \geq 0$
then $ba \geq 0$
Hence, S is symmetric.
Transitive
Take $a = 1, b = 0, c = -1$
Then $ab = 0 \geq 0, bc = 0 \geq 0$
but $ac = -1 < 0$
So, S is not transitive.
Thus, S is only reflexive and symmetric.

Correct Option: B) reflexive and symmetric only

8. Given
Find which relation on $\mathbb{Z}$is not an equivalence relation.
A) $xRy \iff x \leq y$
Reflexive: Yes, since $x \leq x$
Symmetric: No, because $x \leq y$does not imply $y \leq x$
Hence, this is not an equivalence relation.
B) $xRy \iff x = y$
Equality relation is reflexive, symmetric, and transitive.
Hence, it is an equivalence relation.
C) $xRy \iff x - y$is even

Reflexive: $x - x = 0$is even
Symmetric: if $x - y$is even, then $y - x$is even
Transitive: sum of even integers is even
Hence, it is an equivalence relation.
D) $xRy \Leftrightarrow x \equiv y(\text{mod}3)$
Congruence modulo 3is reflexive, symmetric, and transitive.
Hence, it is an equivalence relation.
Correct Option: A

9. Given
Relation R from $\{2,3,4,5\} \rightarrow \{3,6,7,10\}$
defined by $xRy \Leftrightarrow x$ is relatively prime to y
We need the domain of R.
Domain = set of all first elements having at least one related element.
`Check each element:
For $x = 2$
$\gcd(2,3) = 1$
So, 2 belongs to domain.
For $x = 3$
$\gcd(3,7) = 1$
So, 3belongs to domain.
For $x = 4$
$\gcd(4,3) = 1$
So, 4 belongs to domain.
For $x = 5$
$\gcd(5,3) = 1$
So, 5belongs to domain.
Therefore, Domain$(R) = \{2,3,4,5\}$
Correct Option: D

10. Given
Relation $\phi$from $\mathbb{C}$to $\mathbb{R}$is defined by
$x\phi y \Leftrightarrow |x| = y$
Check the options.
A) $(2 + 3i)\phi 3$
$|2 + 3i| = \sqrt{2^2 + 3^2} = \sqrt{13} \neq 3$
False.
B) $\{2, 4, 6, 8\}$
Not a valid ordered relation statement.
False.
C) $(1 + i)\phi 2$
$|1 + i| = \sqrt{1^2 + 1^2} = \sqrt{2} \neq 2$
False.
D) $i\phi 1$
$|i| = 1$
True.
Correct Option: D

ONE OR MORE THAN ONE CORRECT ANSWER TYPE

11. Given Relation

$R = \{(1,2), (2,3), (3,4), (4,5)\}$

Domain = set of first elements

$\{1, 2, 3, 4\}$

Range = set of second elements

$\{2, 3, 4, 5\}$

$(5,4) \notin R$

Since no universal set is given, codomain cannot be concluded as $\{1, 2, 3, 4, 5\}$.

Correct Answers: $\boxed{A, B}$

12. Given Relation

$R = \{(1,2), (2,1), (2,3), (3,2)\}$

For every $(a, b) \in R$, $(b, a) \in R$also exists.

Hence, $R$is symmetric.

Reflexive relation requires

$(1,1), (2,2), (3,3) \in R$

which are not present. So $R$is not reflexive.

$(3,2) \in R$is true.

$(1,3) \notin R$

Correct Answers: $\boxed{A, C}$

13. Given Relation

$R = \{(1,2), (2,3), (1,3), (3,4), (1,4), (2,4)\}$

$(2,4) \in R$, so statement B is true.

$(3,1) \notin R$, so statement C is false.

Since $(1,2) \in R$and $(2,3) \in R$, and $(1,3) \in R$, statement D is true.

Also,

$(2,3), (3,4) \Rightarrow (2,4) \in R$

and

$(1,3), (3,4) \Rightarrow (1,4) \in R$

Hence R is transitive.

Correct Answers: $\boxed{A, B, D}$

14. Given Relation

$R = \{(1,1), (2,2), (3,3), (1,2), (2,1)\}$

All $(a, a)$are present for $a \in \{1,2,3\}$.

Hence, $R$is reflexive.

Since $(1,2) \in R$and $(2,1) \in R$, $R$is symmetric.

Checking transitive:

$(1,2), (2,1) \Rightarrow (1,1) \in R$

$(2,1), (1,2) \Rightarrow (2,2) \in R$

All required pairs are present, so $R$is transitive.

$(3,1) \notin R$

Correct Answers: $\boxed{A, B, C}$

COMPREHENSION TYPE

PASSAGE

15. Given Relation

$A = \{1,2,3\}, R = \{(1,1), (2,2), (3,3), (1,2), (2,1)\}$

All $(a, a)$are present, so $R$is reflexive.

Since $(1,2) \in R$and $(2,1) \in R$, $R$is symmetric.

Hence $R$is reflexive and symmetric.

Correct Answer: $\boxed{C}$

16. Reflexive Relation

A relation on set $A$is reflexive if every element is related to itself.

$(a, a) \in R$ for every $a \in A$

Correct Answer: $\boxed{B}$

17. Symmetric Relation on Integers

A relation is symmetric if:

$aRb \Rightarrow bRa$

$a > b$is not symmetric.

$a = b$is symmetric because if $a = b$, then $b = a$.

$a < b$is not symmetric.

$a = b + 1$is not symmetric.

Correct Answer: $\boxed{B}$

MATRIX MATCHING TYPE

18. Matrix Matching

(a) $R = \{(1,1), (2,2), (3,3)\}$

All elements are related to themselves.

So, $R$is reflexive.

$(a) \rightarrow (ii)$

(b) $R = \{(1,2), (2,1), (2,3), (3,2)\}$

Each pair has its reverse pair.

So, $R$is symmetric but not reflexive.

$(b) \rightarrow (i)$

(c) $R = \{(1,2), (2,3), (1,3)\}$

Since

$(1,2), (2,3) \Rightarrow (1,3) \in R$

$R$is transitive.

$(c) \rightarrow (iii)$

(d) $R = \{(1,2), (2,3)\}$on $A = \{1,2,3\}$

Not reflexive because $(1,1), (2,2), (3,3) \notin R$

Not symmetric because $(1,2) \in R$but $(2,1) \notin R$

$(d) \rightarrow (iv)$

Correct Matching

$\boxed{(a) \rightarrow (ii),\ (b) \rightarrow (i),\ (c) \rightarrow (iii),\ (d) \rightarrow (iv)}$

Integer Type

19. Given:

$A = \{1,2,3,4,5\}$

Relation: $R = \{(a, b): a + b = 6\}$

Possible ordered pairs: $(1,5), (2,4), (3,3), (4,2), (5,1)$

Total ordered pairs: $\boxed{5}$

20. **Integer Type**

$A = \{1,2,3\}, B = \{a, b, c, d\}$

Number of elements in $A \times B$:

$3 \times 4 = 12$

Total number of relations from $A$to B:

$2^{12} = 4096$

Answer: $\boxed{4096}$

STUDENT PRACTISE CORNER

1. Given $A = \{1,2\}$
A symmetric relation on Adepends on pairs in $A \times A$:
$(1,1), (2,2), (1,2), (2,1)$
For symmetry, $(1,2)$and $(2,1)$must occur together.
$(1,1)$: choose or not choose ⇒ 2ways
$(2,2)$: 2ways
Pair $(1,2), (2,1)$: together or absent ⇒ 2ways
Total: $2 \times 2 \times 2 = 8$
Correct Answer: $\boxed{A}$

2. Relation on N:
$2x + y = 9$
For natural numbers:
$x = 1 \Rightarrow y = 7$
$x = 2 \Rightarrow y = 5$
$x = 3 \Rightarrow y = 3$
$x = 4 \Rightarrow y = 1$
So domain: $\{1, 2, 3, 4\}$
Correct Answer:$\boxed{A}$

3. Given: Set $A = \{a, b, c, d\}$
Relation $R = \{(a, b), (b, a), (a, a)\}$
Let's check the properties of the relation:
Reflexive
A relation is reflexive if every element in the set relates to itself.
Meaning, for every $x \in A$, the pair (x, x)must be in R.
For set A, we would need $(a, a), (b, b), (c, c), (d, d)$ to all be in R.
Since (b, b), (c, c), and $(d, d)$are missing, the relation is not reflexive.
Symmetric
A relation is symmetric if whenever $(x, y) \in R$, then (y, x)must also be in R.
$(a, b) \in R$, and its reverse $(b, a) \in R$
$(b, a) \in R$, and its reverse $(a, b) \in R$
$(a, a) \in R$, and its reverse is also $(a, a) \in R$
Every pair has its symmetric counterpart in the relation.
Therefore, the relation is symmetric.
Transitive
A relation is transitive if whenever
$(x, y) \in R$ and $(y, z) \in R$,
then $(x, z) \in R$
We have $(b, a) \in R$ and $(a, b) \in R$
For transitivity, the pair (b, b)
must also belong to R.
Since $(b, b) \notin R$,
the relation is not transitive.
correct option is $\boxed{C)\ \textbf{symmetric only}}$

4. Given $R = \{(1,1), (2,2), (3,3)\}$
Reflexive: yes

Symmetric: yes
Transitive: yes
Correct Answer: $\boxed{D}$

5. Given$R = \{(1,1),(2,2),(3,3),(1,2),(2,1)\}$
All (a,a)present ⇒reflexive.
$(1,2)$and $(2,1)$both present ⇒symmetric.
Correct Answer: $\boxed{A}$

6. Given: $A = \{1,2,3\}$
A relation on Amust contain only ordered pairs whose elements belong to A.
Let's check each option:
Option A $\{(1,2),(2,3)\}$
Both elements in each ordered pair belong to A.
So, this is a relation on A.
Option B $\{(1,1),(2,2)\}$
All elements belong to A.
So, this is a relation on A.
Option C $\{(1,4)\}$
Here, $4 \notin A$.
Therefore, this is not a relation on A.
Option D ϕ
The empty set is always a relation on any set.
Conclusion The set which is NOT a relation on A is $\boxed{C)\ \{(1,4)\}}$

ONE OR MORE THAN ONE CORRECT ANSWER TYPE

7. Given $A = \{1,2,3\}, R = \{(1,1),(2,2),(3,3)\}$
Reflexive
All pairs $(1,1),(2,2),(3,3)$
are present in R.
So, R is reflexive.
Symmetric
Every pair is of the form (a,a), whose reverse is itself.
So, R is symmetric.
Transitive
For any $(a,a),(a,a) \in R$,
we also have $(a,a) \in R$.
So, R is transitive.
Correct options: $\boxed{A, B, C}$

8. Given $A = \{1,2,3\}, R = \{(1,1),(1,2),(2,2),(3,3)\}$
Reflexive
All required pairs $(1,1),(2,2),(3,3)$
are present.
So, R is reflexive.
Symmetric
Since $(1,2) \in R$
but $(2,1) \notin R$,
R is not symmetric.
Checking $(2,1) \in R$
$(2,1) \notin R$

So statement C is false.

Correct options: $\boxed{A,\ D}$

9. Given $A = \{1,2,3\}, R = \{(1,2),(2,3)\}$

Domain

The set of first elements is $\{1,2\}$

So statement A is true.

Range

The set of second elements is $\{2,3\}$

So statement B is true.

Reflexive

For reflexive relation, we need $(1,1),(2,2),(3,3) \in R$

which are missing.

So, R is not reflexive.

Checking $(1,3) \in R$

$(1,3) \notin R$

So statement D is false.

Correct options: $\boxed{A,\ B}$

COMPREHENSION TYPE

PASSAGE

10. Given

$A = \{a,b,c\}, B = \{x,y\}$

$R = \{(a,x),(b,x),(b,y)\}$

Range of R

The range is the set of second elements of the ordered pairs.

From R, the second elements are: $x,\ y$

So, the range is $\{x,y\}$

The correct option is $\boxed{B)\ \{x,y\}}$

11. Given $R = \{(a,x),(b,x),(b,y)\}$

Domain of R

The domain is the set of first elements of the ordered pairs.

From R, the first elements are: $a,\ b$

So, the domain is $\{a,b\}$

The correct option is $\boxed{A)\ \{a,b\}}$

12. Total Number of Relations

If one set has 2elements and the other set has 3elements, then

$|\ A \times B\ | = 2 \times 3 = 6$

A relation is any subset of $A \times B$.

Therefore, total number of relations is 2^6

correct option is $\boxed{B)\ 2^6}$

MATRIX MATCHING TYPE

13. Match the following relations with the correct statements

(a) Reflexive relation

In a reflexive relation, every element is related to itself.

$(a) \rightarrow (iii)$

(b) Symmetric relation

A relation is symmetric if

$(a,b) \in R \Rightarrow (b,a) \in R$

$(b) \rightarrow (i)$

(c) Transitive relation

A relation is transitive if

$(a, b) \in R$ and $(b, c) \in R \Rightarrow (a, c) \in R$

$(c) \rightarrow (ii)$

(d) Antisymmetric relation

A relation is antisymmetric if

$(a, b) \in R$ and $(b, a) \in R \Rightarrow a = b$

$(d) \rightarrow (iv)$

Correct Matching $\boxed{(\boldsymbol{a}) \rightarrow (\boldsymbol{iii}), (\boldsymbol{b}) \rightarrow (\boldsymbol{i}), (\boldsymbol{c}) \rightarrow (\boldsymbol{ii}), (\boldsymbol{d}) \rightarrow (\boldsymbol{iv})}$

INTEGER TYPE

14. Given $A = \{1,2,3\}$

Relation $R = \{(a, b): a < b\}$

List all ordered pairs satisfying $a < b$:

$(1,2), (1,3), (2,3)$

Total ordered pairs: $\boxed{3}$

15. Number of Antisymmetric Relations on $A = \{1,2,3\}$

For antisymmetric relations:

Diagonal pairs:

$(1,1), (2,2), (3,3)$

may be included or not independently.

2^3 ways

For each distinct pair: $(1,2), (2,1)$

at most one can be chosen together.

Choices:

choose neither

choose first

choose second

3 ways

Distinct unordered pairs are:$\{1,2\}, \{1,3\}, \{2,3\}$

So total:$2^3 \times 3^3 = 8 \times 27 = 216$

Therefore, $\boxed{216}$

8.Limits

LEVEL – I

1. Given:

f(x) = { $1 + x^2$, if $0 \le x \le 1$

$2 - x$, if $x > 1$}

Find the left-hand limit of f(x) at x = 1.

To find the left-hand limit, use the expression valid for $x \le 1$.

$\lim_{x\to1^-} f(x) = \lim_{x\to1^-}(1 + x^2)$

$= 1 + (1)^2 = 2$

Answer: B) 2

2. Given:

f(x) = { $5x - 4$, if $0 < x \le 1$

$4x^3 - 3x$, if $1 < x < 2$}

Find $\lim_{x\to0} f(x)$.

For x near 0, use $f(x) = 5x - 4$.

$\lim_{x\to0}(5x - 4)$

$= 5(0) - 4 = -4$

Answer: D) -4

3. $\lim_{x\to1}\left(\frac{1}{1-x} - \frac{3}{1-x^3}\right)$

Factor: $1 - x^3 = (1 - x)(1 + x + x^2)$

Substitute:

$= \lim_{x\to1}\left(\frac{1 + x + x^2 - 3}{(1 - x)(1 + x + x^2)}\right)$

Simplify numerator:

$x^2 + x - 2 = (x - 1)(x + 2)$

$= \lim_{x\to1}\frac{(x - 1)(x + 2)}{-(x - 1)(1 + x + x^2)}$

Cancel (x − 1):

$= \lim_{x\to1}\frac{x + 2}{-(1 + x + x^2)}$

Put x = 1:

$= \frac{3}{-3} = -1$

Answer: B) -1

4. $\lim_{x\to0}\frac{(1+x)(1+2x)(1+3x)-1}{x}$

$(1 + x)(1 + 2x) = 1 + 3x + 2x^2$

$(1 + 3x + 2x^2)(1 + 3x) = 1 + 6x + 11x^2 + 6x^3$

Substitute:

$= \lim_{x\to0}\frac{6x + 11x^2 + 6x^3}{x}$

$= \lim_{x\to0}(6 + 11x + 6x^2)$

Put x = 0: = 6

Answer: C) 6

5. $\lim_{x\to1}\left(\frac{x-1}{x^2-5x+4} + \frac{x-1}{3(x^2-3x+2)}\right)$

Factor denominators:

$x^2 - 5x + 4 = (x-1)(x-4)$
$x^2 - 3x + 2 = (x-1)(x-2)$
Substitute:
$= \lim_{x\to 1}\left(\frac{1}{x-4} + \frac{1}{3(x-2)}\right)$
Put x = 1:
$= \frac{1}{-3} + \frac{1}{-3}$
$= -\frac{2}{3}$

Answer: A) $-\frac{2}{3}$

6. $\lim_{x\to\sqrt{2}} \frac{\sqrt{3+2x}-\sqrt{2}-1}{x^2-2}$
Substitute $x = \sqrt{2}$:
$\sqrt{3+2\sqrt{2}} = \sqrt{2}+1$
So numerator and denominator both become 0.
Rationalize the numerator:
$\frac{\sqrt{3+2x}-(\sqrt{2}+1)}{x^2-2} \cdot \frac{\sqrt{3+2x}+(\sqrt{2}+1)}{\sqrt{3+2x}+(\sqrt{2}+1)}$
$= \frac{2(x-\sqrt{2})}{(x^2-2)\left(\sqrt{3+2x}+\sqrt{2}+1\right)}$
Since $x^2 - 2 = (x-\sqrt{2})(x+\sqrt{2})$,
$= \frac{2}{(x+\sqrt{2})(\sqrt{3+2x}+\sqrt{2}+1)}$
Putting $x = \sqrt{2}$:
$= \frac{2}{2\sqrt{2}\cdot 2(\sqrt{2}+1)} = \frac{1}{2\sqrt{2}(\sqrt{2}+1)}$
$= \frac{\sqrt{2}-1}{2\sqrt{2}}$

Correct Answer: D) $\frac{\sqrt{2}-1}{2\sqrt{2}}$

7. $\lim_{x\to 1} \frac{x^2-\sqrt{x}}{\sqrt{x}-1}$
Let $t = \sqrt{x} \Rightarrow x = t^2$.
Then,
$\frac{t^4 - t}{t-1} = \frac{t(t^3-1)}{t-1}$
$= t(t-1)(t^2+t+1)/(t-1)$
$= t(t^2+t+1)$
Now put $t = 1$:
$1(1+1+1) = 3$

Correct Answer: C) 3

8. $\lim_{x\to -a} \frac{x^9+a^9}{x+a} = 9$
Use identity:
$x^9 + a^9 = (x+a)(x^8 - x^7 a + \cdots - a^8)$
So limit becomes:
$(-a)^8 = a^8$

Given:

$a^8 = 9$

$a = \pm\sqrt[8]{9} = \pm\sqrt{3}$

Among the options, this corresponds to:

$a = \pm 3^{1/2}$

But simplifying from standard factor result:

$9a^8 = 9 \Rightarrow a^8 = 1$

Actually directly using derivative form:

$\lim_{x\to -a} \frac{x^9 - (-a)^9}{x - (-a)} = 9(-a)^8$

$9a^8 = 9 \Rightarrow a^8 = 1 \Rightarrow a = \pm 1$

Correct Answer: A) ± 1

9. $\lim_{x\to 0} \frac{\sqrt{1+x+x^2}-1}{x}$

Rationalize:

$$\frac{\sqrt{1+x+x^2}-1}{x} \cdot \frac{\sqrt{1+x+x^2}+1}{\sqrt{1+x+x^2}+1}$$

$$= \frac{x + x^2}{x(\sqrt{1+x+x^2}+1)}$$

$$= \frac{1+x}{\sqrt{1+x+x^2}+1}$$

Now put $x = 0$:

$$= \frac{1}{1+1} = \frac{1}{2}$$

Correct Answer: D) $\frac{1}{2}$

10. Given: $\lim_{x\to a} \frac{\sqrt{a+3x}-\sqrt{4x}}{\sqrt{4a+x}-\sqrt{5x}}$

Substitute $x = a$:

Numerator: $\sqrt{a+3a} - \sqrt{4a} = \sqrt{4a} - \sqrt{4a} = 0$

Denominator: $\sqrt{4a+a} - \sqrt{5a} = \sqrt{5a} - \sqrt{5a} = 0$

So, it is of the form $\frac{0}{0}$.

Apply L'Hôpital's Rule.

Differentiate numerator and denominator:

Numerator derivative:

$$\frac{3}{2\sqrt{a+3x}} - \frac{2}{\sqrt{4x}}$$

Denominator derivative:

$$\frac{1}{2\sqrt{4a+x}} - \frac{5}{2\sqrt{5x}}$$

Now put $x = a$:

$$\frac{\frac{3}{2\sqrt{4a}} - \frac{2}{\sqrt{4a}}}{\frac{1}{2\sqrt{5a}} - \frac{5}{2\sqrt{5a}}}$$

$$= \frac{\frac{3}{4\sqrt{a}} - \frac{1}{\sqrt{a}}}{\frac{1-5}{2\sqrt{5a}}}$$

$$= \frac{-\frac{1}{4\sqrt{a}}}{-\frac{2}{\sqrt{5a}}}$$

$$= \frac{\sqrt{5}}{8}$$

Correct Answer B: $\boxed{\frac{\sqrt{5}}{8}}$

11. $\lim_{x\to 8} \frac{\sqrt{1+\sqrt{1+x}}-2}{x-8}$

Substitute $x = 8$:

$$\sqrt{1+\sqrt{9}} - 2 = \sqrt{4} - 2 = 0$$

So use derivative idea:

Let $f(x) = \sqrt{1+\sqrt{1+x}}$

Then required limit is $f'(8)$.

$$f'(x) = \frac{1}{2\sqrt{1+\sqrt{1+x}}} \cdot \frac{1}{2\sqrt{1+x}}$$

At $x = 8$:

$$f'(8) = \frac{1}{2(2)} \cdot \frac{1}{2(3)} = \frac{1}{24}$$

Correct Answer: C) $\frac{1}{24}$

12. $\lim_{x\to 0} \frac{\sqrt{1+x^3}-\sqrt{1-x^3}}{x^3}$

Rationalize:

$$\frac{(1+x^3)-(1-x^3)}{x^3\left(\sqrt{1+x^3}+\sqrt{1-x^3}\right)}$$

$$= \frac{2x^3}{x^3\left(\sqrt{1+x^3}+\sqrt{1-x^3}\right)}$$

$$= \frac{2}{\sqrt{1+x^3}+\sqrt{1-x^3}}$$

Putting $x = 0$:

$$= \frac{2}{1+1} = 1$$

Correct Answer: B) 1

13. $\lim_{x\to\infty} \left(\sqrt{x^2+x+1}-\sqrt{x^2+1}\right)$

Rationalize:

$$\frac{(x^2+x+1)-(x^2+1)}{\sqrt{x^2+x+1}+\sqrt{x^2+1}}$$

$$= \frac{x}{\sqrt{x^2+x+1}+\sqrt{x^2+1}}$$

Divide numerator and denominator by x:

$$= \frac{1}{\sqrt{1+\frac{1}{x}+\frac{1}{x^2}}+\sqrt{1+\frac{1}{x^2}}}$$

As $x \to \infty$,

$= \frac{1}{1+1} = \frac{1}{2}$

Correct Answer: D) $\frac{1}{2}$

14. Evaluate

$\lim_{x\to\infty} \frac{-x^3 - 8}{2x^2 + 5x + 7}$

Highest powers are $x^3$and x^2.

$\frac{-x^3}{2x^2} = -\frac{x}{2}$

As $x \to \infty$, $-\frac{x}{2} \to -\infty$.

Correct Answer: C) $-\infty$

15. Evaluate

$\lim_{x\to\infty} \left(\sqrt{x^2 + ax + b} - x\right)$

Rationalize:

$= \lim_{x\to\infty} \frac{x^2 + ax + b - x^2}{\sqrt{x^2 + ax + b} + x}$

$= \lim_{x\to\infty} \frac{ax + b}{\sqrt{x^2 + ax + b} + x}$

Divide by x:

$= \lim_{x\to\infty} \frac{a + \frac{b}{x}}{\sqrt{1 + \frac{a}{x} + \frac{b}{x^2}} + 1}$

$= \frac{a}{2}$

Correct Answer: C) $\frac{a}{2}$

16. Evaluate

$\lim_{x\to\infty} \frac{|\, 3x^2 + 1 \,|}{2x^2 + 1}$

For large x,

$|\, 3x^2 + 1 \,| = 3x^2 + 1$

So,

$\lim_{x\to\infty} \frac{3x^2 + 1}{2x^2 + 1} = \frac{3}{2}$

Correct Answer: A) $\frac{3}{2}$

17. Evaluate $\lim_{n\to\infty} \frac{1^2+2^2+3^2+\cdots+n^2}{n^3}$

Use formula:

$1^2 + 2^2 + \cdots + n^2 = \frac{n(n+1)(2n+1)}{6}$

$= \lim_{n\to\infty} \frac{n(n+1)(2n+1)}{6n^3}$

$= \lim_{n\to\infty} \frac{(n+1)(2n+1)}{6n^2}$

$= \frac{2}{6} = \frac{1}{3}$

Correct Answer: C) $\frac{1}{3}$

18. $\lim_{x\to\infty}\left(\sqrt{x+\sqrt{x+\sqrt{x}}}-\sqrt{x}\right)$

To evaluate this limit, we multiply and divide by the conjugate:

$$\lim_{x\to\infty}\frac{\left(\sqrt{x+\sqrt{x+\sqrt{x}}}-\sqrt{x}\right)\left(\sqrt{x+\sqrt{x+\sqrt{x}}}+\sqrt{x}\right)}{\sqrt{x+\sqrt{x+\sqrt{x}}}+\sqrt{x}}$$

Using $(a-b)(a+b)=a^2-b^2$,

$$=\lim_{x\to\infty}\frac{(x+\sqrt{x+\sqrt{x}})-x}{\sqrt{x+\sqrt{x+\sqrt{x}}}+\sqrt{x}}$$

$$=\lim_{x\to\infty}\frac{\sqrt{x+\sqrt{x}}}{\sqrt{x+\sqrt{x+\sqrt{x}}}+\sqrt{x}}$$

Divide numerator and denominator by $\sqrt{x}$:

$$=\lim_{x\to\infty}\frac{\sqrt{1+\frac{1}{\sqrt{x}}}}{\sqrt{1+\frac{\sqrt{x+\sqrt{x}}}{x}}+1}$$

As $x\to\infty$,

$\frac{1}{\sqrt{x}}\to 0$ and $\frac{\sqrt{x+\sqrt{x}}}{x}\to 0$

Hence,

$$=\frac{\sqrt{1+0}}{\sqrt{1+0}+1}=\frac{1}{2}$$

Correct Option: B) $\frac{1}{2}$

19. Evaluate

$\lim_{x\to 0}\frac{3\,|\,x\,|+x}{7\,|\,x\,|-5x}$

LHL ($x<0$):

$|\,x\,|=-x$

$$\frac{3(-x)+x}{7(-x)-5x}=\frac{-2x}{-12x}=\frac{1}{6}$$

RHL ($x>0$):

$|\,x\,|=x$

$$\frac{3x+x}{7x-5x}=\frac{4x}{2x}=2$$

Since LHL ≠RHL, limit does not exist.

Correct Answer: D) Does not exist

20. Evaluate $\lim_{x\to 1}\frac{x^2-1}{|x-1|}$

Factorize:

$x^2-1=(x-1)(x+1)$

$$=\lim_{x\to 1}\frac{(x-1)(x+1)}{|\,x-1\,|}$$

For $x\to 1^+$:

$|x-1| = x-1 \Rightarrow x+1 \to 2$

For $x \to 1^-$:

$|x-1| = -(x-1) \Rightarrow -(x+1) \to -2$

LHL ≠RHL, so limit does not exist.

Correct Answer: D) Does not exist

LEVEL – II

SINGLE OPTION CORRECT TYPE

1. Given:

$$\lim_{x\to\infty} \frac{[ax+b]}{x}$$

Since $a > 0$, for very large x,

$[ax+b] \approx ax+b$

So,

$$\frac{[ax+b]}{x} \to \frac{ax+b}{x} = a + \frac{b}{x}$$

As $x \to \infty$, $\frac{b}{x} \to 0$.

Therefore,

$$\lim_{x\to\infty} \frac{[ax+b]}{x} = a$$

Correct Answer: C) a

2. Evaluate:

$$\lim_{x\to 1} \frac{(x+x^2+x^3+\cdots+x^n)-n}{x-1}$$

Let $f(x) = x+x^2+x^3+\cdots+x^n$

Then $f(1) = n$.

So the limit becomes

$$\lim_{x\to 1} \frac{f(x)-f(1)}{x-1} = f'(1)$$

Now,

$f'(x) = 1+2x+3x^2+\cdots+nx^{n-1}$

Putting $x = 1$,

$$f'(1) = 1+2+3+\cdots+n = \frac{n(n+1)}{2}$$

Correct Answer: D) $\frac{n(n+1)}{2}$

3. Given

$$f(x) = \begin{cases} bx+c, & x > 1 \\ 3cx-2b+1, & x < 1 \end{cases}$$

For limit to exist at $x = 1$,

LHL = RHL

Right-hand limit:

$$\lim_{x\to 1^+} f(x) = b+c$$

Left-hand limit:

$$\lim_{x\to 1^-} f(x) = 3c-2b+1$$

Equating,

$b+c = 3c-2b+1$

$3b-2c-1 = 0$

Correct Answer: C) $3b-2c-1=0$

4. Evaluate

$$\lim_{n\to\infty}\frac{1+\frac{1}{3}+\frac{1}{3^2}+\cdots+\frac{1}{3^{n-1}}}{1+\frac{1}{2}+\frac{1}{2^2}+\cdots+\frac{1}{2^{n-1}}}$$

Both are geometric series.

Numerator:

$$1+\frac{1}{3}+\frac{1}{3^2}+\cdots=\frac{1}{1-\frac{1}{3}}=\frac{3}{2}$$

Denominator:

$$1+\frac{1}{2}+\frac{1}{2^2}+\cdots=\frac{1}{1-\frac{1}{2}}=2$$

Therefore, $\lim=\frac{\frac{3}{2}}{2}=\frac{3}{4}$

Correct Answer: A) $\frac{3}{4}$

5. Evaluate $\lim_{x\to\infty}\frac{(x+1)^{10}+(x+2)^{10}+\cdots+(x+2018)^{10}}{x^{10}+2018^{10}}$

For large x,

$(x+k)^{10}\sim x^{10}$

There are 2018 terms in the numerator, so

Numerator $\sim 2018x^{10}$

Denominator:

$x^{10}+2018^{10}\sim x^{10}$

Hence,

$$\lim_{x\to\infty}\frac{2018x^{10}}{x^{10}}=2018$$

Correct Answer: B) 2018

6. Evaluate: $\lim_{x\to 2^-}(x-[x])^2$

As $x\to 2^-$, the greatest integer function gives

$[x]=1$

because x is just less than 2.

So,$x-[x]=x-1$

Therefore, $\lim_{x\to 2^-}(x-[x])^2=\lim_{x\to 2^-}(x-1)^2=(2-1)^2=1$

Correct Answer: B) 1

7. Evaluate

$$\lim_{n\to\infty}\frac{1}{n^4}[1^2+(1^2+2^2)+\cdots+(1^2+2^2+\cdots+n^2)]$$

The bracket equals

$$\sum_{k=1}^{n}\sum_{i=1}^{k}i^2$$

Using

$$1^2+2^2+\cdots+k^2=\frac{k(k+1)(2k+1)}{6}$$

Highest power term is

$$\frac{2k^3}{6}=\frac{k^3}{3}$$

Therefore,

$$\sum_{k=1}^{n} \frac{k^3}{3} = \frac{1}{3} \cdot \frac{n^2(n+1)^2}{4} \sim \frac{n^4}{12}$$

Thus,

$$\lim_{n\to\infty} \frac{n^4/12}{n^4} = \frac{1}{12}$$

Correct Answer: A) $\frac{1}{12}$

8. Evaluate $\lim_{n\to\infty} \frac{1\cdot 1!+2\cdot 2!+3\cdot 3!+\cdots+n\cdot n!}{(n+1)!}$

Use

$k \cdot k! = (k+1-1)k! = (k+1)! - k!$

So numerator becomes telescoping:

$(2! - 1!) + (3! - 2!) + \cdots + ((n+1)! - n!)$

All middle terms cancel:

$= (n+1)! - 1$

Hence,

$$\lim_{n\to\infty} \frac{(n+1)! - 1}{(n+1)!} = \lim_{n\to\infty} \left(1 - \frac{1}{(n+1)!}\right) = 1$$

Correct Answer: B) 1

ONE OR MORE THAN ONE CORRECT ANSWER TYPE

9. Evaluate:

$$\lim_{x\to\infty} \left(\frac{x^2+x+1}{x+1} - ax - b\right) = 4$$

First divide:

$$\frac{x^2+x+1}{x+1} = x + \frac{1}{x+1}$$

So, $x + \frac{1}{x+1} - ax - b = (1-a)x - b + \frac{1}{x+1}$

For the limit to be finite,

$1 - a = 0 \Rightarrow a = 1$

Then, $\lim_{x\to\infty} \left(-b + \frac{1}{x+1}\right) = -b$

Given limit = 4,

$-b = 4 \Rightarrow b = -4$

Correct Answers: A, D

10. Let $\alpha(a)$and $\beta(a)$be the roots of

$(\sqrt[3]{1+a} - 1)x^2 + (\sqrt{1+a} - 1)x + \sqrt[6]{1+a} - 1 = 0$

As $a \to 0^+$,

Using expansions:

$$\sqrt[3]{1+a} - 1 \sim \frac{a}{3}$$

$$\sqrt{1+a} - 1 \sim \frac{a}{2}$$

$$\sqrt[6]{1+a} - 1 \sim \frac{a}{6}$$

Equation becomes

$$\frac{a}{3}x^2 + \frac{a}{2}x + \frac{a}{6} = 0$$

Divide by a:

$\frac{1}{3}x^2+\frac{1}{2}x+\frac{1}{6}=0$

Multiply by 6:

$2x^2+3x+1=0$

$(2x+1)(x+1)=0$

Roots:$x=-\frac{1}{2},-1$

Hence,

$\alpha(a)\to-\frac{1}{2},\beta(a)\to-1$

Correct Answers: A, C

11. Evaluate: $\lim_{x\to0}\frac{a-\sqrt{a^2-x^2}-\frac{x^2}{4}}{x^4}=\ell$

Use expansion:

$\sqrt{a^2-x^2}=a\sqrt{1-\frac{x^2}{a^2}}$

$\sqrt{1-t}\approx1-\frac{t}{2}-\frac{t^2}{8}$

So,

$\sqrt{a^2-x^2}\approx a-\frac{x^2}{2a}-\frac{x^4}{8a^3}$

Then numerator:

$a-\left(a-\frac{x^2}{2a}-\frac{x^4}{8a^3}\right)-\frac{x^2}{4}$

$=\left(\frac{1}{2a}-\frac{1}{4}\right)x^2+\frac{x^4}{8a^3}$

For finite limit, coefficient of x^2must be zero:

$\frac{1}{2a}=\frac{1}{4}\Rightarrow a=2$

Then,$\ell=\frac{1}{8(2)^3}=\frac{1}{64}$

Correct Answers: A, C

12. Given:$\lim_{x\to2}\frac{x^2-4}{x-2}$

Factorize the numerator:

$x^2-4=(x-2)(x+2)$

So,$\frac{x^2-4}{x-2}=\frac{(x-2)(x+2)}{x-2}=x+2(x\neq2)$

Now taking limit as $x\to2$:

$\lim_{x\to2}(x+2)=4$

Also, at $x=2$, the original expression becomes $\frac{0}{0}$, which is an indeterminate form.

Therefore:

A) The limit exists → True

B) The limit is equal to 4 → True

C) The expression is indeterminate at $x=2$→ True

D) The limit is equal to 2 → False

Correct Answers: A, B and C

13. Given: $f(x) = \frac{x}{|x|+x^2}$

For $x > 0$,

$|\ x\ | = x$

$f(x) = \frac{x}{x+x^2} = \frac{1}{1+x}$

Thus, $\lim_{x\to 0^+} f(x) = 1$

For $x < 0$,

$|\ x\ | = -x$

$f(x) = \frac{x}{-x+x^2} = \frac{1}{x-1}$

Thus, $\lim_{x\to 0^-} f(x) = -1$

Since LHL ≠RHL, the limit at 0does not exist.

Correct Answers: A, B, D

COMPREHENSION TYPE

Passage – I:

14. Evaluate $\lim_{x\to 0} \frac{\sin^2 2x}{\sin^2 4x}$

Using $\lim_{x\to 0} \frac{\sin x}{x} = 1$,

$\frac{\sin^2 2x}{\sin^2 4x} = \left(\frac{\sin 2x}{\sin 4x}\right)^2$

As $x \to 0$,

$\sin 2x \sim 2x, \sin 4x \sim 4x$

So,

$\left(\frac{2x}{4x}\right)^2 = \left(\frac{1}{2}\right)^2 = \frac{1}{4}$

Correct Answer: C) $\frac{1}{4}$

15. Evaluate $\lim_{x\to a} \frac{\cos\sqrt{x} - \cos\sqrt{a}}{x-a}$

This is of the form

$\frac{f(x)-f(a)}{x-a} = f'(a)$

Let

$f(x) = \cos\sqrt{x}$

Differentiate:

$f'(x) = -\sin\sqrt{x} \cdot \frac{1}{2\sqrt{x}}$

Hence, $f'(a) = -\frac{\sin\sqrt{a}}{2\sqrt{a}}$

Correct Answer: C) $-\frac{\sin\sqrt{a}}{2\sqrt{a}}$

16. Evaluate $\lim_{x\to 0} \frac{x\tan 2x - 2x\tan x}{(1-\cos 2x)^2}$

Using standard limits:

$\tan 2x \sim 2x, \tan x \sim x$

Numerator:

$x(2x) - 2x(x) = 2x^2 - 2x^2 = 0$

Use next approximation:

$1 - \cos 2x \sim \frac{(2x)^2}{2} = 2x^2$

Then

$(1-\cos 2x)^2 \sim (2x^2)^2 = 4x^4$

Also,

$\tan 2x = 2x + \frac{(2x)^3}{3}, \tan x = x + \frac{x^3}{3}$

So numerator:

$x\left(2x+\frac{8x^3}{3}\right) - 2x\left(x+\frac{x^3}{3}\right) = 2x^2 + \frac{8x^4}{3} - 2x^2 - \frac{2x^4}{3} = 2x^4$

Therefore,

$\frac{2x^4}{4x^4} = \frac{1}{2}$

Correct Answer: C) $\frac{1}{2}$

MATRIX MATCHING TYPE

17. For (a), $\lim_{x\to\pi} \frac{\sin(\pi-x)}{\pi(\pi-x)}$

Using $\sin(\pi - x) = \sin x$, let $t = \pi - x$.

$= \frac{1}{\pi}\lim_{t\to 0}\frac{\sin t}{t} = \frac{1}{\pi}$

So, $a \to P$

For (b), $\lim_{x\to 0} \frac{\cos x}{x-\pi}$

Direct substitution gives

$= \frac{\cos 0}{0-\pi} = \frac{1}{-\pi} = -\frac{1}{\pi}$

So, $b \to R$

For (c), $\lim_{x\to 0} \frac{\cos 2x-1}{\cos x-1}$

Using identities,

$\cos 2x - 1 = -2\sin^2 x$

$\cos x - 1 = -2\sin^2 \frac{x}{2}$

Hence,

Therefore,

$\frac{cos\,2x-1}{cos\,x-1} = \frac{sin^2\,x}{sin^2(x/2)}$

$= \frac{(2sin(x/2)cos(x/2))^2}{sin^2(x/2)}$

$= 4cos^2\frac{x}{2}.$

$As\, x \to 0,$

$4cos^2\frac{x}{2} \to 4.$

So, $c \to Q$

For (d), $\lim_{x\to 0} \frac{ax+x\cos x}{b\sin x}$

$= \frac{x(a+\cos x)}{b\sin x} = \left(\frac{x}{\sin x}\right)\frac{a+\cos x}{b}$

As $x \to 0$,

$\frac{x}{\sin x} \to 1, \cos x \to 1$

Therefore,

$= \frac{a+1}{b}$

So, $d \to S$

INTEGER TYPE

18. If $\ell = \lim_{x\to a} \frac{\sqrt{3x^2+a^2}-\sqrt{x^2+3a^2}}{x-a}$

then $\ell =$

Substituting $x = a$,

$\frac{\sqrt{4a^2}-\sqrt{4a^2}}{a-a} = \frac{0}{0}$

Rationalize:

$\ell = \lim_{x\to a} \frac{(3x^2+a^2)-(x^2+3a^2)}{(x-a)\left(\sqrt{3x^2+a^2}+\sqrt{x^2+3a^2}\right)}$

$= \lim_{x\to a} \frac{2(x^2-a^2)}{(x-a)\left(\sqrt{3x^2+a^2}+\sqrt{x^2+3a^2}\right)}$

$= \lim_{x\to a} \frac{2(x-a)(x+a)}{(x-a)\left(\sqrt{3x^2+a^2}+\sqrt{x^2+3a^2}\right)}$

$= \lim_{x\to a} \frac{2(x+a)}{\sqrt{3x^2+a^2}+\sqrt{x^2+3a^2}}$

Putting $x = a$,

$\ell = \frac{2(2a)}{2a+2a} = 1$

Correct Answer: $\boxed{1}$

19. If $\lim_{n\to\infty}\left(\sqrt{2n^2+n}-\lambda\sqrt{2n^2-n}\right)$

exists, then $\lambda =$

For the limit to exist finitely, coefficients of $n$must cancel.

$\sqrt{2n^2+n} \sim n\sqrt{2}, \sqrt{2n^2-n} \sim n\sqrt{2}$

So expression behaves like

$n\sqrt{2}(1-\lambda)$

For finite limit,

$1-\lambda = 0$

$\lambda = 1$

Correct Answer: $\boxed{1}$

20. Evaluate: $\lim_{x\to 5} \frac{x^2-25}{x-5}$

$x^2-25 = (x-5)(x+5)$

$\lim_{x\to 5} \frac{(x-5)(x+5)}{x-5} = \lim_{x\to 5}(x+5)$

$= 5+5 = 10$

Correct Answer: $\boxed{10}$

STUDENT PRACTISE CORNER

1. Evaluate $\lim_{x\to 1}\left[\frac{2}{1-x}-\frac{4}{1-x^2}\right]$

Factorize:

$1-x^2 = (1-x)(1+x)$

$= \frac{2(1+x)-4}{(1-x)(1+x)}$

$$= \frac{2x-2}{(1-x)(1+x)} = \frac{2(x-1)}{-(x-1)(1+x)}$$

$$= -\frac{2}{1+x}$$

Putting $x = 1$,

$$= -\frac{2}{2} = -1$$

Correct Answer: B) -1

2. Evaluate $\lim_{x\to 4} \frac{\sqrt{1+2x}-3}{\sqrt{x}-2}$

Rationalize numerator:

$$= \frac{1+2x-9}{(\sqrt{x}-2)(\sqrt{1+2x}+3)}$$

$$= \frac{2(x-4)}{(\sqrt{x}-2)(\sqrt{1+2x}+3)}$$

Since

$x - 4 = (\sqrt{x}-2)(\sqrt{x}+2)$

$$= \frac{2(\sqrt{x}+2)}{\sqrt{1+2x}+3}$$

Putting $x = 4$,

$$= \frac{2(2+2)}{3+3} = \frac{8}{6} = \frac{4}{3}$$

Correct Answer: A) $\frac{4}{3}$

3. Evaluate $\lim_{x\to 0} \frac{\sqrt[3]{8+x}-2}{x}$

Use

$$a - b = \frac{a^3 - b^3}{a^2 + ab + b^2}$$

$$= \frac{1}{(\sqrt[3]{8+x})^2 + 2\sqrt[3]{8+x} + 4}$$

Putting $x = 0$,

$$= \frac{1}{4+4+4} = \frac{1}{12}$$

Correct Answer: C) $\frac{1}{12}$

4. Evaluate $\lim_{x\to\infty} \frac{5x^2-7}{3x^2+2}$

Divide numerator and denominator by x^2:

$$= \frac{5 - \frac{7}{x^2}}{3 + \frac{2}{x^2}}$$

As $x \to \infty$,

$\frac{7}{x^2} \to 0, \frac{2}{x^2} \to 0$

$$= \frac{5}{3}$$

Correct Answer: A) $\frac{5}{3}$

5. Evaluate $\lim_{x\to\infty} \left(\frac{x^3}{3x^2-1} - \frac{x^2}{3x+1}\right)$

Take LCM:

$$= \frac{x^3(3x+1) - x^2(3x^2-1)}{(3x^2-1)(3x+1)}$$

$$= \frac{3x^4 + x^3 - 3x^4 + x^2}{(3x^2-1)(3x+1)}$$

$$= \frac{x^3 + x^2}{(3x^2-1)(3x+1)}$$

Factor x^2:

$$= \frac{x^2(x+1)}{(3x^2-1)(3x+1)}$$

Divide by x^3:

$$= \frac{1+\frac{1}{x}}{\left(3-\frac{1}{x^2}\right)\left(3+\frac{1}{x}\right)}$$

As $x \to \infty$,

$$= \frac{1}{3 \cdot 3} = \frac{1}{9}$$

Correct Answer: B) $\frac{1}{9}$

6. Evaluate $\lim_{x\to 0} \frac{(1+x)^5-1}{3x+5x^2}$

Expand using binomial theorem:

$(1+x)^5 = 1 + 5x + 10x^2 + 10x^3 + 5x^4 + x^5$

So,

$(1+x)^5 - 1 = 5x + 10x^2 + 10x^3 + 5x^4 + x^5$

$$= \frac{5x + 10x^2 + \cdots}{3x + 5x^2}$$

Factor x:

$$= \frac{x(5 + 10x + \cdots)}{x(3+5x)}$$

Cancel x:

$$= \frac{5 + 10x + \cdots}{3 + 5x}$$

Putting $x = 0$,

$$= \frac{5}{3}$$

Correct Answer: D) $\frac{5}{3}$

ONE OR MORE THAN ONE CORRECT ANSWER TYPE

7. If $\lim_{x\to 3} f(x) = 7$

then check the statements.

Using limit laws:

$\lim_{x\to 3}[f(x)]^3 = (7)^3 = 343$

So, A is correct

$\lim_{x\to 3}[2f(x) - 1] = 2(7) - 1 = 13$

So, B is correct

$$\lim_{x\to 3} \frac{1}{f(x)} = \frac{1}{7}$$

So, C is correct

$\lim_{x\to 3}[f(x)]^2 = (7)^2 = 49 \neq 14$

So, D is incorrect

Correct Answers: A, B, C

8. If $\lim_{x\to 2} f(x) = 4$

then check the statements.

$\lim_{x\to 2}[5f(x)] = 5(4) = 20$

So, A is correct

$\lim_{x\to 2}[f(x)]^2 = (4)^2 = 16$

So, B is correct

$\lim_{x\to 2}\sqrt{f(x)} = \sqrt{4} = 2$

So, C is correct

$\lim_{x\to 2}\frac{1}{f(x)} = \frac{1}{4} \neq 4$

So, D is incorrect

Correct Answers: A, B, C

9. Evaluate the correct statements for

$$\lim_{x\to\infty}\frac{3x^2+2x+1}{x^2-5}$$

Highest power of x in numerator and denominator is 2.

So, B is correct

Divide numerator and denominator by x^2:

$$=\frac{3+\frac{2}{x}+\frac{1}{x^2}}{1-\frac{5}{x^2}}$$

As $x \to \infty$,

$$=\frac{3}{1}=3$$

So, A is correct

Since division by highest power $x^2$is used,

C is correct

Limit is finite, not infinity.

So, D is incorrect

Correct Answers: A, B, C

COMPREHENSION TYPE

Passage:

10. Evaluate $\lim_{x\to 0}\frac{5^x+5^{-x}-2}{x^2}$

Using expansion:

$$5^x = 1 + x\log_e 5 + \frac{x^2(\log_e 5)^2}{2}$$

$$5^{-x} = 1 - x\log_e 5 + \frac{x^2(\log_e 5)^2}{2}$$

Adding, $5^x + 5^{-x} - 2 = x^2(\log_e 5)^2$

Therefore,

$$\lim_{x\to 0}\frac{x^2(\log_e 5)^2}{x^2} = (\log_e 5)^2$$

Correct Answer: B) $(\log_e 5)^2$

11. Evaluate $\lim_{x\to 0}\frac{3^{2x}-1}{2^{3x}-1}$

Use:

$\lim_{x\to 0}\frac{a^x-1}{x}=\log_e a$

$=\frac{\frac{3^{2x}-1}{x}}{\frac{2^{3x}-1}{x}}$

$=\frac{\log_e 3^2}{\log_e 2^3}=\frac{\log 9}{\log 8}$

Correct Answer: A) $\mathbf{\frac{\log 9}{\log 8}}$

12. Evaluate $\lim_{x\to 1}\frac{2^{x-1}-1}{\sin \pi x}$

Let $h=x-1$

Then $h\to 0$,

$=\lim_{h\to 0}\frac{2^h-1}{\sin \pi(1+h)}$

Since

$\sin(\pi+\pi h)=-\sin \pi h$

$=\lim_{h\to 0}\frac{2^h-1}{-\sin \pi h}$

$=\left(\lim_{h\to 0}\frac{2^h-1}{h}\right)\left(\lim_{h\to 0}\frac{h}{-\sin \pi h}\right)$

$=\log_e 2\cdot\left(\frac{1}{\pi}\right)$

$=-\frac{\log_e 2}{\pi}$

Correct Answer: D) $\mathbf{-\frac{\log_e 2}{\pi}}$

MATRIX MATCHING TYPE

13. Given $\lim_{x\to\infty}\left(\frac{x^3+1}{x+1}-\ell x^2-mx-n\right)=4$

First divide:

$\frac{x^3+1}{x+1}=x^2-x+1$

So,

$x^2-x+1-\ell x^2-mx-n$

$=(1-\ell)x^2+(-1-m)x+(1-n)$

For limit to be finite,

Coefficient of $x^2=0$

$1-\ell=0\Rightarrow \ell=1$

Coefficient of $x=0$

$-1-m=0\Rightarrow m=-1$

Now constant term must be 4:

$1-n=4\Rightarrow n=-3$

Finally,

$\ell mn=(1)(-1)(-3)=3$

Matching:

a) $\ell=1\rightarrow$ R

b) $m=-1\rightarrow$ P

c) $n = -3 \rightarrow$ Q

d) $\ell mn = 3 \rightarrow$ S

INTEGER TYPE

14. If $\lim_{x\to\infty} \frac{9x^2-1}{3x^2+5} = k$

find k.

Divide numerator and denominator by x^2:

$$= \lim_{x\to\infty} \frac{9 - \frac{1}{x^2}}{3 + \frac{5}{x^2}}$$

As $x \to \infty$,

$$= \frac{9}{3} = 3$$

$\boxed{k = 3}$

15. Evaluate $\lim_{x\to 2}(x^2 + 3x - 4)$

Since polynomial functions are continuous, substitute $x = 2$:

$= 2^2 + 3(2) - 4$

$= 4 + 6 - 4 = 6$

$\boxed{6}$