

Chapter 2 : Sequences

A. LOUARDANI

National School of Autonomous Systems technology

Definition

A sequence is a function whose domain is the set of natural numbers \mathbb{N} and range a sub-set of real numbers \mathbb{R} , which is defined as follow

$$\mathbb{N} \rightarrow \mathbb{R}$$

$$n \rightarrow u_n$$

We denote such a function as $(u_n)_{n \geq 0}$ or simply (u_n) .
 u_n is called the general term of the sequence (u_n) .

Examples

Real Numbers

A.
LOUARDANI

- $u_n = 1 + \frac{2}{n+1}$, $u_n = (-1)^n \cos\left(\frac{n\pi}{2}\right)$.

- **Classic Sequences :**

- ♠ **Arithmetic Sequence with common difference r** Its general term is given by

Examples

Real Numbers

A.
LOUARDANI

- $u_n = 1 + \frac{2}{n+1}$, $u_n = (-1)^n \cos(\frac{n\pi}{2})$.
- **Classic Sequences :**

♠ **Arithmetic Sequence with common difference r** Its general term is given by

$$u_n = a + nr \text{ and we have } \sum_{k=1}^n u_k = \frac{n}{2}(u_1 + u_n).$$

♠ **Geometric sequence with a common ratio q .** Its general term is given by

Examples

Real Numbers

A.

LOUARDANI

- $u_n = 1 + \frac{2}{n+1}$, $u_n = (-1)^n \cos(\frac{n\pi}{2})$.

- **Classic Sequences :**

♠ **Arithmetic Sequence with common difference r** Its general term is given by

$$u_n = a + nr \text{ and we have } \sum_{k=1}^n u_k = \frac{n}{2}(u_1 + u_n).$$

♠ **Geometric sequence with a common ratio q .** Its general term is given by

$$u_n = aq^n \text{ and we have } \sum_{k=1}^n u_k = u_1 \frac{1 - q^{n+1}}{1 - q} \quad (q \neq 1).$$

♠ **First-order recursive sequence** given by

Examples

Real Numbers

A.
LOUARDANI

- $u_n = 1 + \frac{2}{n+1}$, $u_n = (-1)^n \cos(\frac{n\pi}{2})$.
- **Classic Sequences :**

♠ **Arithmetic Sequence with common difference r** Its general term is given by

$$u_n = a + nr \text{ and we have } \sum_{k=1}^n u_k = \frac{n}{2}(u_1 + u_n).$$

♠ **Geometric sequence with a common ratio q .** Its general term is given by

$$u_n = aq^n \text{ and we have } \sum_{k=1}^n u_k = u_1 \frac{1 - q^{n+1}}{1 - q} \quad (q \neq 1).$$

♠ **First-order recursive sequence** given by

$$\begin{cases} u_0 = a \in \mathbb{R} \\ u_{n+1} = f(u_n) \end{cases}$$

♠ **Second-order recursive sequence** given by

Examples

Real Numbers

A.
LOUARDANI

- $u_n = 1 + \frac{2}{n+1}$, $u_n = (-1)^n \cos(\frac{n\pi}{2})$.

- **Classic Sequences :**

♠ **Arithmetic Sequence with common difference r** Its general term is given by

$$u_n = a + nr \text{ and we have } \sum_{k=1}^n u_k = \frac{n}{2}(u_1 + u_n).$$

♠ **Geometric sequence with a common ratio q .** Its general term is given by

$$u_n = aq^n \text{ and we have } \sum_{k=1}^n u_k = u_1 \frac{1 - q^{n+1}}{1 - q} \quad (q \neq 1).$$

♠ **First-order recursive sequence** given by

$$\begin{cases} u_0 = a \in \mathbb{R} \\ u_{n+1} = f(u_n) \end{cases}$$

♠ **Second-order recursive sequence** given by

$$\begin{cases} u_0 = \alpha, \quad u_1 = \beta \end{cases}$$

Definition 2.

We say that the sequence (u_n) is :

- increasing if and only if

$$\forall n \in \mathbb{N}, \quad u_{n+1} \geq u_n$$

Definition 2.

We say that the sequence (u_n) is :

- increasing if and only if

$$\forall n \in \mathbb{N}, \quad u_{n+1} \geq u_n$$

- decreasing if and only if

$$\forall n \in \mathbb{N}, \quad u_{n+1} \leq u_n$$

Definition 2.

We say that the sequence (u_n) is :

- increasing if and only if

$$\forall n \in \mathbb{N}, \quad u_{n+1} \geq u_n$$

- decreasing if and only if

$$\forall n \in \mathbb{N}, \quad u_{n+1} \leq u_n$$

- strictly increasing if and only if

$$\forall n \in \mathbb{N}, \quad u_{n+1} > u_n$$

Definition 2.

We say that the sequence (u_n) is :

- increasing if and only if

$$\forall n \in \mathbb{N}, \quad u_{n+1} \geq u_n$$

- decreasing if and only if

$$\forall n \in \mathbb{N}, \quad u_{n+1} \leq u_n$$

- strictly increasing if and only if

$$\forall n \in \mathbb{N}, \quad u_{n+1} > u_n$$

- strictly decreasing if and only if

$$\forall n \in \mathbb{N}, \quad u_{n+1} < u_n$$

Definition 2.

We say that the sequence (u_n) is :

- increasing if and only if

$$\forall n \in \mathbb{N}, \quad u_{n+1} \geq u_n$$

- decreasing if and only if

$$\forall n \in \mathbb{N}, \quad u_{n+1} \leq u_n$$

- strictly increasing if and only if

$$\forall n \in \mathbb{N}, \quad u_{n+1} > u_n$$

- strictly decreasing if and only if

$$\forall n \in \mathbb{N}, \quad u_{n+1} < u_n$$

Definition 2

- monotonic if and only if it is either increasing or decreasing.

Definition 2

- monotonic if and only if it is either increasing or decreasing.
- strictly monotonic if and only if it is either strictly increasing or strictly decreasing.

Example 2.

Consider the sequence $u_n = \sum_{k=1}^n \frac{1}{k^2}$, let's study its monotonicity.

Definition 2

- monotonic if and only if it is either increasing or decreasing.
- strictly monotonic if and only if it is either strictly increasing or strictly decreasing.

Example 2.

Consider the sequence $u_n = \sum_{k=1}^n \frac{1}{k^2}$, let's study its monotonicity. We have

$$u_{n+1} - u_n = \sum_{k=1}^{n+1} \frac{1}{k^2} - \sum_{k=1}^n \frac{1}{k^2} = \frac{1}{(n+1)^2} > 0 \quad \forall n \geq 1.$$

The sequence (u_n) is strictly increasing.

Definition 2

- monotonic if and only if it is either increasing or decreasing.
- strictly monotonic if and only if it is either strictly increasing or strictly decreasing.

Example 2.

Consider the sequence $u_n = \sum_{k=1}^n \frac{1}{k^2}$, let's study its monotonicity. We have

$$u_{n+1} - u_n = \sum_{k=1}^{n+1} \frac{1}{k^2} - \sum_{k=1}^n \frac{1}{k^2} = \frac{1}{(n+1)^2} > 0 \quad \forall n \geq 1.$$

The sequence (u_n) is strictly increasing.

$$2. \quad u_n = \sum_{k=1}^n \frac{1}{n+k}.$$

Definition 3

Let (u_n) be a sequence of real numbers. We say that the sequence (u_n) is :

- Upper bounded if the subset of \mathbb{R} , $A = \{u_n, n \in \mathbb{N}\}$ is bounded above , i.e.,

$$\exists M \in \mathbb{R}, \forall n \in \mathbb{N}, u_n \leq M$$

Definition 3

Let (u_n) be a sequence of real numbers. We say that the sequence (u_n) is :

- Upper bounded if the subset of \mathbb{R} , $A = \{u_n, n \in \mathbb{N}\}$ is bounded above , i.e.,

$$\exists M \in \mathbb{R}, \forall n \in \mathbb{N}, u_n \leq M$$

- lower bounded if the subset of \mathbb{R} , $A = \{u_n, n \in \mathbb{N}\}$ is bounded below, i.e.,

$$\exists m \in \mathbb{R}, \forall n \in \mathbb{N}, u_n \geq m$$

Definition 3

Let (u_n) be a sequence of real numbers. We say that the sequence (u_n) is :

- Upper bounded if the subset of \mathbb{R} , $A = \{u_n, n \in \mathbb{N}\}$ is bounded above , i.e.,

$$\exists M \in \mathbb{R}, \forall n \in \mathbb{N}, u_n \leq M$$

- lower bounded if the subset of \mathbb{R} , $A = \{u_n, n \in \mathbb{N}\}$ is bounded below, i.e.,

$$\exists m \in \mathbb{R}, \forall n \in \mathbb{N}, u_n \geq m$$

- bounded if and only if it is both lower and upper bounded.

Example 3 Show that the sequence $u_n = \sum_{k=1}^n \frac{1}{k+n}$.

Example 3 Show that the sequence $u_n = \sum_{k=1}^n \frac{1}{k+n}$.

Example 4. Show that the sequence $u_n = \sum_{k=1}^n \frac{1}{k^2}$ is upper bounded. We have for $2 \leq k \leq n$

$$\frac{1}{k^2} \leq \frac{1}{k(k-1)} = \frac{1}{k-1} - \frac{1}{k}$$

Thus,

Example 3 Show that the sequence $u_n = \sum_{k=1}^n \frac{1}{k+n}$.

Example 4. Show that the sequence $u_n = \sum_{k=1}^n \frac{1}{k^2}$ is upper bounded. We have for $2 \leq k \leq n$

$$\frac{1}{k^2} \leq \frac{1}{k(k-1)} = \frac{1}{k-1} - \frac{1}{k}$$

Thus,

$$u_n \leq 1 + \sum_{k=2}^n \left(\frac{1}{k-1} - \frac{1}{k} \right) = 1 + \left(1 - \frac{1}{2} \right) + \left(\frac{1}{2} - \frac{1}{3} \right) + \dots +$$

$$\left(\frac{1}{n-2} - \frac{1}{n-1} \right) + \left(\frac{1}{n-1} - \frac{1}{n} \right) = 2 - \frac{1}{n} \leq 2$$

Thus, for all $n \in \mathbb{N}^*$, the sequence $u_n \leq 2$.

Proposition.

The sequence (u_n) is bounded if and only if the sequence $(|u_n|)$ is upper bounded , i.e., there exists $M \in \mathbb{R}$ such that $|u_n| \leq M$ for all $n \in \mathbb{N}$.

Proof.

Proof. \Leftarrow The sequence $|u_n|$ is upper bounded, which means there exists $M \in \mathbb{R}$ such that $|u_n| \leq M \Leftrightarrow -M \leq u_n \leq M$, meaning that (u_n) is bounded.

Proof. \Leftarrow The sequence $|u_n|$ is upper bounded, which means there exists $M \in \mathbb{R}$ such that $|u_n| \leq M \Leftrightarrow -M \leq u_n \leq M$, meaning that (u_n) is bounded. \Rightarrow

Proof. \Leftarrow The sequence $|u_n|$ is upper bounded, which means there exists $M \in \mathbb{R}$ such that $|u_n| \leq M \Leftrightarrow -M \leq u_n \leq M$, meaning that (u_n) is bounded. \Rightarrow The sequence (u_n) is bounded, meaning there exist M and m in \mathbb{R} such that $m \leq u_n \leq M$ for all $n \in \mathbb{N}$.

Proof. \Leftarrow The sequence $|u_n|$ is upper bounded, which means there exists $M \in \mathbb{R}$ such that $|u_n| \leq M \Leftrightarrow -M \leq u_n \leq M$, meaning that (u_n) is bounded. \Rightarrow The sequence (u_n) is bounded, meaning there exist M and m in \mathbb{R} such that $m \leq u_n \leq M$ for all $n \in \mathbb{N}$. We take $k = \max(|m|, |M|)$, which means that $|u_n| \leq k$.

Proof. \Leftarrow The sequence $|u_n|$ is upper bounded, which means there exists $M \in \mathbb{R}$ such that $|u_n| \leq M \Leftrightarrow -M \leq u_n \leq M$, meaning that (u_n) is bounded. \Rightarrow The sequence (u_n) is bounded, meaning there exist M and m in \mathbb{R} such that $m \leq u_n \leq M$ for all $n \in \mathbb{N}$. We take $k = \max(|m|, |M|)$, which means that $|u_n| \leq k$.

Note. We say that the sequence (u_n) is bounded (upper bounded, lower bounded) from some term onwards if there exist $N \in \mathbb{N}$ and $M \in \mathbb{R}$ (or $m \in \mathbb{R}$) such that for all $n \geq N$, we have $u_n \leq M$ (or $u_n \geq m$).

Convergent Sequence

Real Numbers

A.
LOUARDANI

Definition 4. We say that the sequence $(u_n)_{n \geq 0}$ has the limit $l \in \mathbb{R}$ if and only if for every

$\forall \epsilon > 0, \exists N \in \mathbb{N}$, such that


$\forall n \in \mathbb{N}, n \geq N$ implies $|u_n - l| \leq \epsilon$

In other words, from a certain rank N , the terms of the sequence (u_n) get closer to the limit l (the terms of the sequence are in the interval $[l - \epsilon, l + \epsilon]$).

If such an l exists, we say that the sequence (u_n) is convergent (or the sequence (u_n) converges to l), and we write

$$\lim_{n \rightarrow \infty} u_n = l \text{ or } \lim u_n = l \text{ or } u_n \rightarrow l$$

Otherwise, we say that the sequence (u_n) is divergent.



Convergent Sequence

Real Numbers

A.
LOUARDANI

Remark

- If the sequence (u_n) has infinity as its limit, we say it diverges, and the divergence is of the first kind.

Convergent Sequence

Real Numbers

A.
LOUARDANI

Remark

- If the sequence (u_n) has infinity as its limit, we say it diverges, and the divergence is of the first kind.
- If the sequence (u_n) has no limit, we say it diverges, and the divergence is of the second kind.

Convergent Sequence

Real Numbers

A.
LOUARDANI

Remark

- If the sequence (u_n) has infinity as its limit, we say it diverges, and the divergence is of the first kind.
- If the sequence (u_n) has no limit, we say it diverges, and the divergence is of the second kind.

Example 4. Show that the sequence $u_n = \frac{1}{n}$ ($n \in \mathbb{N}^*$) converges to 0.

Convergent Sequence

Real Numbers

A.
LOUARDANI

Remark

- If the sequence (u_n) has infinity as its limit, we say it diverges, and the divergence is of the first kind.
- If the sequence (u_n) has no limit, we say it diverges, and the divergence is of the second kind.

Example 4. Show that the sequence $u_n = \frac{1}{n}$ ($n \in \mathbb{N}^*$) converges to 0.

Let $\epsilon > 0$, we want to find $N = N(\epsilon) \in \mathbb{N}$ such that $|u_n - l| \leq \epsilon$. We often reduce this to solving inequalities.

We have $|u_n - l| = \frac{1}{n} \leq \epsilon \Leftrightarrow n \geq \frac{1}{\epsilon}$. We take $N = E(\frac{1}{\epsilon}) + 1$.

Thus, for $n \geq N$ we have $|u_n - 0| = \frac{1}{n} \leq \frac{1}{N} \leq \epsilon$.

Convergent Sequence

Real Numbers

A.
LOUARDANI

Remark. Instead of using a non-strict inequality in the definition of a convergent sequence, we can use a strict inequality, meaning :

$$u_n \rightarrow l \Leftrightarrow \forall \varepsilon > 0, \exists N \in \mathbb{N}, \forall n \geq N \text{ we have } |u_n - l| < \varepsilon.$$

Convergent Sequence

Real Numbers

A.
LOUARDANI

Remark. Instead of using a non-strict inequality in the definition of a convergent sequence, we can use a strict inequality, meaning :

$$u_n \rightarrow l \Leftrightarrow \forall \varepsilon > 0, \exists N \in \mathbb{N}, \forall n \geq N \text{ we have } |u_n - l| < \varepsilon.$$

Indeed, the implication in the reverse direction \Leftarrow is obvious.

Convergent Sequence

Real Numbers

A.
LOUARDANI

Remark. Instead of using a non-strict inequality in the definition of a convergent sequence, we can use a strict inequality, meaning :

$$u_n \rightarrow l \Leftrightarrow \forall \varepsilon > 0, \exists N \in \mathbb{N}, \forall n \geq N \text{ we have } |u_n - l| < \varepsilon.$$

Indeed, the implication in the reverse direction \Leftarrow is obvious. For the other direction of the implication (\Rightarrow), let $\varepsilon > 0$, we set $\varepsilon' = \frac{\varepsilon}{2} > 0$ (which is arbitrary), by the definition, we have

$$u_n \rightarrow l \Rightarrow \exists N \in \mathbb{N}, \forall n \geq N, |u_n - l| \leq \varepsilon' < \varepsilon$$

Convergent Sequence

Real Numbers

A.
LOUARDANI

Theorem 1.

The limit $l \in \mathbb{R}$ of a real sequence, if it exists, is unique.

Convergent Sequence

Real Numbers

A.
LOUARDANI

Theorem 1.

The limit $l \in \mathbb{R}$ of a real sequence, if it exists, is unique.

Proof. We assume by contradiction that the sequence (u_n) has two different limits l_1 and l_2 ($l_1 \neq l_2$).

Convergent Sequence

Real Numbers

A.
LOUARDANI

Theorem 1.

The limit $l \in \mathbb{R}$ of a real sequence, if it exists, is unique.

Proof. We assume by contradiction that the sequence (u_n) has two different limits l_1 and l_2 ($l_1 \neq l_2$).

Let $\epsilon = \frac{|l_1 - l_2|}{2} > 0$.

- Since $u_n \rightarrow l_1$ then there exists $N_1 \in \mathbb{N}$ from which we have $|u_n - l_1| < \epsilon$.
- Since $u_n \rightarrow l_2$ then there exists $N_2 \in \mathbb{N}$ from which we have $|u_n - l_2| < \epsilon$.

Now we consider the integer $N = \max(N_1, N_2)$, from this rank, both of the last two inequalities $|u_n - l_1| < \epsilon$, $|u_n - l_2| < \epsilon$ are satisfied. Using the triangle inequality, we obtain

$$|l_1 - l_2| = |l_1 - u_n + u_n - l_2| \leq |u_n - l_1| + |u_n - l_2| < 2\epsilon = |l_1 - l_2|$$

which is absurd.

Convergent Sequence

Real Numbers

A.
LOUARDANI

Theorem 2.

Every convergent sequence is bounded.

Convergent Sequence

Real Numbers

A.
LOUARDANI

Theorem 2.

Every convergent sequence is bounded.

Proof. Let (u_n) be a real sequence that converges to l . By definition, we have

$$\forall \epsilon > 0, \exists N \in \mathbb{N}, \forall n \geq N, |u_n - l| \leq \epsilon \Leftrightarrow -\epsilon \leq u_n - l \leq \epsilon \Leftrightarrow$$

$$l - \epsilon \leq u_n \leq l + \epsilon,$$

which means that $|u_n| \leq |l + \epsilon|$ or $|u_n| \leq |l - \epsilon|$. Therefore, let $M = \max(|l + \epsilon|, |l - \epsilon|)$, we have for every $n \geq N$, $|u_n| \leq M$.

The first terms of the sequence remain, so we set

$M' = \max(|u_0|, |u_1|, \dots, |u_{N-1}|, M)$, we have $|u_n| \leq M'$ for every $n \in \mathbb{N}$.

Properties. Let (u_n) and (v_n) be two real sequences with respective limits l_1, l_2 . The table below summarizes the properties of the limits of the sum, product, and quotient of two sequences

$\lim u_n$	$\lim v_n$	$\lim(u_n + v_n)$	$\lim(u_n \times v_n)$	$\lim \frac{u_n}{v_n}$
l_1	l_2	$l_1 + l_2$	$l_1 \times l_2$	$\frac{l_1}{l_2} \ (l_2 \neq 0)$
0	l_2	l_2	0	0
l_1	0	l_1	0	∞
0	0	0	0	F.I
∞	0	∞	F.I	∞
0	∞	∞	F.I	0
l_1	∞	∞	∞	0

$\lim u_n$	$\lim v_n$	$\lim(u_n + v_n)$	$\lim(u_n \times v_n)$	$\lim \frac{u_n}{v_n}$
∞	l_2	∞	∞	∞
$+\infty$	$+\infty$	$+\infty$	$+\infty$	F.I
$-\infty$	$-\infty$	$-\infty$	$+\infty$	F.I
$+\infty$	$-\infty$	F.I	$-\infty$	F.I

Positive Limit

Real Numbers

A.
LOUARDANI

Proposition Let (u_n) be a convergent real sequence, such that :

- i) There exists $N \in \mathbb{N}$ from which $u_n > 0$
- ii) $\lim_{n \rightarrow \infty} u_n = l$

Then $l \geq 0$.

Positive Limit

Real Numbers

A.
LOUARDANI

Proposition Let (u_n) be a convergent real sequence, such that :

- i) There exists $N \in \mathbb{N}$ from which $u_n > 0$
- ii) $\lim_{n \rightarrow \infty} u_n = l$

Then $l \geq 0$.

Proof Let's assume, by contradiction, that $l < 0$. Set $\varepsilon = -l > 0$. Since $u_n \rightarrow l$, there exists $N_1 \in \mathbb{N}$ such that $|u_n - l| \leq \varepsilon$, and there exists N_2 such that for all $n \geq N_2$ we have $u_n > 0$. Let's take $N = \max(N_1, N_2)$, so for all $n \geq N$, we have :

$$l \leq u_n - l \leq -l \text{ and } u_n > 0 \Leftrightarrow 2l \leq u_n \leq 0 \text{ and } u_n > 0$$

which is absurd.

Passing to the Limit in Inequalities

Real Numbers

A.
LOUARDANI

Proposition Let (u_n) and (v_n) be two real sequences, such that :

i) $u_n \rightarrow l$

ii) $v_n \rightarrow l'$

iii) There exists $N \in \mathbb{N}$ such that for all $n \geq N$ we have $u_n \leq v_n$.

Then $l \leq l'$.

Proof We can proceed in a similar way to the previous proof. We just need to consider the sequence $w_n = v_n - u_n$.

Squeeze theorem

Real Numbers

A.
LOUARDANI

Theorem 3.

Let (u_n) , (v_n) , and (w_n) be three real sequences, such that :

i) $\lim v_n = \lim w_n = l$

ii) There exists $N \in \mathbb{N}$ such that for all $n \geq N$ we have
 $v_n \leq u_n \leq w_n$.

Then $\lim u_n = l$.

Squeeze theorem

Real Numbers

A.
LOUARDANI

Proof. We have :

- $v_n \rightarrow l$, so there exists $N_1 \in \mathbb{N}$ from which we have $|v_n - l| \leq \varepsilon \Leftrightarrow l - \varepsilon \leq v_n \leq l + \varepsilon$
- $w_n \rightarrow l$, so there exists $N_2 \in \mathbb{N}$ from which we have $|w_n - l| \leq \varepsilon \Leftrightarrow l - \varepsilon \leq w_n \leq l + \varepsilon$
- There exists $N_3 \in \mathbb{N}$ such that $n \geq N$ implies $v_n \leq u_n \leq w_n$.
Let's take $N = \max(N_1, N_2, N_3)$. Then, for all $n \geq N$, we have :

$$l - \varepsilon \leq v_n \leq u_n \leq w_n \leq l + \varepsilon$$

so

$$-\varepsilon \leq u_n - l \leq \varepsilon$$

Consequently, for all $n \geq N$, we have $|u_n - l| \leq \varepsilon$.

Squeeze theorem

Real Numbers

A.
LOUARDANI

Corollary Let (u_n) and (v_n) be two numerical sequences, such that :

- The sequence (u_n) converges to 0.
- The sequence (v_n) is bounded.

Then, the sequence $w_n = u_n v_n$ converges to 0.

Proof It is enough to apply Gendarme's Theorem to the sequence (w_n) .

Real Sequences and Monotonicity

Real Numbers

A.
LOUARDANI

Theorem 4

(Convergence Criterion for Monotonic Sequences)

Let (u_n) be a monotonic sequence, then :

- If it is increasing, it is convergent if and only if it is bounded. Moreover, $\lim u_n = \sup\{u_n, n \in \mathbb{N}\}$.
- If it is decreasing, it is convergent if and only if it is bounded. Moreover, $\lim u_n = \inf\{u_n, n \in \mathbb{N}\}$.

Real Sequences and Monotonicity

Real Numbers

A.
LOUARDANI

Remark

- If the sequence (u_n) is increasing and unbounded, then $\lim u_n = +\infty$.

Real Sequences and Monotonicity

Real Numbers

A.
LOUARDANI

Remark

- If the sequence (u_n) is increasing and unbounded, then $\lim u_n = +\infty$.
- If the sequence (u_n) is decreasing and unbounded, then $\lim u_n = -\infty$.

Real Sequences and Monotonicity

Real Numbers

A.
LOUARDANI

Remark

- If the sequence (u_n) is increasing and unbounded, then $\lim u_n = +\infty$.
- If the sequence (u_n) is decreasing and unbounded, then $\lim u_n = -\infty$.
- If the increasing sequence (u_n) converges to l , then for all $n \in \mathbb{N}$, $u_n \leq l$.

Real Sequences and Monotonicity

Real Numbers

A.
LOUARDANI

Remark

- If the sequence (u_n) is increasing and unbounded, then $\lim u_n = +\infty$.
- If the sequence (u_n) is decreasing and unbounded, then $\lim u_n = -\infty$.
- If the increasing sequence (u_n) converges to l , then for all $n \in \mathbb{N}$, $u_n \leq l$.
- If the decreasing sequence (u_n) converges to l , then for all $n \in \mathbb{N}$, $u_n \geq l$.

Real Sequences and Monotonicity

Real Numbers

A.
LOUARDANI

Proof We define A as a subset of \mathbb{R} by $A = \{u_n, n \in \mathbb{N}\}$. A is bounded, so according to the supremum property, it has a supremum l . We will show that $l = \lim u_n$.

Using the supremum property, for all $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that $l - \varepsilon < u_N \leq l$. As the sequence (u_n) is increasing, for all $n \geq N$ we have $u_n \geq u_N$. Since l is an upper bound for A , we have :

$$l - \varepsilon \leq u_N \leq u_n \leq l \Leftrightarrow -\varepsilon \leq u_n - l \leq 0 \Leftrightarrow |u_n - l| \leq \varepsilon.$$

Real Sequences and Monotonicity

Real Numbers

A.
LOUARDANI

Example 5

1- We have seen in the previous examples that the sequence $u_n = \sum_{k=0}^n \frac{1}{k^2}$ is bounded by 2 and increasing.

Real Sequences and Monotonicity

Real Numbers

A.
LOUARDANI

Example 5

1- We have seen in the previous examples that the sequence $u_n = \sum_{k=0}^n \frac{1}{k^2}$ is bounded by 2 and increasing.

Therefore, according to the monotone sequence convergence criterion, the sequence (u_n) is convergent.

Real Sequences and Monotonicity

Real Numbers

A.
LOUARDANI

2- Study the nature of the numerical sequence (u_n) defined as :

$$u_n = \sum_{k=1}^n \frac{1}{n+k}$$

We have :

$$u_{n+1} - u_n = \sum_{k=1}^{n+1} \frac{1}{n+1+k} - \sum_{k=1}^n \frac{1}{n+k} = -\frac{1}{n+1} + \frac{1}{2n+2} + \frac{1}{2n+1} - \frac{1}{2n+1}$$
$$\frac{1}{2(n+1)(2n+1)} > 0 \text{ for all } n \in \mathbb{N}$$

So, the sequence (u_n) is strictly increasing.

Real Sequences and Monotonicity

Real Numbers

A.
LOUARDANI

2- Study the nature of the numerical sequence (u_n) defined as :

$$u_n = \sum_{k=1}^n \frac{1}{n+k}$$

We have :

$$u_{n+1} - u_n = \sum_{k=1}^{n+1} \frac{1}{n+1+k} - \sum_{k=1}^n \frac{1}{n+k} = -\frac{1}{n+1} + \frac{1}{2n+2} + \frac{1}{2n+1} - \frac{1}{2n+1}$$
$$\frac{1}{2(n+1)(2n+1)} > 0 \text{ for all } n \in \mathbb{N}$$

So, the sequence (u_n) is strictly increasing. Furthermore, we have, for all $1 \leq k \leq n$:

$$u_n \leq \sum_{k=1}^n \frac{1}{n+1} = \frac{n}{n+1} < 1$$

Real Sequences and Monotonicity

Real Numbers

A.
LOUARDANI

Using the following inequalities :

$$\ln\left(\frac{x+1}{x}\right) \leq \frac{1}{x} \leq \ln\left(\frac{x}{x-1}\right) \text{ for all } x > 1$$

Calculate the limit of (u_n) .

Real Sequences and Monotonicity

Real Numbers

A.
LOUARDANI

Using the following inequalities :

$$\ln\left(\frac{x+1}{x}\right) \leq \frac{1}{x} \leq \ln\left(\frac{x}{x-1}\right) \text{ for all } x > 1$$

Calculate the limit of (u_n) .

For $1 \leq k \leq n$, we have :

$$\ln\left(\frac{n+k+1}{n+k}\right) \leq \frac{1}{n+k} \leq \ln\left(\frac{n+k}{n+k-1}\right)$$

Summing for k from 1 to n , we get :

$$\sum_{k=1}^n \ln\left(\frac{n+k+1}{n+k}\right) \leq u_n \leq \sum_{k=1}^n \ln\left(\frac{n+k}{n+k-1}\right)$$

$$\Leftrightarrow \sum_{k=1}^n \ln(n+k+1) - \ln(n+k) \leq u_n \leq \sum_{k=1}^n \ln(n+k) - \ln(n+k-1)$$

The sequences $v_n = \sum_{k=1}^n \ln(n+k+1) - \ln(n+k)$ and $w_n = \sum_{k=1}^n \ln(n+k) - \ln(n+k-1)$ are called telescopic series, and we can easily calculate their sums. In fact, we have :

$$\begin{aligned} v_n &= \sum_{k=1}^n \ln(n+k+1) - \ln(n+k) = (\ln(n+2) - \ln(n+1)) + \\ &(\ln(n+3) - \ln(n+2)) + \dots + (\ln(2n) - \ln(2n-1)) + (\ln(2n+1) - \ln(2n)) \\ &= \ln(2n+1) - \ln(n+1) = \ln\left(\frac{2n+1}{n+1}\right) \rightarrow \ln(2) \end{aligned}$$

Doing the same for the sequence (w_n) , we find that $w_n \rightarrow \ln(2)$. According to the squeeze theorem, we deduce that the sequence (u_n) converges to $\ln(2)$.

Adjacent Sequences

Real Numbers

A.
LOUARDANI

Definition 4. Let (u_n) and (v_n) be two real sequences. We say that they are adjacent if and only if :

1. (u_n) is increasing.
2. (v_n) is decreasing.
3. $\lim(u_n - v_n) = 0$.

Adjacent Sequences

Real Numbers

A.
LOUARDANI

Theorem 5.

If (u_n) and (v_n) are two adjacent sequences, then :

- The sequences (u_n) and (v_n) converge to the same limit.
- For all $n \in \mathbb{N}$, we have $u_n \leq l \leq v_n$, where l is the common limit.

Adjacent Sequences

Real Numbers

A.
LOUARDANI

Proof. Let (u_n) be an increasing sequence, and (v_n) be a decreasing sequence. We have :

$$u_n \leq u_{n+1} \Leftrightarrow -u_n \geq -u_{n+1} \text{ and } v_n \geq v_{n+1}$$

Thus, $v_n - u_n \geq v_{n+1} - u_{n+1}$, which means that the sequence $v_n - u_n$ is decreasing.

Since $\lim(u_n - v_n) = 0$, we have $v_n - u_n \geq 0 \Leftrightarrow v_n \geq u_n$.

We now have :

$$u_n \leq v_n \leq v_0 \longrightarrow$$

The sequence (u_n) is increasing and bounded, so it converges to l and

$$v_n \geq u_n \geq u_0 \longrightarrow \text{The sequence } (v_n)$$

is decreasing and bounded, so it converges to l' .

Since $\lim(v_n - u_n) = 0$, we have $l - l' = 0 \Leftrightarrow l = l'$.

Adjacent Sequences

Real Numbers

A.
LOUARDANI

Example 6.

Consider the sequence $w_n = \sum_{k=1}^n \frac{(-1)^{k+1}}{k}$. Show that the sequences $u_n = w_{2n}$ and $v_n = w_{2n+1}$ are adjacent.

Adjacent Sequences

Real Numbers

A.
LOUARDANI

Example 6.

Consider the sequence $w_n = \sum_{k=1}^n \frac{(-1)^{k+1}}{k}$. Show that the sequences $u_n = w_{2n}$ and $v_n = w_{2n+1}$ are adjacent.

Monotonicity of (u_n) and (v_n) : We have :

$$\begin{aligned} u_{n+1} - u_n &= w_{2n+2} - w_{2n} = \sum_{k=1}^{2n+2} \frac{(-1)^{k+1}}{k} - \sum_{k=1}^{2n} \frac{(-1)^{k+1}}{k} \\ &= \frac{1}{2n+1} - \frac{1}{2n+2} \geq 0, \quad \forall n \in \mathbb{N}^* \end{aligned}$$

and

$$\begin{aligned} v_{n+1} - v_n &= w_{2n+3} - w_{2n+1} = \sum_{k=1}^{2n+3} \frac{(-1)^{k+1}}{k} - \sum_{k=1}^{2n+1} \frac{(-1)^{k+1}}{k} \\ &= -\frac{1}{2n+2} + \frac{1}{2n+3} \leq 0, \quad \forall n \in \mathbb{N}^* \end{aligned}$$

Thus, the sequence (u_n) is increasing, and the sequence (v_n) is

Furthermore, we have :

$$u_n - v_n = \frac{1}{2n+1} \rightarrow 0$$

Therefore, the sequences (u_n) and (v_n) are adjacent, and they converge to the same limit.

Subsequence of a Sequence

Real Numbers

A.
LOUARDANI

Definition 6. Let (u_n) be a numerical sequence. We call a subsequence of (u_n) a sequence $(u_{\phi(n)})$ where $\phi : \mathbb{N} \rightarrow \mathbb{N}$ is a strictly increasing function.

Subsequence of a Sequence

Real Numbers

A.
LOUARDANI

Definition 6. Let (u_n) be a numerical sequence. We call a subsequence of (u_n) a sequence $(u_{\phi(n)})$ where $\phi : \mathbb{N} \rightarrow \mathbb{N}$ is a strictly increasing function.

Example 7.

- For $u_n = \frac{1}{n}$, we have $u_{n^2} = \frac{1}{n^2}$.

Subsequence of a Sequence

Real Numbers

A.
LOUARDANI

Definition 6. Let (u_n) be a numerical sequence. We call a subsequence of (u_n) a sequence $(u_{\phi(n)})$ where $\phi : \mathbb{N} \rightarrow \mathbb{N}$ is a strictly increasing function.

Example 7.

- For $u_n = \frac{1}{n}$, we have $u_{n^2} = \frac{1}{n^2}$.
- For $u_n = (-1)^n$, we have $u_{2n} = 1$ and $u_{2n+1} = -1$.

Subsequence of a Sequence

Real Numbers

A.
LOUARDANI

Definition 6. Let (u_n) be a numerical sequence. We call a subsequence of (u_n) a sequence $(u_{\phi(n)})$ where $\phi : \mathbb{N} \rightarrow \mathbb{N}$ is a strictly increasing function.

Example 7.

- For $u_n = \frac{1}{n}$, we have $u_{n^2} = \frac{1}{n^2}$.
- For $u_n = (-1)^n$, we have $u_{2n} = 1$ and $u_{2n+1} = -1$.
- For $u_n = \sin(\frac{2n\pi}{17})$, we have $u_{17n} = \sin(2n\pi) = 0$ and $u_{17n+1} = \sin(\frac{2\pi}{17}) \neq 0$.

Proposition.

Let (u_n) be a numerical sequence. Then, (u_n) converges to l if and only if every subsequence of (u_n) also converges to l .

Proof \Leftarrow This direction is obvious. We just need to take the subsequence $(u_{\phi(n)})$ with $\phi(n) = n$.

\Rightarrow To prove this direction, we need to show that for every $n \in \mathbb{N}$, we have $\phi(n) \geq n$. We will proceed by induction. For $n = 0$, we have $\phi(0) \geq 0$ because $\phi(0) \in \mathbb{N}$.

Suppose that $\phi(n) \geq n$ and let's show that $\phi(n+1) \geq n+1$. Since ϕ is strictly increasing, for $n+1 > n$, we have $\phi(n+1) > \phi(n) \geq n$. Thus, $\phi(n+1) \geq n+1$ ($\phi(n+1), n+1 \in \mathbb{N}$).

The sequence (u_n) converges to l , which means that for every $\varepsilon > 0$, there exists an $N \in \mathbb{N}$ such that for all $n \geq N$, we have $|u_n - l| \leq \varepsilon$. But we also have $\phi(n) \geq n \geq N$, so there exists $N \in \mathbb{N}$ such that $\phi(n) \geq N$, and in this case, we have $|u_{\phi(n)} - l| \leq \varepsilon$.

This proposition is often used to show that a sequence (u_n) is divergent.

Corollary

Let (u_n) be a numerical sequence. If subsequence $(u_{\psi(n)})$ diverges, then the sequence (u_n) also diverges.

Example 8. Consider the sequence $u_n = \cos(\frac{n\pi}{3})$. If we take the subsequence (u_{3n}) , we have $u_{3n} = \cos(n\pi) = (-1)^n$, which diverges. Thus, the sequence (u_n) also diverges.