1. The probability of a leap year selected at random contain 53									
Sunday is:									
(a) 53/366	(b) 1/7	(c) 2/7	(d) 53/365						
_			marble is drawn at						
random. The prob	-	_							
(a) 3/5 (b) 2/5 (c) 0/5 (d) 1/5									
3. The probability			0.85. What is the						
probability that it									
• •	1 7		(d) none of these						
•	•		ted from the numbers						
(1, 2, 3,,15)			4.0						
(a) 1/5	(b) 4/5	(c) 2/15	(d) 1/3						
5. What are the t	otal outcomes	when we thro	w three coins? {HHH,HHT,THH,HTH						
(a) 4	(D) 5	(C) 8	(a) / ,TTT,HTT,THT,TTH}						
-	•	number select	ted at random from the						
numbers (1,2,3,	35) is:	() 10 (0)	- (I) (II						
			5 (d) none of these						
7. The sum of the	_								
` '	(b) 1 (c)	` '							
•	•	re given; choo	se the correct answer						
for that which is r		() 7.15	(1)						
* *	, ,		(d) none of these.						
		litaneousiy, tr	nan the probability of						
getting at least tw	(b) 3/8	(0) 1/	(A) 1 (O)						
10. A letter is ch									
(a) 6/12	(b) 7/12		e letter chosen has: (d) none of these.						
(a) 0/13	(b) // 13	(6) 1	(d) Hotte of tilese.						
11 A dica is throu	11. A dice is thrown. Find the probability of getting an even number.								
	_		(D) 1/2						
(A) 2/3	(D) I	(C) 3/0	(D) 1/2						
12. Two coins are thrown at the same time. Find the probability of									
getting both head			p. 0000, 0.						
(A) 3/4 (B) 1/4		(D) 0							
() -, .	(-) -, -	(-)							
13. Two dice are thrown simultaneously. The probability of getting a									

1

sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1/9	(D) 4/	9					
14. 100 cards are numbered from 1 to 100. Find the probability of getting a prime number.									
• • •	(B) 27/50	(C) 1/4	(D)	29/100					
of drawing a	15. A bag contains 5 red balls and some blue balls .If the probability of drawing a blue ball is double that of a red ball, then the number of blue balls in a bag is:								
(A) 5	(B) 10	(C) 15	(D) $20 = x$	= 10					
	random from	ontains 12 defenthis box. Ther	the probabi	lity that it is					
non-defective bulb is: (A) 143/150 (B) 147/150 (C) 1/25 (D) 1/50 =0.98OR 147/150 17. Cards marked with numbers 2 to 101 are placed in a box and mixed thoroughly. One card is drawn from this box randomly, then the probability that the number on card is a perfect square. (A) 9/100 (B) 1/10 (C) 3/10 (D) 19/100									
18. What is (A) 1/7	-	ity of getting 53 (C) 2/7	Mondays in (D) 7/						
		n a well shuffled ing of red suit.							
			\ 1 /1 0)38					
equally likel 1,2,312 ;	y to come to	onsists of spinn rest pointing to bability that it w (C) 7/12	one of the n	umber n odd number is:					
21. A game its outcome result i.e. th probability t	consists of t each time. A ree heads or	cossing a one ru Aryan wins if all three tails and ill lose the game	pee coin 3 tir the tosses g loses otherw	mes and noting ive the same					

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:									
-	(B) 31/365		(D) [*]	1/133225					
2. Then the pr	x is chosen at ra robability that x ² < 2/5 (C) 3/5	2 is?	numbers -2,	-1, 0 , 1,					
24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is: (A) 10 (B) 6 (C) 8 (D) 7									
Then the prob	is selected at rar ability that it is a 4/25 (C) 1/25	multiple of 3 and		umbers.					
with n dots sh	26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?								
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) $\frac{4}{21}$	1 + 2 + 3 + 4 + 5 + 6 = 21 =4/21					
	red by batsman ir ne standard devia	n 5 one day mato							
a) 25.79	b) 25.49		d) 25.69	9					
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.									
a) 13, 15	b) 13, 18	c) 18, 15		d) 13, 16					
3 cases is a) $^1/_2$ 30. X is a varia	b) $^1\!/_3$ ate between 0 and	c) $\frac{1}{4}$ d 3. The value of		d) $\frac{1}{16}$					
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?									

32.Out of the following values, which one is not possible in probability?									
a) $P(x) = 1$ c) $P(x) = 0.5$		b) $\sum x P(x) =$	x(x) = 3 x(x) = 0.5						
33.If E(x) =	2 and l b) 6	E(z) = 4, th	n en E(z - c) 0	- x) =?	d) Ins	ufficient d	ata		
34.The cova	ariance	of two inc	depende	ent rando	m variab	le is			
a) 1	b) 0		c) - 1		d) Un	defined			
35.If Σ P(x)	35.If $\Sigma P(x) = k^2 - 8$ then, the value of k is? a) 0 b) 1 c) 3 d) Insufficient data								
36.If P(x) = 0.5 and x = 4 , then E(x) = ? a) 1 b) 0.5 c) 4 d) 2									
37.In a disc	rete pro	obability d	istributi	on, the s	um of all	probabilit	ies		
is always? a) 0	b) Infinite c) 1 d) Undefined								
38.If the pr	obabili	ty of hittin	g the ta	rget is 0.	4, find m	ean and			
variance. a) 0.4, 0.24		b) 0.6, 0.2	4	c) 0.4	, 0.16	d) 0.6, 0).16		
39.If the probability that a bomb dropped from a place will strike the target is 60% and if 10 bombs are dropped, find mean and variance? a) 0.6, 0.24 b) 6, 2.4 c) 0.4, 0.16 d) 4, 1.6									
40. Find the mean of tossing 8 coins. a) 2 b) 4 c) 8 d) 1 41. What is the mean and variance for standard normal distribution?									

c) 5

d) 7

a) 3

b) 4

a) Mean is 0 and variance is 1 b) Mean is 1 and variance is 0 c) Mean is 0 and variance is ∞ d) Mean is ∞ and variance is 0									
42. Variance of a random variable X is given by a) $E(X)$ b) $E(X2)$ c) $E(X2)$ - $E(X)$ d) $E(X)$									
43.Mean o	a) E(X) b) E(X2) c) E(X2) - (E(X))2 d) (E(X))2								
44.Mean o a) 0	44.Mean of a constant 'a' is a) 0								
45. Variance of a constant 'a' is a) 0									
46.Find the mean and variance of X?									
x 0 1 2 3 4									
f(x)	1/9	2/9	3/9	2/9	1/9				
a) 2, 4/3 b) 3, 4/3 c) 2, 2/3 d) 3, 2/3									
47. Find the expectation of a random variable X?									

	X	0	1	2	3			
	f(x)	1/6	2/6	2/6	1/6			
a) ().5		b) 1.5			c) 2.5		d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by



c) np2q

d) npq2

49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**

a)
$$P(X = x) = nCx px qx$$

- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx

50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a) \sqrt{np}
- b) \sqrt{pq} c) (np)2
- d) \sqrt{npq}