



# Data Structures and Algorithms

## *1st Session*

Lecturer: Dr.Katanforoush

Author: Muhammad Karbalaee

October 14, 2022

## Contents

<b>1</b>	<b>Different sorts of problem statement in algorithm theory</b>	<b>2</b>
1.1	What is an algorithm? . . . . .	2
1.2	Importance of problem categorization	2
1.2.1	Satisfaction Problems . . . .	2
1.2.2	Root finding Problems . . . .	2
1.2.3	Enumeration Problems . . . .	2
1.2.4	Optimization Problems . . . .	2
<b>2</b>	<b>Exercises</b>	<b>3</b>
<b>3</b>	<b>Solutions</b>	<b>4</b>

# 1 Different sorts of problem statement in algorithm theory

## 1.1 What is an algorithm?

Algorithm is the method of solving an specific problem.

## 1.2 Importance of problem categorization

The definition of algorithm is mostly relied on what a problem is. Therefore, knowing the exact definition of problem is vital to understand what an algorithm is. In theory of algorithms, only specific statements are valid and every problem should get pigeonholed in one of the categories. There are four major problem sorts.

### 1.2.1 Satisfaction Problems

Problems in this category are stated in the following pattern.

*$P(X)$  is a proposition in the predicate logic <sup>1</sup> and the question is whether there is an  $X$  that  $P(X)=True$ .*

The algorithm which solves a satisfaction problem can only have two outputs, True or False. It's crucial to know that in satisfaction problems, the algorithm is not concerned with finding the wanted  $X$ , but is focused on validating the existence of  $X$ .

Example 1 *Is 2 a prime number?*

In this example  $P$  is being prime number and  $X=2$

This problem could be stated in other words too, which is still a satisfaction problem.

*Is  $P(X)=True$  by the given value  $X$ ?*

Example 2 *Consider graph  $G$  with  $V$  as its vertex set and  $E$  as its edge set. Is there a path between  $s$  and  $t$  where  $s, t \in V$*

### 1.2.2 Root finding Problems

Remember in the previous category the emphasis was on the existence of  $X$ ?

<sup>1</sup>Predicate logic in simple words means a logic system that contains variables in its statements and is concerned with the existence of that variable not the value of the variable itself for example: other than propositions such as "Socrates is a man", one can have expressions in the form "there exists  $x$  such that  $x$  is Socrates and  $x$  is a man", where "there exists" is a quantifier, while  $x$  is a variable.

In contrast with that, root finding problems are concerned with finding the  $X$  itself not checking whether it exists or not.

Formally stating:

*Find an  $X$ , if exists, such that  $P(X)=True$*

Example *Consider graph  $G$  with  $V$  as its vertex set and  $E$  as its edge set. Find path between  $s$  and  $t$  such as  $P = \langle v_1, v_2, v_3, \dots, v_k \rangle, k \in N$  where  $s, t \in V$*

### 1.2.3 Enumeration Problems

These problems aim to find all  $X$ s that  $P(X)=True$ , not just one. Each enumeration problem can be reduced to a root finding problem and root finding problems can be reduced to a satisfaction problem if the working space is countable. <sup>2</sup>

Example 1 *Find all two-digit prime numbers*

### 1.2.4 Optimization Problems

Consider  $f$  as a function defined this way,  $f : S \Rightarrow R$  and  $P(X)$  as a satisfiable proposition.

Optimization problems are formed this way then:  
*Find  $X$  such that  $f(X)$  reaches its minimum/maximum value and  $P(X)=True$*

$f$ : Object function

$P$ : constraints

Example  $G(V,E)$  is a graph and  $w$  is an edge weighing function such as  $w : E \Rightarrow R$ .

For given  $s$  and  $t$  such that  $s, t \in V$ , find the path with minimum total weight from  $s$  to  $t$ .

$S$ : The space of all paths in  $G(V,E)$

$X \in S, Path \in S$

$f(x) = \sum_{i=1}^k w(v_i, v_{i+1})$   
 $(v_i, v_{i+1}) \in Path$

$P : v_1 = s, v_k = t$

<sup>2</sup>Problem reduction means, by having an algorithm that solves problem A, it's possible to come up with an algorithm that solves problem B, this way it is said that problem B is reduced to problem A. For example: To find all two-digit prime numbers, we should be able to firstly, find a prime number. So the first problem is reduced to the second one.

## 2 Exercises

Here are a couple exercises related to the subject of the first session.

*Categorize each problem into one of the four categories you learnt in this session.*

1. Consider  $M$  as 3 by 5 matrix. Is  $M$  invertible?
2. Find an  $X$  such that  $2x + 10 = 0$
3. Find all divisors of 35.
4. All prime numbers are odd.
5. What is the fastest way to get from the dorm to the campus.
6. Consider  $V$  as a vector two-dimensional space. Find all vectors in  $V$  such as  $t$  that  $t = \langle 0, 0 \rangle$

### 3 Solutions

- 1.satisfaction problem
- 2.root finding problem
- 3.enumeration problem
- 4.satisfaction problem
- 5.Optimization problem
- 6.enumeration problem