Exponential functions

1

a Yes **b** No

2 **A**

3

a Decreasing **b** Increasing **c** Decreasing **d** Increasing

e Increasing **f** Increasing **g** Decreasing **h** Decreasing

i Decreasing **j** Increasing **k** Increasing **l** Increasing

4 **B**

5

a $y = 5^x$ **b** $y = 3 \times 2^x$

6

a 1 **b** -2 **c** 3 **d** 3

7

a -243 **b** -3 **c** $-\frac{1}{3}$ **d** 4

8 67

9

a 1 **b** $\frac{9}{16}$

10 $50\,313.28$

11 6885.29

12 3313.46

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

13

a

i $\frac{1}{16}$ ii 1 iii 16

b None. For all values of x, it will have a positive value.

c It gets larger, approaching infinity.

d It gets smaller, approaching zero.

14

a

x	-5	-4	-3	-2	-1	0	1	2	3	4	5	10
y	$\frac{1}{243}$	$\frac{1}{81}$	$\frac{1}{27}$	$\frac{1}{9}$	$\frac{1}{3}$	1	3	9	27	81	243	$59\,049$

b As x increases, the function increases at a faster and faster rate.

c All real x

d $y > 0$

15

a

x	-5	-4	-3	-2	-1	0	1	2	3	4	5	10
y	32	16	8	4	2	1	$\frac{1}{2}$	$\frac{1}{4}$	$\frac{1}{8}$	$\frac{1}{16}$	$\frac{1}{32}$	$\frac{1}{1024}$

b As x increases, the function decreases at a slower and slower rate.

c 1

d All real x

e $y > 0$

16

a

x	-5	-4	-3	-2	-1	0	1	2	3	4	5	10
y	$-\frac{1}{32}$	$-\frac{1}{16}$	$-\frac{1}{8}$	$-\frac{1}{4}$	$-\frac{1}{2}$	-1	-2	-4	-8	-16	-32	-1024

b No. For $a > 0$ the expression a^x is greater than zero for all real x. So, $-(a^x)$ will be less than or equal to zero.

c Decreasing

d As x increases, the function decreases at a faster and faster rate.

e All real x.

f $y < 0$

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

17

a Positive

b 0

c $y = 0$ is a horizontal asymptote of the curve.

18

a $(0, 1)$ **b** No **c** All real x **d** $y > 0$


e $y = 128$ **f** $x = 8$

19

a

x	-3	-2	-1	0	1	2	3
$f(x)$	0.05	0.14	0.37	1.00	2.72	7.39	20.09

b



20


a

x	-2	-1	0	1	2	3
y	$-\frac{1}{100}$	$-\frac{1}{10}$	-1	-10	-100	-1000

b Yes. For any value of x, 10^x is always positive. Therefore, -10^x is always negative.

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

c



21

a No. For $a > 0$ the expression a^x is greater than zero for all real x.


b 0

c ∞

d 1

e 0

f



© ClipSAT: By Mr. Mohamed Abdallah +966597688647


22

a $(0, 4)$

b No

c ∞

d



23

a 2


b No. For $a > 0$ the expression a^x is greater than zero for all real x. So, $2 \times (a^x)$ will be greater than or equal to zero.

c All real x.

d $y > 0$

e ∞

f



24

a Graph R

b For $x < 0$, the graph of $y = 6^x$ is below the graph of $y = 4^x$. This is because negative values of x result in fractional function values, and for any negative value of x, 6^x will result in a smaller value than 4^x.

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

25 a For $x > 0$, the graph of $y = e^x$ will lie above the graph of $y = (2)^x$ and below the graph of $y = (3)^x$.


b For $x < 0$, the graph of $y = e^x$ will lie above the graph of $y = (3)^x$ and below the graph of $y = (2)^x$.

Transformations of exponential functions

26

a A reflection over the x-axis


b



27

a Vertical translation

b




© ClipSAT: By Mr. Mohamed Abdallah +966597688647

28

a Reflection over the y-axis.


b



29

a Reflection over the x-axis and y-axis.

b



30

a $y = 0$ **b** $y = 2$ **c** 1 **d** All real x.

e $y < 2$

31

a 1 **b** -1 **c** $y = 0$ **d** $y = -2$

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

32

a

i 4 units to the left

ii 2 units

iii 1

b

i 1 units to the left

ii 1 units down

iii $\frac{1}{3}$

c

i $\frac{1}{5}$ units to the right

ii 2 units

iii 5

33

a $y = 2^x - 7$

b $y = 11^{x+10}$

34


a $y = 7^{x-3} - 5$

b $y = -7^{x+3} + 5$


c $y = -7^{x+3} - 5$

35

a



b



© ClipSAT: By Mr. Mohamed Abdallah +966597688647

36


a

i 4

ii All real x

iii $y > 3$

iv




b

i -4

ii All real x

iii $y > -5$

iv



© ClipSAT: By Mr. Mohamed Abdallah +966597688647


c

i 16

ii All real x

iii $y > 0$

iv




d

i $\frac{1}{3}$

ii All real x

iii $y > 0$

iv



© ClipSAT: By Mr. Mohamed Abdallah +966597688647


e

i 2

ii All real x

iii $y < 3$

iv




37

a $y = 0$ **b** $y = 2$ **c** 1

38

a


i $(0, 3)$ ii $y = 2$ iii



© ClipSAT: By Mr. Mohamed Abdallah +966597688647


b

i $(0, -1)$ ii $y = -2$ iii




c

i $(0, 1)$ ii $y = 2$ iii



d


i $(0, 0)$ ii $y = -1$ iii



© ClipSAT: By Mr. Mohamed Abdallah +966597688647

e

i $(0, 2)$ ii $y = 1$ iii




39

a

Point on original graph	Point on new graph
$\left(-1, \frac{1}{3}\right)$	$\left(-1, \frac{2}{3}\right)$
$(0, 1)$	$(0, 2)$
$(1, 3)$	$(1, 6)$
$(2, 9)$	$(2, 18)$

b $y = 2 \times 3^x$

c



d For all values of x, $y = 2 \times 3^x$ lies above $y = 3^x$.

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

40

a
i No ii No iii No

b
i No ii Yes iii Yes

c
i Yes ii Yes iii Yes

d
i No ii Yes iii Yes

Exponential equations

41

a $y = 2^x$ **b** $y = 2^{-x}$ **c** $y = 9^x$ **d** $a = 5$

42

a

x	1	2	3	4
y	2	4	8	16

b 2

c $y = 2^x$

43

a 10 **b** $y = 10^x$ **c** $10\,000\,000\,000$

44

a $a = 5$ **b** $b = 4$ **c** $y = \frac{5}{16}$

© ClipSAT: By Mr. Mohamed Abdallah +966597688647