Properties of logarithms

1 Simplify each of the following expressions:

a $\log 4 + \log 9$

b $\log_{10}(10) + \log_{10}(10)$

c $\log_{10} 11 + \log_{10} 2 + \log_{10} 9$

d $\log_{10} 12 - (\log_{10} 2 + \log_{10} 3)$

e $\dfrac{\log_{10} 4}{\log_{10} 2}$

f $\dfrac{\log_4 125}{\log_4 5}$

g $\dfrac{\log a^8}{\log a^4}$

h $\dfrac{\log a^3}{\log \sqrt[3]{a}}$

i $\log 10x + \log 10y$

j $\log x^2 + \log x^3$

k $\dfrac{\log\left(\dfrac{1}{x^4}\right)}{\log x}$

l $\log_{10} 10 + \dfrac{\log_{10}(15^{20})}{\log_{10}(15^5)}$

m $\dfrac{5\log m^2}{6\log \sqrt[3]{m}}$

n $\dfrac{8\log_{10}(\sqrt{10})}{\log_{10}(100)}$

o $10^{\log w}$

p $x^{4\log_x 3 - 6\log_x 2}$

2 Simplify each of the following expressions:

a $\dfrac{\ln a^6}{\ln a^3}$

b $\dfrac{12\ln(m^4)}{10\ln\sqrt{m}}$

c $\dfrac{\ln(a^4)}{\ln \sqrt[3]{a}}$

d $\dfrac{2\ln x}{10\ln y}$

3 Rewrite the following as the sum or difference of logarithms without any powers, fractions or radicals:

a $\log_9 uv$

b $\log(x^4)$

c $\log_4(x^7)$

d $\log_5\left(\dfrac{9}{7}\right)$

e $\log\left(x^{\frac{2}{5}}\right)$

f $\log_b(x^2)$

g $\log(3x^{-1})$

h $\log\left((3x)^5\right)$

i $\log(7x^{-4})$

j $\log\left((5x)^{-7}\right)$

k $\log\left((2x)^{-1}\right)$

l $\log\left(\dfrac{1}{xy}\right)$

m $\log\left(\dfrac{pq}{r}\right)$

n $\log\left((3x+7)^{-1}\right)$

o $\log\left((3x+4)^{-8}\right)$

p $\log\left((5x+7)^{\frac{1}{2}}\right)$

q $\log\left((5x+2)^6\right)$

r $\log\left((x+6)^5\right)$

s $\log_2(5x)$

t $\log_{10}\left(\dfrac{2}{x}\right)$

u $\log(m^2)$

v $\log\left(5x^{\frac{2}{3}}\right)$

w $\log\left((14x)^{\frac{1}{3}}\right)$

x $\log\left(\sqrt{\dfrac{c^8}{d}}\right)$

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

4 Rewrite the following as the sum or difference of logarithms without any powers, fractions or radicals:

a $\ln\left(\frac{15}{8}\right)$ **b** $\ln\left(\frac{pq}{r}\right)$ **c** $\ln\left(\frac{19}{7}\right)$

5 Rewrite each of the following as a single logarithm or integer:

a $5\log x^3 - 4\log x^2$ **b** $5\log x + 3\log y$

c $8\log x - \frac{1}{3}\log y$ **d** $7\log x - \log\left(\frac{1}{x}\right) - \log y$

e $7\log_{10} 5 - 21\log_{10} 25$ **f** $5\log_{10} 8 - 3\log_{10} 4$

g $2\log_6 3 + \frac{1}{3}\log_6 64$ **h** $\log_2 36 - 2\log_2 3$

6 Rewrite each of the following expressions as a single logarithm:

a $\ln 3 + \ln 5$ **b** $\ln 24 - \ln 4$

c $\ln 5 - \ln 7$ **d** $\ln 11 + \ln 5 + \ln 8$

e $\ln 24 - (\ln 2 + \ln 6)$ **f** $\ln 7 + \ln 2 + \ln 3 - \ln 5$

7 Rewrite each of the following expressions as a single logarithm:

a $6\ln x - \frac{1}{6}\ln y$ **b** $\ln(3x) + \ln(5y)$

c $\ln\left(x^2\right) + \ln\left(x^3\right)$ **d** $3\ln\left(x^5\right) - 4\ln\left(x^2\right)$

8 Use the properties of logarithms to express each of the following without any powers or radicals:

a $\ln y^2$ **b** $\ln\frac{1}{y^2}$ **c** $\ln\sqrt{y}$ **d** $\ln\sqrt[3]{y}$

e $\ln y^{-5}$ **f** $\ln y^{\frac{2}{5}}$ **g** $\ln y^{\frac{1}{2}}$ **h** $\ln\sqrt{w}$

9 Use the properties of logarithms to rewrite each of the following in the form $\ln b^a$ where $a > 0$:

a $-2\ln x$ **b** $\frac{\ln x}{2}$

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

Change of base

10 Rewrite the following in terms of base 10 logarithms:

a $\log_4 16$ **b** $\log_3 0.9$ **c** $\log_3 \sqrt{5}$ **d** $\log_a B$

11 Rewrite $\log_3 20$ in terms of base 4 logarithms.

12 Consider the following logarithmic expressions:

i Rewrite the expression in terms of base 10 logarithms.

ii Now, evaluate each to two decimal places.

a $\log_8 21$ **b** $\log_4 \sqrt{5}$ **c** $\log_2 3^3$ **d** $\log_\pi 105$

© ClipSAT: By Mr. Mohamed Abdallah +966597688647