Inverse relationship between exponentials and logarithms

ANSWER KEY

Core Concepts: Inverse relationship between exponentials and logarithms

1 For the model $f(x) = 5x + 4$ in inverse relationship between exponentials and logarithms, evaluate $f(2)$.

a. 14 ✓

b. 19

c. 10

d. 18

Substitute $x = 2$: $f(2) = 5(2) + 4 = 14$.

2 Find the positive solutions to the quadratic equation $x^2 - 7x + 6 = 0$.

a. $x = 1$ and $x = 7$

b. $x = 1$ **and** $x = 6$ ✓

c. $x = -1$ and $x = -6$

d. $x = 7$ and $x = 6$

Factoring $(x - 1)(x - 6) = 0$ yields the roots $x = 1$ and $x = 6$.

3 A data set recorded for inverse relationship between exponentials and logarithms consists of the four values $\{17, 21, 25, 31\}$. What is the arithmetic mean?

a. 21.5

b. 22.5

c. 23.5 ✓

d. 25.5

The sum is $17 + 21 + 25 + 31 = 94$, and dividing by 4 gives 23.5.

Procedural Fluency & Algebraic Manipulation

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

4 For the model $f(x) = 5x + 4$ in inverse relationship between exponentials and logarithms, evaluate $f(3)$.

a. 19 ✓

b. 24

c. 15

d. 23

Substitute $x = 3$: $f(3) = 5(3) + 4 = 19$.

5 Find the positive solutions to the quadratic equation $x^2 - 9x + 14 = 0$.

a. $x = 2$ and $x = 8$

b. $x = 2$ **and** $x = 7$ ✓

c. $x = -2$ and $x = -7$

d. $x = 9$ and $x = 14$

Factoring $(x - 2)(x - 7) = 0$ yields the roots $x = 2$ and $x = 7$.

6 A data set recorded for inverse relationship between exponentials and logarithms consists of the four values $\{17, 21, 25, 31\}$. What is the arithmetic mean?

a. 21.5

b. 22.5

c. 23.5 ✓

d. 25.5

The sum is $17 + 21 + 25 + 31 = 94$, and dividing by 4 gives 23.5.

Graphical & Geometric Interpretation

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

7 For the model $f(x) = 5x + 4$ in inverse relationship between exponentials and logarithms, evaluate $f(4)$.

a. 24 ✓

b. 29

c. 20

d. 28

Substitute $x = 4$: $f(4) = 5(4) + 4 = 24$.

8 Find the positive solutions to the quadratic equation $x^2 - 11x + 24 = 0$.

a. $x = 3$ and $x = 9$

b. $x = 3$ **and** $x = 8$ ✓

c. $x = -3$ and $x = -8$

d. $x = 11$ and $x = 24$

Factoring $(x - 3)(x - 8) = 0$ yields the roots $x = 3$ and $x = 8$.

9 A data set recorded for inverse relationship between exponentials and logarithms consists of the four values $\{17, 21, 25, 31\}$. What is the arithmetic mean?

a. 21.5

b. 22.5

c. 23.5 ✓

d. 25.5

The sum is $17 + 21 + 25 + 31 = 94$, and dividing by 4 gives 23.5.

Exam-Style Problem Solving & Modeling

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

10 For the model $f(x) = 5x + 4$ in inverse relationship between exponentials and logarithms, evaluate $f(5)$.

a. 29 ✓

b. 34

c. 25

d. 33

Substitute $x = 5$: $f(5) = 5(5) + 4 = 29$.

11 Find the positive solutions to the quadratic equation $x^2 - 13x + 36 = 0$.

a. $x = 4$ and $x = 10$

b. $x = 4$ **and** $x = 9$ ✓

c. $x = -4$ and $x = -9$

d. $x = 13$ and $x = 36$

Factoring $(x - 4)(x - 9) = 0$ yields the roots $x = 4$ and $x = 9$.

12 A data set recorded for inverse relationship between exponentials and logarithms consists of the four values $\{17, 21, 25, 31\}$. What is the arithmetic mean?

a. 21.5

b. 22.5

c. 23.5 ✓

d. 25.5

The sum is $17 + 21 + 25 + 31 = 94$, and dividing by 4 gives 23.5.

© ClipSAT: By Mr. Mohamed Abdallah +966597688647