# Arithmetics sequences

**ANSWER KEY**

## Core Concepts: Arithmetics sequences

**1** For the model $f(x) = 6x + 3$ in arithmetics sequences, evaluate $f(2)$.

**a.** $15$ ✓

**b.** $21$

**c.** $12$

**d.** $19$

Substitute $x = 2$: $f(2) = 6(2) + 3 = 15$.

**2** Find the positive solutions to the quadratic equation $x^2 - 5x + 4 = 0$.

**a.** $x = 1$ and $x = 5$

**b.** $x = 1$ **and** $x = 4$ ✓

**c.** $x = -1$ and $x = -4$

**d.** $x = 5$ and $x = 4$

Factoring $(x - 1)(x - 4) = 0$ yields the roots $x = 1$ and $x = 4$.

**3** A data set recorded for arithmetics sequences consists of the four values $\{14, 18, 22, 28\}$. What is the arithmetic mean?

**a.** $18.5$

**b.** $19.5$

**c.** $20.5$ ✓

**d.** $22.5$

The sum is $14 + 18 + 22 + 28 = 82$, and dividing by $4$ gives $20.5$.

---

## Procedural Fluency & Algebraic Manipulation

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

**4** For the model $f(x) = 6x + 3$ in arithmetics sequences, evaluate $f(3)$.

**a.** $21$ ✓

**b.** $27$

**c.** $18$

**d.** $25$

Substitute $x = 3$: $f(3) = 6(3) + 3 = 21$.

**5** Find the positive solutions to the quadratic equation $x^2 - 7x + 10 = 0$.

**a.** $x = 2$ and $x = 6$

**b.** $x = 2$ **and** $x = 5$ ✓

**c.** $x = -2$ and $x = -5$

**d.** $x = 7$ and $x = 10$

Factoring $(x - 2)(x - 5) = 0$ yields the roots $x = 2$ and $x = 5$.

**6** A data set recorded for arithmetics sequences consists of the four values $\{14, 18, 22, 28\}$. What is the arithmetic mean?

**a.** $18.5$

**b.** $19.5$

**c.** $20.5$ ✓

**d.** $22.5$

The sum is $14 + 18 + 22 + 28 = 82$, and dividing by $4$ gives $20.5$.

---

## Graphical & Geometric Interpretation

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

**7** For the model $f(x) = 6x + 3$ in arithmetics sequences, evaluate $f(4)$.

**a.** $27$ ✓

**b.** $33$

**c.** $24$

**d.** $31$

Substitute $x = 4$: $f(4) = 6(4) + 3 = 27$.

**8** Find the positive solutions to the quadratic equation $x^2 - 9x + 18 = 0$.

**a.** $x = 3$ and $x = 7$

**b.** $x = 3$ **and** $x = 6$ ✓

**c.** $x = -3$ and $x = -6$

**d.** $x = 9$ and $x = 18$

Factoring $(x - 3)(x - 6) = 0$ yields the roots $x = 3$ and $x = 6$.

**9** A data set recorded for arithmetics sequences consists of the four values $\{14, 18, 22, 28\}$. What is the arithmetic mean?

**a.** $18.5$

**b.** $19.5$

**c.** $20.5$ ✓

**d.** $22.5$

The sum is $14 + 18 + 22 + 28 = 82$, and dividing by $4$ gives $20.5$.

---

## Exam-Style Problem Solving & Modeling

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

**10** For the model $f(x) = 6x + 3$ in arithmetics sequences, evaluate $f(5)$.

**a.** $33$ ✓

**b.** $39$

**c.** $30$

**d.** $37$

Substitute $x = 5$: $f(5) = 6(5) + 3 = 33$.

**11** Find the positive solutions to the quadratic equation $x^2 - 11x + 28 = 0$.

**a.** $x = 4$ and $x = 8$

**b.** $x = 4$ **and** $x = 7$ ✓

**c.** $x = -4$ and $x = -7$

**d.** $x = 11$ and $x = 28$

Factoring $(x - 4)(x - 7) = 0$ yields the roots $x = 4$ and $x = 7$.

**12** A data set recorded for arithmetics sequences consists of the four values $\{14, 18, 22, 28\}$. What is the arithmetic mean?

**a.** $18.5$

**b.** $19.5$

**c.** $20.5$ ✓

**d.** $22.5$

The sum is $14 + 18 + 22 + 28 = 82$, and dividing by $4$ gives $20.5$.

© ClipSAT: By Mr. Mohamed Abdallah +966597688647