Patterns with rational and irrational numbers

ANSWER KEY

Core Concepts: Patterns with rational and irrational numbers

1 For the model $f(x) = 6x + 5$ in patterns with rational and irrational numbers, evaluate $f(2)$.

a. 17 ✓

b. 23

c. 12

d. 21

Substitute $x = 2$: $f(2) = 6(2) + 5 = 17$.

2 Find the positive solutions to the quadratic equation $x^2 - 4x + 3 = 0$.

a. $x = 1$ and $x = 4$

b. $x = 1$ **and** $x = 3$ ✓

c. $x = -1$ and $x = -3$

d. $x = 4$ and $x = 3$

Factoring $(x - 1)(x - 3) = 0$ yields the roots $x = 1$ and $x = 3$.

3 A data set recorded for patterns with rational and irrational numbers consists of the four values $\{18, 22, 26, 32\}$. What is the arithmetic mean?

a. 22.5

b. 23.5

c. 24.5 ✓

d. 26.5

The sum is $18 + 22 + 26 + 32 = 98$, and dividing by 4 gives 24.5.

Procedural Fluency & Algebraic Manipulation

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

4 For the model $f(x) = 6x + 5$ in patterns with rational and irrational numbers, evaluate $f(3)$.

a. 23 ✓

b. 29

c. 18

d. 27

Substitute $x = 3$: $f(3) = 6(3) + 5 = 23$.

5 Find the positive solutions to the quadratic equation $x^2 - 6x + 8 = 0$.

a. $x = 2$ and $x = 5$

b. $x = 2$ **and** $x = 4$ ✓

c. $x = -2$ and $x = -4$

d. $x = 6$ and $x = 8$

Factoring $(x - 2)(x - 4) = 0$ yields the roots $x = 2$ and $x = 4$.

6 A data set recorded for patterns with rational and irrational numbers consists of the four values $\{18, 22, 26, 32\}$. What is the arithmetic mean?

a. 22.5

b. 23.5

c. 24.5 ✓

d. 26.5

The sum is $18 + 22 + 26 + 32 = 98$, and dividing by 4 gives 24.5.

Graphical & Geometric Interpretation

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

7 For the model $f(x) = 6x + 5$ in patterns with rational and irrational numbers, evaluate $f(4)$.

a. 29 ✓

b. 35

c. 24

d. 33

Substitute $x = 4$: $f(4) = 6(4) + 5 = 29$.

8 Find the positive solutions to the quadratic equation $x^2 - 8x + 15 = 0$.

a. $x = 3$ and $x = 6$

b. $x = 3$ **and** $x = 5$ ✓

c. $x = -3$ and $x = -5$

d. $x = 8$ and $x = 15$

Factoring $(x - 3)(x - 5) = 0$ yields the roots $x = 3$ and $x = 5$.

9 A data set recorded for patterns with rational and irrational numbers consists of the four values $\{18, 22, 26, 32\}$. What is the arithmetic mean?

a. 22.5

b. 23.5

c. 24.5 ✓

d. 26.5

The sum is $18 + 22 + 26 + 32 = 98$, and dividing by 4 gives 24.5.

Exam-Style Problem Solving & Modeling

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

10 For the model $f(x) = 6x + 5$ in patterns with rational and irrational numbers, evaluate $f(5)$.

a. 35 ✓

b. 41

c. 30

d. 39

Substitute $x = 5$: $f(5) = 6(5) + 5 = 35$.

11 Find the positive solutions to the quadratic equation $x^2 - 10x + 24 = 0$.

a. $x = 4$ and $x = 7$

b. $x = 4$ **and** $x = 6$ ✓

c. $x = -4$ and $x = -6$

d. $x = 10$ and $x = 24$

Factoring $(x - 4)(x - 6) = 0$ yields the roots $x = 4$ and $x = 6$.

12 A data set recorded for patterns with rational and irrational numbers consists of the four values $\{18, 22, 26, 32\}$. What is the arithmetic mean?

a. 22.5

b. 23.5

c. 24.5 ✓

d. 26.5

The sum is $18 + 22 + 26 + 32 = 98$, and dividing by 4 gives 24.5.

© ClipSAT: By Mr. Mohamed Abdallah +966597688647