Regions defined by inequalities (Extended)

ANSWER KEY

Core Concepts: Regions defined by inequalities (Extended)

1 For the model $f(x) = 2x + 7$ in regions defined by inequalities (extended), evaluate $f(2)$.

a. 11 ✓

b. 13

c. 4

d. 15

Substitute $x = 2$: $f(2) = 2(2) + 7 = 11$.

2 Find the positive solutions to the quadratic equation $x^2 - 4x + 3 = 0$.

a. $x = 1$ and $x = 4$

b. $x = 1$ **and** $x = 3$ ✓

c. $x = -1$ and $x = -3$

d. $x = 4$ and $x = 3$

Factoring $(x - 1)(x - 3) = 0$ yields the roots $x = 1$ and $x = 3$.

3 A data set recorded for regions defined by inequalities (extended) consists of the four values $\{12, 16, 20, 26\}$. What is the arithmetic mean?

a. 16.5

b. 17.5

c. 18.5 ✓

d. 20.5

The sum is $12 + 16 + 20 + 26 = 74$, and dividing by 4 gives 18.5.

Procedural Fluency & Algebraic Manipulation

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

4 For the model $f(x) = 2x + 7$ in regions defined by inequalities (extended), evaluate $f(3)$.

a. 13 ✓

b. 15

c. 6

d. 17

Substitute $x = 3$: $f(3) = 2(3) + 7 = 13$.

5 Find the positive solutions to the quadratic equation $x^2 - 6x + 8 = 0$.

a. $x = 2$ and $x = 5$

b. $x = 2$ **and** $x = 4$ ✓

c. $x = -2$ and $x = -4$

d. $x = 6$ and $x = 8$

Factoring $(x - 2)(x - 4) = 0$ yields the roots $x = 2$ and $x = 4$.

6 A data set recorded for regions defined by inequalities (extended) consists of the four values $\{12, 16, 20, 26\}$. What is the arithmetic mean?

a. 16.5

b. 17.5

c. 18.5 ✓

d. 20.5

The sum is $12 + 16 + 20 + 26 = 74$, and dividing by 4 gives 18.5.

Graphical & Geometric Interpretation

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

7 For the model $f(x) = 2x + 7$ in regions defined by inequalities (extended), evaluate $f(4)$.

a. 15 ✓

b. 17

c. 8

d. 19

Substitute $x = 4$: $f(4) = 2(4) + 7 = 15$.

8 Find the positive solutions to the quadratic equation $x^2 - 8x + 15 = 0$.

a. $x = 3$ and $x = 6$

b. $x = 3$ **and** $x = 5$ ✓

c. $x = -3$ and $x = -5$

d. $x = 8$ and $x = 15$

Factoring $(x - 3)(x - 5) = 0$ yields the roots $x = 3$ and $x = 5$.

9 A data set recorded for regions defined by inequalities (extended) consists of the four values $\{12, 16, 20, 26\}$. What is the arithmetic mean?

a. 16.5

b. 17.5

c. 18.5 ✓

d. 20.5

The sum is $12 + 16 + 20 + 26 = 74$, and dividing by 4 gives 18.5.

Exam-Style Problem Solving & Modeling

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

10 For the model $f(x) = 2x + 7$ in regions defined by inequalities (extended), evaluate $f(5)$.

a. 17 ✓

b. 19

c. 10

d. 21

Substitute $x = 5$: $f(5) = 2(5) + 7 = 17$.

11 Find the positive solutions to the quadratic equation $x^2 - 10x + 24 = 0$.

a. $x = 4$ and $x = 7$

b. $x = 4$ **and** $x = 6$ ✓

c. $x = -4$ and $x = -6$

d. $x = 10$ and $x = 24$

Factoring $(x - 4)(x - 6) = 0$ yields the roots $x = 4$ and $x = 6$.

12 A data set recorded for regions defined by inequalities (extended) consists of the four values $\{12, 16, 20, 26\}$. What is the arithmetic mean?

a. 16.5

b. 17.5

c. 18.5 ✓

d. 20.5

The sum is $12 + 16 + 20 + 26 = 74$, and dividing by 4 gives 18.5.

© ClipSAT: By Mr. Mohamed Abdallah +966597688647