Solving contextual problems with integers

ANSWER KEY

Core Concepts: Solving contextual problems with integers

1 For the model $f(x) = 4x + 5$ in solving contextual problems with integers, evaluate $f(2)$.

a. 13 ✓

b. 17

c. 8

d. 17

Substitute $x = 2$: $f(2) = 4(2) + 5 = 13$.

2 Find the positive solutions to the quadratic equation $x^2 - 5x + 4 = 0$.

a. $x = 1$ and $x = 5$

b. $x = 1$ **and** $x = 4$ ✓

c. $x = -1$ and $x = -4$

d. $x = 5$ and $x = 4$

Factoring $(x - 1)(x - 4) = 0$ yields the roots $x = 1$ and $x = 4$.

3 A data set recorded for solving contextual problems with integers consists of the four values $\{11, 15, 19, 25\}$. What is the arithmetic mean?

a. 15.5

b. 16.5

c. 17.5 ✓

d. 19.5

The sum is $11 + 15 + 19 + 25 = 70$, and dividing by 4 gives 17.5.

Procedural Fluency & Algebraic Manipulation

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

4 For the model $f(x) = 4x + 5$ in solving contextual problems with integers, evaluate $f(3)$.

a. 17 ✓

b. 21

c. 12

d. 21

Substitute $x = 3$: $f(3) = 4(3) + 5 = 17$.

5 Find the positive solutions to the quadratic equation $x^2 - 7x + 10 = 0$.

a. $x = 2$ and $x = 6$

b. $x = 2$ **and** $x = 5$ ✓

c. $x = -2$ and $x = -5$

d. $x = 7$ and $x = 10$

Factoring $(x - 2)(x - 5) = 0$ yields the roots $x = 2$ and $x = 5$.

6 A data set recorded for solving contextual problems with integers consists of the four values $\{11, 15, 19, 25\}$. What is the arithmetic mean?

a. 15.5

b. 16.5

c. 17.5 ✓

d. 19.5

The sum is $11 + 15 + 19 + 25 = 70$, and dividing by 4 gives 17.5.

Graphical & Geometric Interpretation

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

7 For the model $f(x) = 4x + 5$ in solving contextual problems with integers, evaluate $f(4)$.

a. 21 ✓

b. 25

c. 16

d. 25

Substitute $x = 4$: $f(4) = 4(4) + 5 = 21$.

8 Find the positive solutions to the quadratic equation $x^2 - 9x + 18 = 0$.

a. $x = 3$ and $x = 7$

b. $x = 3$ **and** $x = 6$ ✓

c. $x = -3$ and $x = -6$

d. $x = 9$ and $x = 18$

Factoring $(x - 3)(x - 6) = 0$ yields the roots $x = 3$ and $x = 6$.

9 A data set recorded for solving contextual problems with integers consists of the four values $\{11, 15, 19, 25\}$. What is the arithmetic mean?

a. 15.5

b. 16.5

c. 17.5 ✓

d. 19.5

The sum is $11 + 15 + 19 + 25 = 70$, and dividing by 4 gives 17.5.

Exam-Style Problem Solving & Modeling

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

10 For the model $f(x) = 4x + 5$ in solving contextual problems with integers, evaluate $f(5)$.

a. 25 ✓

b. 29

c. 20

d. 29

Substitute $x = 5$: $f(5) = 4(5) + 5 = 25$.

11 Find the positive solutions to the quadratic equation $x^2 - 11x + 28 = 0$.

a. $x = 4$ and $x = 8$

b. $x = 4$ **and** $x = 7$ ✓

c. $x = -4$ and $x = -7$

d. $x = 11$ and $x = 28$

Factoring $(x - 4)(x - 7) = 0$ yields the roots $x = 4$ and $x = 7$.

12 A data set recorded for solving contextual problems with integers consists of the four values $\{11, 15, 19, 25\}$. What is the arithmetic mean?

a. 15.5

b. 16.5

c. 17.5 ✓

d. 19.5

The sum is $11 + 15 + 19 + 25 = 70$, and dividing by 4 gives 17.5.

© ClipSAT: By Mr. Mohamed Abdallah +966597688647