
Trigonometry

ClipSAT Printable Practice Suite (clipsat.org) • Topic 6.01

Part I — Free-Response Questions (FRQs): Foundational Fluency (1–7)

1 [Foundational] • **Target: Right triangle trigonometric ratios**


In a right triangle, angle θ has an opposite side of 8 and hypotenuse 17. Find $\sin\theta$, $\cos\theta$, and $\tan\theta$.



© ClipSAT — clipsat.org | All Rights Reserved

2 [Foundational] • Target: Right triangle trigonometric ratios


A ladder 15 feet long leans against a wall, making a 70° angle with the ground. Find the height reached on the wall, to the nearest tenth.



© ClipSAT — clipsat.org | All Rights Reserved


3 [Foundational] • Target: The unit circle and radian measure

Convert 150° to radians, in terms of π.



4 [Foundational] • Target: The unit circle and radian measure


Find the exact value of $\sin\left(\frac{\pi}{6}\right)$ and $\cos\left(\frac{\pi}{4}\right)$, using the unit circle.



© ClipSAT — clipsat.org | All Rights Reserved


5 [Foundational] • Target: Graphs of sine and cosine functions

State the amplitude and period of $y = 3\sin(2x)$.




6 [Foundational] • Target: Graphs of sine and cosine functions

State the amplitude, period, and vertical shift of $y = 4\cos(x) + 2$.



7 [Foundational] • Target: Law of Sines

In triangle ABC, $A = 40^\circ$, $B = 65^\circ$, and $a = 10$. Find side b, to the nearest tenth, using the Law of Sines.




© ClipSAT — clipsat.org | All Rights Reserved

Part II — Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)


8 **[Intermediate]** • **Target: Law of Cosines**

In triangle ABC, $a = 7$, $b = 9$, and $C = 60^\circ$. Find side c, using the Law of Cosines.



9 **[Intermediate]** • **Target: Applications of trigonometry to real-world problems**


This question has two parts. A surveyor stands 200 feet from the base of a tower and measures the angle of elevation to the top as 38°. (a) Find the height of the tower, to the nearest foot. (b) If the surveyor moves 50 feet closer to the tower, find the new angle of elevation, to the nearest degree, using the height found in part (a).



© ClipSAT — clipsat.org | All Rights Reserved


10 [Intermediate] • Target: Reciprocal trigonometric functions

If $\sin\theta = \frac{5}{13}$, find $\csc\theta$.



11 [Intermediate] • Target: Reciprocal trigonometric functions


If $\cos\theta = \frac{3}{5}$, find $\sec\theta$ and $\tan\theta$ (assuming θ is in the first quadrant).



© ClipSAT — clipsat.org | All Rights Reserved


12 [Intermediate] • Target: Angle of elevation and depression

From the top of a 100-foot cliff, the angle of depression to a boat is 25°. Find the horizontal distance from the base of the cliff to the boat, to the nearest foot.



13 [Intermediate] • Target: A multi-part triangle problem combining Law of Sines and Law of Cosines


This question has two parts. Triangle ABC has $a = 8$, $b = 11$, and $C = 50^\circ$. (a) Use the Law of Cosines to find side c, to the nearest tenth. (b) Use the Law of Sines to find angle A, to the nearest degree.



© ClipSAT — clipsat.org | All Rights Reserved

14 **[Intermediate]** • **Target: Free-Response Questions (FRQs): Foundational Fluency (1–7)**


Find the exact value of $\sin\left(\frac{\pi}{6}\right)$ and $\cos\left(\frac{\pi}{4}\right)$, using the unit circle.



Part III — Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20)

15 **[Challenging]** • **Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)** • **Cross-Lesson: 7.01 Statistics & Probability**


If $\sin\theta = \frac{5}{13}$, find $\csc\theta$.



© ClipSAT — clipsat.org | All Rights Reserved

16 [Challenging] • Target: Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20) • Cross-Lesson: 8.01 Matrices, Vectors & Complex Numbers

If $\cos\theta = \frac{3}{5}$, find $\sec\theta$ and $\tan\theta$ (assuming θ is in the first quadrant).



17 [Challenging] • Target: The unit circle and radian measure • Cross-Lesson: 1.01 Numbers, Percents & Ratios

Convert $\frac{3\pi}{4}$ radians to degrees.

18 [Challenging] • Target: Basic trigonometric identities • Cross-Lesson: 2.01 Elementary & Intermediate Algebra

Use the Pythagorean identity to find $\cos\theta$, given that $\sin\theta = \frac{3}{5}$ and θ is in the first quadrant.

19 [Challenging] • Target: Basic trigonometric identities • Cross-Lesson: 3.01 Functions & Graphs

Simplify $\frac{\sin\theta}{\cos\theta} \cdot \cos\theta$.

20 [Challenging] • Target: Trigonometric equations • Cross-Lesson: 4.01 Coordinate Geometry

Solve $2\sin\theta = 1$ for $0^\circ \le \theta < 360^\circ$.

Part IV — Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

© ClipSAT — clipsat.org | All Rights Reserved

21 [Foundational] • Target: Right triangle trigonometric ratios

In a right triangle, the angle θ has an opposite side of length 20, adjacent side 21, and hypotenuse 29. Find $\tan\theta$.

a. $\frac{29}{21}$

b. $\frac{20}{21}$

c. $\frac{29}{20}$

d. $\frac{21}{20}$

22 [Foundational] • Target: The unit circle and radian measure

In a right triangle, the angle θ has an opposite side of length 9, adjacent side 12, and hypotenuse 15. Find $\sin\theta$.

a. $\frac{4}{3}$

b. $\frac{3}{5}$

c. $\frac{5}{4}$

d. $\frac{5}{3}$

23 [Foundational] • Target: Basic trigonometric identities

In a right triangle, the angle θ has an opposite side of length 20, adjacent side 21, and hypotenuse 29. Find $\cos\theta$.

a. $\frac{29}{20}$

b. $\frac{21}{20}$

c. $\frac{21}{29}$

d. $\frac{29}{21}$

24 [Foundational] • Target: Graphs of sine and cosine functions

In a right triangle, the angle θ has an opposite side of length 9, adjacent side 12, and hypotenuse 15. Find $\cos\theta$.

a. $\frac{5}{4}$

b. $\frac{5}{3}$

c. $\frac{4}{5}$

d. $\frac{4}{3}$

25 [Foundational] • Target: Law of Sines

In a right triangle, the angle θ has an opposite side of length 12, adjacent side 16, and hypotenuse 20. Find $\tan\theta$.

a. $\frac{4}{3}$

b. $\frac{3}{4}$

c. $\frac{5}{3}$

d. $\frac{5}{4}$

© ClipSAT — clipsat.org | All Rights Reserved

26 [Foundational] • Target: Law of Cosines

In a right triangle, the angle θ has an opposite side of length 9, adjacent side 12, and hypotenuse 15. Find $\tan\theta$.

a. $\frac{5}{4}$

b. $\frac{5}{3}$

c. $\frac{3}{4}$

d. $\frac{4}{3}$

27 [Foundational] • Target: Trigonometric equations

In a right triangle, the angle θ has an opposite side of length 12, adjacent side 16, and hypotenuse 20. Find $\cos\theta$.

a. $\frac{5}{4}$

b. $\frac{5}{3}$

c. $\frac{4}{5}$

d. $\frac{4}{3}$

28 [Foundational] • Target: Applications of trigonometry to real-world problems

In a right triangle, the angle θ has an opposite side of length 7, adjacent side 24, and hypotenuse 25. Find $\sin\theta$.

a. $\frac{7}{25}$

b. $\frac{25}{7}$

c. $\frac{24}{7}$

d. $\frac{25}{24}$

Part V — Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

29 [Intermediate] • Target: Reciprocal trigonometric functions

Convert 170° to radians, in terms of π.

a. $\frac{18}{17}\pi$

b. $\frac{17}{18}\pi$

c. $\frac{10}{9}\pi$

d. $\frac{17}{9}\pi$

30 [Intermediate] • Target: Angle of elevation and depression

Convert 350° to radians, in terms of π.

a. $\frac{35}{18}\pi$

b. $\frac{18}{35}\pi$

c. $\frac{19}{9}\pi$

d. $\frac{35}{9}\pi$

© ClipSAT — clipsat.org | All Rights Reserved

31 [Intermediate] • Target: Solving trigonometric equations over a restricted domain

Convert 60° to radians, in terms of π.

a. $\frac{2}{3}\pi$

b. $\frac{1}{3}\pi$

c. $\frac{1}{2}\pi$

d. 3π

32 [Intermediate] • Target: A multi-part triangle problem combining Law of Sines and Law of Cosines

Convert 210° to radians, in terms of π.

a. $\frac{4}{3}\pi$

b. $\frac{6}{7}\pi$

c. $\frac{7}{3}\pi$

d. $\frac{7}{6}\pi$

33 [Intermediate] • Target: Cofunction and negative angle identities

Convert 45° to radians, in terms of π.

a. $\frac{5}{12}\pi$

b. $\frac{1}{2}\pi$

c. 4π

d. $\frac{1}{4}\pi$

34 [Intermediate] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7)

Convert 90° to radians, in terms of π.

a. π

b. $\frac{1}{2}\pi$

c. $\frac{2}{3}\pi$

d. 2π

35 [Intermediate] • Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

Convert 15° to radians, in terms of π.

a. 12π

b. $\frac{1}{12}\pi$

c. $\frac{1}{4}\pi$

d. $\frac{1}{6}\pi$

36 [Intermediate] • Target: Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20)

Convert 160° to radians, in terms of π.

a. $\frac{9}{8}\pi$

b. $\frac{16}{9}\pi$

c. $\frac{19}{18}\pi$

d. $\frac{8}{9}\pi$

© ClipSAT — clipsat.org | All Rights Reserved

37 [Intermediate] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

In triangle ABC, $A = 32^\circ$, $B = 57^\circ$, and $a = 9$. Find side b, to the nearest tenth, using the Law of Sines.

a. 14.2

b. 12.2

c. 16.2

d. 18.5

Part VI — Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45)

38 [Challenging] • Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37) • Cross-Lesson: 2.01 Elementary & Intermediate Algebra

In triangle ABC, $A = 48^\circ$, $B = 34^\circ$, and $a = 5$. Find side b, to the nearest tenth, using the Law of Sines.

a. 3.8

b. 1.8

c. 5.8

d. 4.9

39 [Challenging] • Target: Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45) • Cross-Lesson: 3.01 Functions & Graphs

In triangle ABC, $A = 34^\circ$, $B = 36^\circ$, and $a = 17$. Find side b, to the nearest tenth, using the Law of Sines.

a. 17.9

b. 23.2

c. 15.9

d. 19.9

40 [Challenging] • Target: Right triangle trigonometric ratios • Cross-Lesson: 4.01 Coordinate Geometry

In triangle ABC, $a = 10$, $b = 12$, and $C = 105^\circ$. Find side c, to the nearest tenth, using the Law of Cosines.

a. 17.5

b. 21.0

c. 16.0

d. 19.0

© ClipSAT — clipsat.org | All Rights Reserved

41 [Challenging] • Target: The unit circle and radian measure • Cross-Lesson: 5.01 Plane Geometry

In triangle ABC, $a = 6$, $b = 12$, and $C = 30^\circ$. Find side c, to the nearest tenth, using the Law of Cosines.

a. 5.9

b. 7.4

c. 7.9

d. 8.9

42 [Challenging] • Target: Basic trigonometric identities • Cross-Lesson: 7.01 Statistics & Probability

Solve $2\cos\theta - 1 = 0$ for $0^\circ \le \theta < 360^\circ$.

a. $270^\circ, 300^\circ$

b. $30^\circ, 210^\circ$

c. $45^\circ, 315^\circ$

d. $60^\circ, 300^\circ$

43 [Challenging] • Target: Graphs of sine and cosine functions • Cross-Lesson: 8.01 Matrices, Vectors & Complex Numbers

Solve $\tan\theta + 1 = 0$ for $0^\circ \le \theta < 360^\circ$.

a. $135^\circ, 315^\circ$

b. $135^\circ, 240^\circ$

c. $60^\circ, 240^\circ$

d. $90^\circ, 300^\circ$

44 [Challenging] • Target: Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45) • Cross-Lesson: 1.01 Numbers, Percents & Ratios

What is the exact value of $\cos(\frac{5\pi}{6})$?

a. $\frac{\sqrt{3}}{2}$

b. $-\frac{\sqrt{3}}{2}$

c. $\frac{1}{2}$

d. $-\frac{1}{2}$

45 [Challenging] • Target: Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45) • Cross-Lesson: 2.01 Elementary & Intermediate Algebra

In triangle PQR, sides $p = 5$, $q = 8$, and the included angle $R = 60^\circ$. What is the length of side r?

a. 6

b. 7

c. 8

d. $\sqrt{49}$

© ClipSAT — clipsat.org | All Rights Reserved