Vectors and Vector-Valued Functions

ClipSAT Printable Practice Suite (clipsat.org) • Topic 13.01

ANSWER KEY — COMPLETE WORKED SOLUTIONS

Part I — Free-Response Questions (FRQs): Foundational Fluency (1–7)

1 **[Foundational] • Target: 3D Vectors, Dot and Cross Products**

Consider the region R in the first quadrant bounded by the curves $y = \sqrt{x}$, $y = 0$, and $x = 4$. A solid is formed by revolving R about the line $x = 5$. Determine the volume of this solid using the method of cylindrical shells. The volume can be expressed in the form $V = k\pi \int_a^b (5 - x)\sqrt{x}\, dx$. Find the value of k.

The region R is bounded by $y = \sqrt{x}$, $y = 0$, and $x = 4$. We are revolving this region about the vertical line $x = 5$. We will use the method of cylindrical shells. The radius of a cylindrical shell at a given x is the distance from the axis of revolution $x = 5$ to x, which is $5 - x$. The height of the shell is the difference in the y-values of the bounding curves, which is $\sqrt{x} - 0 = \sqrt{x}$. The thickness of the shell is dx. The volume of a single cylindrical shell is.

2 **[Foundational] • Target: Derivatives and Integrals of Vector-Valued Functions**

Consider the vector $\mathbf{v}$ with initial point $P = (1, 2)$ and terminal point $Q = (4, 6)$. What are the components of $\mathbf{v}$?

To find the components of a vector $\mathbf{v}$ with initial point (x_1, y_1) and terminal point (x_2, y_2), we subtract the coordinates of the initial point from the coordinates of the terminal point. That is, $\mathbf{v} = \langle x_2 - x_1, y_2 - y_1 \rangle$. In this case, the initial point is $P = (1, 2)$ and the terminal point is $Q = (4, 6)$. Therefore, the components of $\mathbf{v}$ are $\langle 4 - 1, 6 - 2 \rangle = \langle 3, 4 \rangle$.

3 **[Foundational] • Target: Arc Length and Curvature of Space Curves**

Find the curvature of $\mathbf{r}(t) = \langle 3\cos(t), 3\sin(t), 3t \rangle$ at $t = \pi$.

$\kappa = \frac{||\mathbf{r}' \times \mathbf{r}''||}{||\mathbf{r}'||^3}$. At $t = \pi$: $\mathbf{r}' = \langle 0, -3, 3 \rangle$, $\mathbf{r}'' = \langle 3, 0, 0 \rangle$, giving $\kappa = \frac{1}{6}$.

4 **[Foundational] • Target: Velocity, Acceleration, and Projectile Motion**

The figure shows the graph of the function $f(x) = x^2 - 2x - 1$ and the point $P(3, 2)$ on the curve. What is the slope of the tangent line to the graph of f at point P?

The slope of the tangent line to the graph of $f(x)$ at any point is given by its derivative, $f'(x)$. 1. Find the derivative of $f(x) = x^2 - 2x - 1$ using the power rule: $f'(x) = 2x - 2$ 2. Evaluate the derivative at the x -coordinate of point $P(3, 2)$, which is $x = 3$: $f'(3) = 2(3) - 2 = 6 - 2 = 4$ Thus, the slope of the tangent line at P is 4. Misconception Analysis: - 2 results from mistakenly reading the y-coordinate of point P as the slope, since $f(3) = 2$. - 3 results from taking the x-c.

© ClipSAT — clipsat.org | All Rights Reserved

5 [Foundational] • Target: 3D Vectors, Dot and Cross Products

The graph of the quadratic function $f(x) = x^2 - 6x + k$ is shown. The vertex of the parabola is at the point $(3, -5)$. Find the value of the parameter k.

The vertex of a parabola in the form $f(x) = ax^2 + bx + c$ occurs at $x = -\frac{b}{2a}$. For the given function $f(x) = x^2 - 6x + k$, we have $a = 1$ and $b = -6$. Thus, the x-coordinate of the vertex is $x = -\frac{-6}{2(1)} = 3$. This matches the given vertex $(3, -5)$. Since the vertex $(3, -5)$ lies on the graph of the function, it must satisfy the function's equation. Substitute $x = 3$ and $f(x) = -5$ into the equation: $-5 = (3)^2 - 6(3) + k$ $-5 = 9 - 18 + k$ $-5 = -9 + k$ To solve for k, add 9 to both sides.

6 [Foundational] • Target: Derivatives and Integrals of Vector-Valued Functions

The graph of a differentiable function $f(x)$ is shown below. The function has a tangent line at $x = 1$ with equation $y = 2x + c$. If $f(1) = 3$, find the value of c.

The equation of the tangent line to $f(x)$ at $x = 1$ is given as $y = 2x + c$. We are given that $f(1) = 3$. This means the point $(1, 3)$ lies on both the function $f(x)$ and its tangent line at $x = 1$. Since the point $(1, 3)$ lies on the tangent line $y = 2x + c$, we can substitute the coordinates into the equation: $3 = 2(1) + c$ $3 = 2 + c$ $c = 3 - 2$ $c = 1$ The derivative of the function at $x = 1$ is the slope of the tangent line, which is 2. This information is consistent with the problem but not direct.

7 [Foundational] • Target: Arc Length and Curvature of Space Curves

The region bounded by the curve $y = x^2 - 4$ and the x-axis is rotated about the x-axis. If the volume of the resulting solid is $V = \pi\int_a^b (x^2 - 4)^2 dx$, find the value of $b - a$.

The region is bounded by the curve $y = x^2 - 4$ and the x-axis. To find the limits of integration, we need to find the x-intercepts of the curve, where $y = 0$. Setting $x^2 - 4 = 0$, we get $x^2 = 4$, which gives $x = \pm 2$. Therefore, the bounds of integration are $a = -2$ and $b = 2$. The volume of the solid of revolution is given by $V = \pi\int_a^b [f(x)]^2 dx$. In this case, $f(x) = x^2 - 4$, and the limits are $a = -2$ and $b = 2$. The question asks for the value of $b - a$. $b - a = 2 - (-2) = 2 + 2 = 4$.

Part II — Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

© ClipSAT — clipsat.org | All Rights Reserved

8 [Intermediate] • Target: Velocity, Acceleration, and Projectile Motion

Consider the function $f(x) = x^2 - 6x + 5$. The function has a local minimum at which value of x?

To find the local minimum of the function $f(x) = x^2 - 6x + 5$, we first find the derivative of the function with respect to x: $f'(x) = 2x - 6$. Next, we set the derivative equal to zero to find the critical points: $2x - 6 = 0$. Solving for x, we get $2x = 6$, which means $x = 3$. To determine if this critical point is a local minimum, we can use the second derivative test. The second derivative of $f(x)$ is $f''(x) = 2$. Since $f''(3) = 2 > 0$, the function has a local minimum at $x = 3$.

9 [Intermediate] • Target: 3D Vectors, Dot and Cross Products

Consider the function $f(x) = x^2 - 5x + 6$. What is the derivative of $f(x)$ with respect to x?

To find the derivative of the function $f(x) = x^2 - 5x + 6$, we apply the power rule for differentiation, which states that the derivative of ax^n is anx^{n-1}. The derivative of a constant is 0. Applying the power rule to each term: 1. The derivative of x^2 is $2x^{2-1} = 2x$. 2. The derivative of $-5x$ is $-5 \cdot 1 \cdot x^{1-1} = -5x^0 = -5$. 3. The derivative of the constant 6 is 0. Combining these, the derivative of $f(x)$ is $f'(x) = 2x - 5 + 0 = 2x - 5$.

10 [Intermediate] • Target: Derivatives and Integrals of Vector-Valued Functions

The graph of the function $f(x) = x^2 - 4x + 3$ is shown below. The function has a local minimum. What are the coordinates of the local minimum?

The function is a parabola opening upwards, so its local minimum is also its vertex. The x-coordinate of the vertex of a parabola given by $f(x) = ax^2 + bx + c$ is $x = -b/(2a)$. For $f(x) = x^2 - 4x + 3$, we have $a = 1$ and $b = -4$. Therefore, the x-coordinate of the vertex is $x = -(-4)/(2 * 1) = 4/2 = 2$. To find the y-coordinate, we substitute $x = 2$ into the function: $f(2) = (2)^2 - 4(2) + 3 = 4 - 8 + 3 = -1$. Thus, the coordinates of the local minimum are (2, -1).

11 [Intermediate] • Target: Arc Length and Curvature of Space Curves

Consider the arithmetic sequence with the first term $a_1 = 5$ and a common difference $d = 3$. What is the value of the 5th term, a_5?

The formula for the nth term of an arithmetic sequence is given by $a_n = a_1 + (n - 1)d$, where a_1 is the first term and d is the common difference. Given $a_1 = 5$ and $d = 3$, we want to find the 5th term, a_5. Using the formula: $a_5 = a_1 + (5 - 1)d$ $a_5 = 5 + (4)(3)$ $a_5 = 5 + 12$ $a_5 = 17$ Therefore, the 5th term of the sequence is 17.

© ClipSAT — clipsat.org | All Rights Reserved

12 [Intermediate] • Target: Velocity, Acceleration, and Projectile Motion

Consider the graph of the function $f(x)$ shown below. What is the limit of $f(x)$ as x approaches 1?

The limit of a function as x approaches a certain value is the value that $f(x)$ gets arbitrarily close to as x gets arbitrarily close to that value. Observing the graph, as x approaches 1 from both the left and the right, the value of $f(x)$ approaches 1. Therefore, $\lim_{x \to 1} f(x) = 1$.

13 [Intermediate] • Target: 3D Vectors, Dot and Cross Products

Solve $y' + 2y = x$, $y(0) = -3$. Find $y(1)$.

Integrating factor e^{2x}: $\frac{d}{dx}(e^{2x}y) = xe^{2x}$. Integrating and applying $y(0) = -3$ gives $y = \frac{x}{2} - \frac{1}{4} - \frac{11e^{-2x}}{4}$, so $y(1) = -\frac{2^{\frac{87}{92}} \cdot 3^{\frac{367}{920}} \cdot 5^{\frac{63}{115}} \cdot 7^{\frac{707}{920}}}{210}$.

14 [Intermediate] • Target: Derivatives and Integrals of Vector-Valued Functions

Consider the function $z = f(x, y) = x^2 + 2y^2$. What is the value of the partial derivative $\frac{\partial f}{\partial x}$ at the point $(1, 2)$?

To find the partial derivative of $f(x, y) = x^2 + 2y^2$ with respect to x, we treat y as a constant and differentiate with respect to x. The derivative of x^2 with respect to x is $2x$, and the derivative of $2y^2$ (a constant with respect to x) is 0. Therefore, $\frac{\partial f}{\partial x} = 2x$. Now, we evaluate this partial derivative at the point $(1, 2)$. Substituting $x = 1$ into the expression for $\frac{\partial f}{\partial x}$: $\frac{\partial f}{\partial x}$ at $(1, 2) = 2 \times 1 = 2$. Th.

Part III — Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20)

15 [Challenging] • Target: Arc Length and Curvature of Space Curves • Cross-Lesson: 8.01 Area Between Curves

Consider the region bounded by the curve $y = x^2$, the x-axis, and the vertical lines $x = 1$ and $x = 3$. What is the area of this region?

The area of a region bounded by a curve $y = f(x)$, the x-axis, and the vertical lines $x = a$ and $x = b$ is given by the definite integral $\int_a^b f(x)\,dx$. In this case, $f(x) = x^2$, $a = 1$, and $b = 3$. The area A is given by: $A = \int_1^3 x^2\,dx$ To evaluate this integral, we find the antiderivative of x^2, which is $\frac{x^3}{3}$. Now, we apply the Fundamental Theorem of Calculus.

16 [Challenging] • Target: Velocity, Acceleration, and Projectile Motion • Cross-Lesson: 8.02 Volumes of Revolution

$\frac{d}{dx}\left[\int_{\cos x}^{\sin x} e^{t^2}\,dt\right] =$

Chain rule FTC: $e^{\sin^2 x}\cos x - e^{\cos^2 x} \cdot (-\sin x) = e^{\sin^2 x}\cos x - e^{\cos^2 x}(-\sin x)$... upper bound: $+e^{\sin^2 x}\cos x$; lower bound: $-e^{\cos^2 x}(-\sin x) = -e^{\cos^2 x} \cdot (-\sin x)$. Final: $e^{\sin^2 x}\cos x - e^{\cos^2 x}(-\sin x)$. Correct: $e^{\sin^2 x}\cos x - e^{\cos^2 x}(-\sin x) = e^{\sin^2 x}\cos x + e^{\cos^2 x}\sin x$. Check sign: $\frac{d}{dx}\int_L^U = f(U)U' - f(L)L'$. So $e^{\sin^2 x}\cos x - e^{\cos^2 x}(-\sin x) = e^{\sin^2 x}\cos x + e^{\cos^2 x}\sin x$. Answer is A.

© ClipSAT — clipsat.org | All Rights Reserved

17 [Challenging] • Target: 3D Vectors, Dot and Cross Products • Cross-Lesson: 8.03 Arc Length and Improper Integrals

Consider the function $z = f(x, y) = x^2 - 3xy + 2y^2$. What is the partial derivative of f with respect to y, denoted as $\frac{\partial f}{\partial y}$?

To find the partial derivative of $f(x, y) = x^2 - 3xy + 2y^2$ with respect to y, we treat x as a constant. \\ \\ The derivative of x^2 with respect to y is 0, as x is considered constant. \\ \\ The derivative of $-3xy$ with respect to y is $-3x \cdot 1 = -3x$. \\ \\ The derivative of $2y^2$ with respect to y is $2 \cdot 2y = 4y$. \\ \\ Therefore, the partial derivative $\frac{\partial f}{\partial y}$ is $0 - 3x + 4y = -3x + 4y$.

18 [Challenging] • Target: Derivatives and Integrals of Vector-Valued Functions • Cross-Lesson: 9.01 Separable Differential Equations

A particle's position in the xy-plane is given by the parametric equations $x(t) = 2t + 1$ and $y(t) = t^2 - 3$, where t is time. What is the Cartesian equation of the path traced by the particle?

To find the Cartesian equation, we need to eliminate the parameter t from the given parametric equations. Given: $x(t) = 2t + 1$ $y(t) = t^2 - 3$ From the first equation, we can solve for t in terms of x: $x - 1 = 2t$ $t = \frac{x-1}{2}$ Now, substitute this expression for t into the second equation: $y = \left(\frac{x-1}{2}\right)^2 - 3$ This is the Cartesian equation of the path traced by the particle.

19 [Challenging] • Target: Arc Length and Curvature of Space Curves • Cross-Lesson: 1.01 Precalculus Review for Calculus

A particle's position is described by the parametric equations $x(t) = 3t + 1$ and $y(t) = 2t - 2$. What are the coordinates of the particle when $t = 3$?

To find the coordinates of the particle at a specific time t, substitute the value of t into the parametric equations for x and y. Given $x(t) = 3t + 1$ and $y(t) = 2t - 2$. We need to find the coordinates when $t = 3$. For the x-coordinate: $x(3) = 3(3) + 1 = 9 + 1 = 10$ For the y-coordinate: $y(3) = 2(3) - 2 = 6 - 2 = 4$ Therefore, the coordinates of the particle when $t = 3$ are (10, 4).

20 [Challenging] • Target: Velocity, Acceleration, and Projectile Motion • Cross-Lesson: 10.01 Calculus with Parametric and Polar Curves

Reduction formula: $\int \sin^n x\,dx = -\frac{\sin^{n-1} x \cos x}{n} + \frac{n-1}{n}\int \sin^{n-2} x\,dx$. Then $\int_0^{\pi/2} \sin^6 x\,dx =$

Wallis: $\frac{5}{6} \cdot \frac{3}{4} \cdot \frac{1}{2} \cdot \frac{\pi}{2} = \frac{5\pi \cdot 3}{6 \cdot 4 \cdot 2 \cdot 2} = \frac{15\pi}{96} = \frac{5\pi}{32}$. Actually Wallis: $\int_0^{\pi/2} \sin^{2n} x\,dx = \frac{(2n-1)!!}{(2n)!!}\frac{\pi}{2}$. For $n = 3$: $\frac{5 \cdot 3 \cdot 1}{6 \cdot 4 \cdot 2}\frac{\pi}{2} = \frac{15}{48}\frac{\pi}{2} = \frac{15\pi}{96} = \frac{5\pi}{32}$.

Part IV — Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

© ClipSAT — clipsat.org | All Rights Reserved

21 [Foundational] • Target: 3D Vectors, Dot and Cross Products

Given $\mathbf{u} = \langle 2, -1, 3 \rangle$ and $\mathbf{v} = \langle 4, 2, -1 \rangle$, find $\mathbf{u} \cdot \mathbf{v}$.

a. 3 ✓

b. 7

c. 13

d. -1

$\mathbf{u} \cdot \mathbf{v} = (2)(4) + (-1)(2) + (3)(-1) = 8 - 2 - 3 = 3$.

22 [Foundational] • Target: 3D Vectors, Dot and Cross Products

Find the cross product $\mathbf{u} \times \mathbf{v}$ for $\mathbf{u} = \langle 1, 0, 2 \rangle$ and $\mathbf{v} = \langle 0, 3, -1 \rangle$.

a. $\langle -6, 1, 3 \rangle$ ✓

b. $\langle 6, -1, -3 \rangle$

c. $\langle -6, -1, 3 \rangle$

d. $\langle 6, 1, 3 \rangle$

$\mathbf{u} \times \mathbf{v} = \langle 0(-1) - 2(3), 2(0) - 1(-1), 1(3) - 0(0) \rangle = \langle -6, 1, 3 \rangle$.

23 [Foundational] • Target: 3D Vectors, Dot and Cross Products

Find the area of the parallelogram determined by $\mathbf{u} = \langle 1, 2, 0 \rangle$ and $\mathbf{v} = \langle 0, 3, 4 \rangle$.

a. $\sqrt{89}$ ✓

b. 9

c. $\sqrt{73}$

d. 10

$\mathbf{u} \times \mathbf{v} = \langle 8, -4, 3 \rangle$. Magnitude $||\mathbf{u} \times \mathbf{v}|| = \sqrt{64 + 16 + 9} = \sqrt{89}$.

24 [Foundational] • Target: Derivatives and Integrals of Vector-Valued Functions

Given $\mathbf{r}(t) = \langle t^2, \sin(2t), e^{3t} \rangle$, find $\mathbf{r}'(t)$.

a. $\langle 2t, 2\cos(2t), 3e^{3t} \rangle$ ✓

b. $\langle 2t, \cos(2t), e^{3t} \rangle$

c. $\langle t, 2\cos(2t), 3e^{3t} \rangle$

d. $\langle 2t, -2\cos(2t), 3e^{3t} \rangle$

Differentiate component-wise: $\mathbf{r}'(t) = \langle \frac{d}{dt}(t^2), \frac{d}{dt}(\sin 2t), \frac{d}{dt}(e^{3t}) \rangle = \langle 2t, 2\cos(2t), 3e^{3t} \rangle$.

© ClipSAT — clipsat.org | All Rights Reserved

25 [Foundational] • Target: Derivatives and Integrals of Vector-Valued Functions

Find $\int_0^1 \langle 2t, 3t^2, 4t^3 \rangle \, dt$.

a. $\langle 1, 1, 1 \rangle$ ✓

b. $\langle 2, 3, 4 \rangle$

c. $\langle 1/2, 1/3, 1/4 \rangle$

d. $\langle 1, 2, 3 \rangle$

Integrate component-wise: $\langle [t^2]_0^1, [t^3]_0^1, [t^4]_0^1 \rangle = \langle 1, 1, 1 \rangle$.

26 [Foundational] • Target: Derivatives and Integrals of Vector-Valued Functions

Find the unit tangent vector $\mathbf{T}(t)$ to $\mathbf{r}(t) = \langle 3\cos t, 3\sin t, 4t \rangle$.

a. $\langle -\frac{3}{5}\sin t, \frac{3}{5}\cos t, \frac{4}{5} \rangle$ ✓

b. $\langle -3\sin t, 3\cos t, 4 \rangle$

c. $\langle -\frac{3}{4}\sin t, \frac{3}{4}\cos t, 1 \rangle$

d. $\frac{1}{5}\langle 3\cos t, 3\sin t, 4t \rangle$

$\mathbf{r}'(t) = \langle -3\sin t, 3\cos t, 4 \rangle$. $||\mathbf{r}'(t)|| = \sqrt{9\sin^2 t + 9\cos^2 t + 16} = \sqrt{25} = 5$. Thus $\mathbf{T}(t) = \frac{\mathbf{r}'(t)}{5}$.

27 [Foundational] • Target: Arc Length and Curvature of Space Curves

Find the arc length of the circular helix $\mathbf{r}(t) = \langle 3\cos t, 3\sin t, 4t \rangle$ for $0 \le t \le 2\pi$.

a. 10π ✓

b. 5π

c. 20π

d. 6π

Speed $||\mathbf{r}'(t)|| = \sqrt{(-3\sin t)^2 + (3\cos t)^2 + 4^2} = \sqrt{9 + 16} = 5$. Arc length $L = \int_0^{2\pi} 5 \, dt = 10\pi$.

28 [Foundational] • Target: Arc Length and Curvature of Space Curves

State the formula for curvature $\kappa(t)$ of a space curve in terms of $\mathbf{r}'$ and $\mathbf{r}''$.

a. $\kappa(t) = \frac{||\mathbf{r}'(t) \times \mathbf{r}''(t)||}{||\mathbf{r}'(t)||^3}$ ✓

b. $\kappa(t) = \frac{||\mathbf{r}''(t)||}{||\mathbf{r}'(t)||^2}$

c. $\kappa(t) = \frac{\mathbf{r}'(t) \cdot \mathbf{r}''(t)}{||\mathbf{r}'(t)||^3}$

d. $\kappa(t) = ||\mathbf{r}''(t)||$

The standard vector formula for the curvature of a twice-differentiable space curve is $\kappa = \frac{||\mathbf{r}' \times \mathbf{r}''||}{||\mathbf{r}'||^3}$.

© ClipSAT — clipsat.org | All Rights Reserved

Part V — Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

29 **[Intermediate]** • **Target: Arc Length and Curvature of Space Curves**

Find the curvature of a circle of radius R.

a. $\kappa = 1/R$ ✓

b. $\kappa = R$

c. $\kappa = 2\pi R$

d. $\kappa = 1/R^2$

A circle of radius R has constant curvature equal to the reciprocal of its radius: $\kappa = 1/R$.

30 **[Intermediate]** • **Target: Velocity, Acceleration, and Projectile Motion**

If position is $\mathbf{r}(t) = \langle t^2, 4t, -16t^2 + 64t \rangle$, find the velocity and acceleration vectors at $t = 1$.

a. $\mathbf{v}(1) = \langle 2, 4, 32 \rangle, \mathbf{a}(1) = \langle 2, 0, -32 \rangle$ ✓

b. $\mathbf{v}(1) = \langle 1, 4, 48 \rangle, \mathbf{a}(1) = \langle 2, 0, -16 \rangle$

c. $\mathbf{v}(1) = \langle 2, 4, 0 \rangle, \mathbf{a}(1) = \langle 2, 0, -32 \rangle$

d. $\mathbf{v}(1) = \langle 2, 4, -32 \rangle$

$\mathbf{v}(t) = \mathbf{r}'(t) = \langle 2t, 4, -32t + 64 \rangle$. At $t = 1$: $\mathbf{v}(1) = \langle 2, 4, 32 \rangle$. $\mathbf{a}(t) = \langle 2, 0, -32 \rangle$.

31 **[Intermediate]** • **Target: Velocity, Acceleration, and Projectile Motion**

Find the speed at $t = 0$ for a particle with velocity $\mathbf{v}(t) = \langle 3\cos t, 4, 3\sin t \rangle$.

a. 5 ✓

b. 7

c. 4

d. $\sqrt{7}$

Speed is the magnitude of velocity: $||\mathbf{v}(0)|| = \sqrt{3^2 + 4^2 + 0^2} = \sqrt{9 + 16} = 5$.

© ClipSAT — clipsat.org | All Rights Reserved

32 [Intermediate] • Target: Velocity, Acceleration, and Projectile Motion

Decompose acceleration into tangential and normal components: $\mathbf{a} = a_T\mathbf{T} + a_N\mathbf{N}$. What is a_T?

a. $a_T = \frac{\mathbf{v} \cdot \mathbf{a}}{||\mathbf{v}||}$ ✓

b. $a_T = \frac{||\mathbf{v} \times \mathbf{a}||}{||\mathbf{v}||}$

c. $a_T = ||\mathbf{a}||$

d. $a_T = \frac{\mathbf{a}}{||\mathbf{v}||}$

The tangential component of acceleration represents the rate of change of speed: $a_T = \frac{d}{dt}||\mathbf{v}|| = \frac{\mathbf{v} \cdot \mathbf{a}}{||\mathbf{v}||}$.

33 [Intermediate] • Target: 3D Vectors, Dot and Cross Products

Find the arc length of $\mathbf{r}(t) = \langle 2t, t^2, \ln(t) \rangle$ for $1 \le t \le 3$.

a. $2\ln(3) + 16$

b. $\ln(3) + 8$ ✓

c. $\frac{\ln(3)}{2} + 4$

d. $\ln(3) + 9$

$L = \int_a^b ||\mathbf{r}'(t)||\, dt$; here $||\mathbf{r}'(t)||$ simplifies to a function with an elementary antiderivative, and the integral equals $\ln(3) + 8$.

34 [Intermediate] • Target: Derivatives and Integrals of Vector-Valued Functions

Find the arc length of $\mathbf{r}(t) = \langle 2t, t^2, \ln(t) \rangle$ for $1 \le t \le 2$.

a. $2\ln(2) + 6$

b. $\ln(2) + 3$ ✓

c. $\frac{\ln(2)}{2} + \frac{3}{2}$

d. $\ln(2) + 4$

$L = \int_a^b ||\mathbf{r}'(t)||\, dt$; here $||\mathbf{r}'(t)||$ simplifies to a function with an elementary antiderivative, and the integral equals $\ln(2) + 3$.

© ClipSAT — clipsat.org | All Rights Reserved

35 [Intermediate] • Target: Arc Length and Curvature of Space Curves

Find the arc length of $\mathbf{r}(t) = \langle e^t, e^{-t}, \sqrt{2}\,t\rangle$ for $0 \le t \le \ln(2)$.

a. $\frac{3}{2}$ ✓

b. 3

c. $\frac{3}{4}$

d. $\frac{5}{2}$

$L = \int_a^b ||\mathbf{r}'(t)||\,dt$; here $||\mathbf{r}'(t)||$ simplifies to a function with an elementary antiderivative, and the integral equals $\frac{3}{2}$.

36 [Intermediate] • Target: Velocity, Acceleration, and Projectile Motion

Find the arc length of $\mathbf{r}(t) = \langle e^t, e^{-t}, \sqrt{2}\,t\rangle$ for $0 \le t \le \ln(3)$.

a. $\frac{16}{3}$

b. $\frac{8}{3}$ ✓

c. $\frac{4}{3}$

d. $\frac{11}{3}$

$L = \int_a^b ||\mathbf{r}'(t)||\,dt$; here $||\mathbf{r}'(t)||$ simplifies to a function with an elementary antiderivative, and the integral equals $\frac{8}{3}$.

37 [Intermediate] • Target: 3D Vectors, Dot and Cross Products

Arc length of $\mathbf{r}(t) = (e^t\cos t, e^t\sin t, e^t)$ from $t = 0$ to $t = 1$:

a. $e - 1$

b. $\sqrt{3}(e-1)$ ✓

c. $\sqrt{2}(e-1)$

d. $e\sqrt{3}$

$|r'| = \sqrt{3}e^t \Rightarrow L = \sqrt{3}(e-1)$.

Part VI — Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45)

38 [Challenging] • Target: Derivatives and Integrals of Vector-Valued Functions • Cross-Lesson: 9.01 Separable Differential Equations

Arc length of $\mathbf{r}(t) = (t, t, t)$ from $t = 0$ to $t = 1$:

a. 1

b. $\sqrt{2}$

c. $\sqrt{3}$ ✓

d. 3

$|r'| = \sqrt{3} \Rightarrow L = \sqrt{3}$.

© ClipSAT — clipsat.org | All Rights Reserved

39 [Challenging] • Target: Arc Length and Curvature of Space Curves • Cross-Lesson: 1.01 Precalculus Review for Calculus

Find the curvature of $\mathbf{r}(t) = \langle e^t, e^{-t}, \sqrt{2}\,t \rangle$ at $t = 0$.

a. $\frac{\sqrt{2}}{2}$

b. $\sqrt{2}$

c. $\frac{\sqrt{2}}{4}$ ✓

d. $2\sqrt{2}$

$\kappa = \frac{||\mathbf{r}' \times \mathbf{r}''||}{||\mathbf{r}'||^3}$. At $t = 0$: $\mathbf{r}' = \langle 1, -1, \sqrt{2} \rangle$, $\mathbf{r}'' = \langle 1, 1, 0 \rangle$, giving $\kappa = \frac{\sqrt{2}}{4}$.

40 [Challenging] • Target: Velocity, Acceleration, and Projectile Motion • Cross-Lesson: 10.01 Calculus with Parametric and Polar Curves

Find the curvature of $\mathbf{r}(t) = \langle 3\cos(t), 3\sin(t), 3t \rangle$ at $t = \frac{\pi}{2}$.

a. $\frac{1}{6}$ ✓

b. $\frac{\sqrt{2}}{2}$

c. 3

d. $\frac{1}{3}$

$\kappa = \frac{||\mathbf{r}' \times \mathbf{r}''||}{||\mathbf{r}'||^3}$. At $t = \frac{\pi}{2}$: $\mathbf{r}' = \langle -3, 0, 3 \rangle$, $\mathbf{r}'' = \langle 0, -3, 0 \rangle$, giving $\kappa = \frac{1}{6}$.

41 [Challenging] • Target: 3D Vectors, Dot and Cross Products • Cross-Lesson: 11.01 Sequences and Series

Find the curvature of $\mathbf{r}(t) = \langle 3\cos(t), 3\sin(t), 4t \rangle$ at $t = \pi$.

a. $\frac{3}{25}$ ✓

b. $\frac{3}{5}$

c. 3

d. $\frac{6}{25}$

$\kappa = \frac{||\mathbf{r}' \times \mathbf{r}''||}{||\mathbf{r}'||^3}$. At $t = \pi$: $\mathbf{r}' = \langle 0, -3, 4 \rangle$, $\mathbf{r}'' = \langle 3, 0, 0 \rangle$, giving $\kappa = \frac{3}{25}$.

42 [Challenging] • Target: Derivatives and Integrals of Vector-Valued Functions • Cross-Lesson: 12.01 Convergence Tests

Find the curvature of $\mathbf{r}(t) = \langle \cos(t), \sin(t), 2t \rangle$ at $t = 0$.

a. $\frac{\sqrt{5}}{5}$

b. 1

c. $\frac{2}{5}$

d. $\frac{1}{5}$ ✓

$\kappa = \frac{||\mathbf{r}' \times \mathbf{r}''||}{||\mathbf{r}'||^3}$. At $t = 0$: $\mathbf{r}' = \langle 0, 1, 2 \rangle$, $\mathbf{r}'' = \langle -1, 0, 0 \rangle$, giving $\kappa = \frac{1}{5}$.

© ClipSAT — clipsat.org | All Rights Reserved

43 [Challenging] • Target: Arc Length and Curvature of Space Curves • Cross-Lesson: 12.02 Power Series and Taylor Series

Find the curvature of $\mathbf{r}(t) = (t, t^2, 0)$ at $t = 0$:

a. 2 ✓

b. 1

c. 0

d. 1/2

$\kappa = \frac{|\mathbf{r}' \times \mathbf{r}''|}{|\mathbf{r}'|^3}$; $\mathbf{r}' = (1, 2t, 0)$, $\mathbf{r}'' = (0, 2, 0)$; at $t = 0$: $|(1, 0, 0) \times (0, 2, 0)|/|(1, 0, 0)|^3 = |(0, 0, 2)|/1 = 2$.

44 [Challenging] • Target: Velocity, Acceleration, and Projectile Motion • Cross-Lesson: 14.01 Partial Derivatives and the Gradient

For $\mathbf{r}(t) = \langle 2t, 3t^2, t \rangle$, find the normal component of acceleration at $t = 1$.

a. $\frac{6\sqrt{205}}{41}$ ✓

b. $\frac{36\sqrt{41}}{41}$

c. 6

d. 36

$\mathbf{v} = \langle 2, 6, 1 \rangle$, $\mathbf{a} = \langle 0, 6, 0 \rangle$. $a_N = \frac{||\mathbf{v} \times \mathbf{a}||}{||\mathbf{v}||} = \frac{6\sqrt{205}}{41}$.

45 [Challenging] • Target: 3D Vectors, Dot and Cross Products • Cross-Lesson: 15.01 Double Integrals

For $\mathbf{r}(t) = \langle 2t, 2t^2, t \rangle$, find the normal component of acceleration at $t = 1$.

a. $\frac{16\sqrt{21}}{21}$

b. 4

c. 16

d. $\frac{4\sqrt{105}}{21}$ ✓

$\mathbf{v} = \langle 2, 4, 1 \rangle$, $\mathbf{a} = \langle 0, 4, 0 \rangle$. $a_N = \frac{||\mathbf{v} \times \mathbf{a}||}{||\mathbf{v}||} = \frac{4\sqrt{105}}{21}$.

© ClipSAT — clipsat.org | All Rights Reserved