

The Normal Distribution




Properties of the normal distribution

- 1 State three key properties of a normal distribution's shape.
 - 2 Explain what the two parameters μ and σ control in a normal distribution's shape.
-

The 68-95-99.7 rule

- 3 Adult male heights are approximately normal with mean 70 inches and standard deviation 3 inches. Using the 68-95-99.7 rule, find the range of heights containing the middle 95% of adult men.




- 4 Using the same distribution, find the approximate percentage of adult men taller than 76 inches.
-

Standardizing and using the normal distribution

- 5 Using the same height distribution (mean 70, standard deviation 3), find the z-score for a height of 77.5 inches, and use a standard normal table value of $P(Z < 2.5) \approx 0.9938$ to find the proportion of men shorter than 77.5 inches.
 - 6 Find the proportion of men between 65 and 75 inches tall, given $P(Z < 1.67) \approx 0.9525$ and $P(Z < -1.67) \approx 0.0475$.
-

Visualizing the standard normal distribution


- 7 The graph shows the standard normal distribution ($\mu = 0$, $\sigma = 1$), with the area to the left of $z = 1$ shaded (though shading itself is not shown, imagine it visually).



- a Using $P(Z < 1) \approx 0.8413$, find the area under the curve to the right of $z = 1$.
- b Explain, using the curve's symmetry, why $P(Z < -1) = P(Z > 1)$.

Working backward: finding values from percentiles


- 8 SAT scores are approximately normal with mean 1050 and standard deviation 150. A student scores at the 90th percentile, corresponding to $z \approx 1.28$. Find that student's approximate score.



- 9 Using the same distribution, find the score that separates the bottom 10% of scores, given that $z \approx -1.28$ corresponds to the 10th percentile.


Assessing normality

- 10 Explain how a normal probability plot (or normal quantile plot) can be used to assess whether a data set is approximately normal, and what a roughly straight-line pattern indicates.



Applying the normal distribution to real data

- 11 Pregnancy lengths are approximately normal with mean $\mu = 266$ days and standard deviation $\sigma = 16$ days. Find the z-score for a pregnancy lasting 250 days.



- 12 Using the same pregnancy-length distribution and the 68-95-99.7 rule, find the approximate percentage of pregnancies lasting between 234 and 298 days.

Combining two independent normal random variables

- 13 A vending machine dispenses a soda whose volume is normally distributed with mean 355 mL and standard deviation 4 mL, independently of a snack bag whose weight is normally distributed with mean 50 g and standard deviation 2 g. If T represents the combined "total" formed by adding the soda volume (in mL) and snack weight (in g) as a single number, find the mean of T . (This is a simplified illustrative combination, not a real physical unit.)
- 14 Battery lifespans are normally distributed with mean 500 hours and standard deviation 40 hours. Find the z-score for a battery lasting 460 hours, and describe its relative position.

- 15** A company manufactures resistors with resistance values approximately normal, mean 100 ohms and standard deviation 5 ohms. A resistor is considered defective if its resistance is below 90 ohms or above 110 ohms. Using the 68-95-99.7 rule, estimate the percentage of resistors that are defective, and explain your reasoning step by step.
- 16** Two students are comparing their standardized test scores from different tests. Student X scored 78 on a test with mean 70 and standard deviation 8. Student Y scored 165 on a different test with mean 150 and standard deviation 12. Using z-scores, determine who performed relatively better compared to their own test's distribution of test-takers, showing full reasoning.

